

A CENSUS OF THE VASCULAR PLANTS OF TASMANIA, INCLUDING MACQUARIE ISLAND

MF de Salas & ML Baker

2019 edition

Tasmanian Vascular Plant Census 2019

A Census of the Vascular Plants of Tasmania, including Macquarie Island.

2019 edition

MF de Salas and ML Baker

Postal address:

Tasmanian Herbarium
PO Box 5058
UTAS LPO
Sandy Bay, Tasmania 7005
Australia

Street address:

College Road
Sandy Bay, Tasmania 7005
Australia

© Tasmanian Herbarium, Tasmanian Museum and Art Gallery

Published by the Tasmanian Herbarium, Tasmanian Museum and Art Gallery

GPO Box 1164
Hobart, Tasmania 7001
Australia

<https://www.tmag.tas.gov.au>

Cite as: de Salas, MF, Baker, ML (2019) *A Census of the Vascular Plants of Tasmania, including Macquarie Island*. (Tasmanian Herbarium, Tasmanian Museum and Art Gallery, Hobart) <https://flora.tmag.tas.gov.au/resources/census/>

Introduction

The *Census of the Vascular Plants of Tasmania* is a checklist of every native and naturalised vascular plant taxon for which there is physical evidence of its presence in Tasmania. It includes the correct nomenclature and authorship of the taxon's name, as well as the reference of its original publication.

According to this *Census*, the Tasmanian flora contains 2726 vascular plants, of which 1920 (70%) are considered native and 808 (30%) have naturalised from elsewhere. Among the native taxa, 533 (28%) are endemic to the State. Forty-eight of the State's exotic taxa are considered sparingly naturalised, and are known only from a small number of populations. Twenty-three native taxa are recognised as extinct, whereas eight naturalised taxa are considered to have either not persisted in Tasmania or have been eradicated. The sub-Antarctic Macquarie Island, considered part of Tasmania, supports 49 species of vascular plants, of which 42 are considered native and seven naturalised. For some basic statistics on the Tasmanian flora see Tables 1–3.

No new native species are recognised in the 2019 edition (Appendix 1a). The names of several taxa have changed since the previous edition (Appendix 1b), some of which now are recognised as infraspecies. Finally, one taxon (*Isolepis alpina*) has had its status changed since the previous edition (Appendix 1c). *Viola sieberiana* is no longer considered part of the Tasmanian flora and has been removed from this edition of the *Census*.

The *Census* incorporates every name (including synonyms) used to refer to Tasmanian plants in the major taxonomic publications about Tasmania, and aims to account for every species name attributed to Tasmania, whether erroneously or correctly. Names from the following publications have been included: Jacques-Julien Houtou de Labillardière's *Novae Hollandiae Plantarum Specimen* (1804-1807), Robert Brown's *Prodromus Florae Novae Hollandiae et Insulae van-Diemen* (1810), Joseph Dalton Hooker's *The Botany of the Antarctic Voyage of H. M. Discovery Ships Erebus and Terror in the Years 1839 - 1843 Under the Command of Captain Sir James Clark Ross. Part III: Flora Tasmaniae* (1855-1859, complete work published in 1860), all volumes of George Bentham's *Flora Australiensis* (1863-1878) and Leonard Rodway's *The Tasmanian Flora* (1903). In addition, all names used in Cheeseman's (1919) *The Vascular Flora of Macquarie Island* have been accounted for in the Macquarie Island section.

The classification systems used in this *Census* follows APG IV (2016) for flowering plants (Angiosperms) and McCarthy (1998) for conifers, ferns and their allies. In contrast, the system used to arrange the botanical collections of the Tasmanian Herbarium and in the *Flora of Australia* series, which is published by the Australian Biological Resources Study (ABRS), follows Cronquist (1981), and the *Flora of Tasmania Online* (Duretto 2009+) follows the older APG II (2003) system. To determine the families in which genera are placed in this edition, please refer to Appendix 2 at the end of this document.

This census also serves as an index to *The Student's Flora of Tasmania* (Curtis 1963, 1967, 1979; Curtis & Morris 1975, 1994). Species accounts can be found in *The Student's Flora of Tasmania* by referring to the volume and page number reference that is given in the far right column (e.g. 3:539). Families with accounts completed in the *Flora of Tasmania Online* (de Salas 2009+) are highlighted at the family entry (e.g. FTO 1).

Acknowledgments

This checklist was developed using BRAHMS (Botanical Research And Herbarium Management System) database management system. The authors would like to thank the BRAHMS team for making possible (and infinitely easier) the publication of an annual up-to-date census. In addition the authors would like to thank the APC (Australian Plant Census) group, whose work brings to our attention a myriad of taxonomic updates we might otherwise miss. We would like to thank Lynette Cave of the Tasmanian Herbarium for her assistance in checking the validity and publication details of numerous taxa. Mark Wapstra (ECOtas) provided useful comments on a draft of this publication and for these we are grateful.

TABLE 1: SOME BASIC STATISTICS ON THE STATUS OF TASMANIAN VASCULAR PLANTS

| | | Native | | Naturalised |
|-------------------|-------------|-------------|------------|-------------|
| | | non-endemic | endemic | |
| Basal angiosperms | 2 | 1 | 1 | 0 |
| Magnoliids | 8 | 7 | 1 | 0 |
| Eudicots | 1736 | 792 | 363 | 581 |
| Monocotyledons | 862 | 490 | 151 | 221 |
| Gymnosperms | 15 | 2 | 9 | 4 |
| Pteridophytes | 105 | 95 | 8 | 2 |
| TOTAL | 2728 | 1387 | 533 | 808 |

TABLE 2: SOME BASIC STATISTICS ON THE STATUS OF MACQUARIE ISLAND VASCULAR PLANTS

| | | Native | | Naturalised |
|-------------------|-----------|-------------|----------|-------------|
| | | non-endemic | endemic | |
| Basal angiosperms | 0 | 0 | 0 | 0 |
| Magnoliids | 0 | 0 | 0 | 0 |
| Eudicots | 24 | 20 | 1 | 3 |
| Monocotyledons | 20 | 13 | 3 | 4 |
| Gymnosperms | 0 | 0 | 0 | 0 |
| Pteridophytes | 5 | 5 | 0 | 0 |
| Total | 49 | 38 | 4 | 7 |

TABLE 3: STATISTICS OF TASMANIAN AND MACQUARIE ISLAND VASCULAR PLANT FAMILIES AND GENERA

| | Tasmanian families | Tasmanian genera | Macquarie Island families | Macquarie Island genera |
|-------------------|--------------------|------------------|---------------------------|-------------------------|
| Basal angiosperms | 1 | 1 | 0 | 0 |
| Magnoliids | 4 | 4 | 0 | 0 |
| Eudicots | 99 | 551 | 14 | 19 |
| Monocotyledons | 38 | 242 | 4 | 12 |
| Gymnosperms | 3 | 10 | 0 | 0 |
| Pteridophytes | 26 | 43 | 5 | 5 |
| Total | 171 | 851 | 23 | 36 |

Symbols used in species list

- e endemic in Tasmania
- t within Australia, occurs only in Tasmania
- x considered by the Tasmanian Herbarium to be extinct or eradicated in Tasmania or Macquarie Island
- i introduced and naturalised in Tasmania
- # sparingly naturalised or known from only one or two populations or collections
- * doubtfully naturalised: collections exist but no evidence exists to establish if a taxon is naturalised in Tasmania
- n a change since the 2018 edition (see Appendix I for summary)

no symbol indicates that the taxon is native both in Tasmania and the Australian mainland

NOTE: The 2019 *Census* no longer includes information on the listing status of species listed on the *Tasmanian Threatened Species Protection Act 1995*, the *Commonwealth Environment Protection and Biodiversity Conservation Act 1999*, or the *Tasmanian Weed Management Act 1999*. For this information please refer to current legislation.

TABLE OF CONTENTS

A CENSUS OF THE VASCULAR PLANTS OF TASMANIA, INCLUDING MACQUARIE ISLAND..... 1

Introduction..... 3

Acknowledgments..... 4

Table 1: Some basic statistics on the status of
Tasmanian vascular plants..... 4

Table 2: Some basic statistics on the status of
Macquarie Island vascular plants 4

Table 3: Statistics of Tasmanian and Macquarie
Island vascular plant families and genera..... 4

Symbols used in species list..... 5

TABLE OF CONTENTS 6

Tasmanian Vascular Plant Census 9

Basal angiosperms 9

Hydatellaceae FTO 1 9

Magnoliids 9

Atherospermataceae FTO 3 9

Lauraceae FTO 5 9

Monimiaceae FTO 4 9

Winteraceae FTO 2 9

Eudicots..... 9

Acanthaceae FTO 120..... 9

Adoxaceae 9

Aizoaceae FTO 101..... 9

Amaranthaceae FTO 98..... 10

Apiaceae (UMBELLIFERAE)..... 12

Apocynaceae 13

Aquifoliaceae 13

Araliaceae..... 13

Asclepidaceae..... 14

Asteraceae (COMPOSITAE) 14

Basellaceae FTO 99..... 29

Berberidaceae FTO 46 29

Betulaceae 30

Bignoniaceae FTO 121 30

Boraginaceae 30

Brassicaceae (CRUCIFERAE)..... 31

Campanulaceae..... 33

Cannabaceae 34

Caprifoliaceae 34

Caryophyllaceae 34

Casuarinaceae FTO 67 37

Celastraceae..... 37

Cistaceae 37

Convolvulaceae FTO 111 37

Crassulaceae 38

Cucurbitaceae FTO 65 39

Cunoniaceae FTO 72 39

Dilleniaceae FTO 92 39

Dipsacaceae FTO 134..... 40

Donatiaceae FTO 126..... 40

Droseraceae FTO 96 40

Elaeocarpaceae FTO 71 41

Elatinaceae FTO 76 41

Ericaceae FTO 105 41

Escalloniaceae FTO 131 47

Euphorbiaceae 47

Fabaceae (LEGUMINOSAE) 48

Frankeniaceae FTO 93..... 55

Fumariaceae FTO 45 55

Gentianaceae 55

Geraniaceae FTO 54 56

Goodeniaceae FTO 129..... 57

Griselinaceae FTO 136..... 57

Grossulariaceae FTO 50 57

Gunneraceae FTO 49 57

Gyrostemonaceae FTO 83 57

Haloragaceae 58

Hypericaceae 59

Lamiaceae (LABIATAE) 59

Lentibulariaceae FTO 119..... 60

Linaceae FTO 77 60

Loganiaceae..... 60

Lythraceae FTO 55 61

Malvaceae 61

Mazaceae 61

Menyanthaceae FTO 128 61

Myrsinaceae 62

Myrtaceae FTO 57 62

Nothofagaceae 65

Nyctaginaceae FTO 101.1 65

Oleaceae FTO 113 65

Onagraceae 65

Orobanchaceae 66

Oxalidaceae FTO 70 67

Papaveraceae FTO 44 68

Passifloraceae FTO 81 68

| | | | |
|------------------------------|-----------|---------------------------|------------|
| Phrymaceae | 68 | Burmanniaceae FTO 16..... | 89 |
| Phyllanthaceae..... | 68 | Campynemataceae | 89 |
| Phytolaccaceae..... | 68 | Centrolepidaceae | 89 |
| Picrodendraceae..... | 68 | Colchicaceae..... | 90 |
| Pittosporaceae FTO 138..... | 69 | Commelinaceae | 90 |
| Plantaginaceae | 69 | Cymodoceaceae | 90 |
| Plumbaginaceae FTO 94..... | 71 | Cyperaceae | 90 |
| Polemoniaceae FTO 102..... | 71 | Haemodoraceae | 95 |
| Polygalaceae FTO 60..... | 71 | Hemerocallidaceae..... | 95 |
| Polygonaceae FTO 95..... | 72 | Hyacinthaceae | 96 |
| Portulacaceae FTO 100..... | 73 | Hydrocharitaceae..... | 96 |
| Proteaceae FTO 48..... | 73 | Hypoxidaceae..... | 96 |
| Ranunculaceae FTO 47 | 75 | Iridaceae | 97 |
| Resedaceae FTO 84..... | 77 | Juncaceae | 98 |
| Rhamnaceae..... | 77 | Juncaginaceae..... | 99 |
| Rosaceae | 78 | Laxmanniaceae | 100 |
| Rubiaceae | 80 | Luzuriagaceae | 100 |
| Rutaceae FTO 87..... | 81 | Melanthiaceae..... | 100 |
| Salicaceae FTO 80..... | 83 | Orchidaceae | 100 |
| Santalaceae FTO 91 | 83 | Poaceae (GRAMINEAE) | 110 |
| Sapindaceae FTO 86..... | 83 | Posidoniaceae..... | 125 |
| Scrophulariaceae | 83 | Potamogetonaceae..... | 125 |
| Solanaceae FTO 112..... | 84 | Restionaceae..... | 126 |
| Stylidiaceae FTO 127..... | 84 | Ruppiaceae | 127 |
| Tetracarpaeaceae FTO 52..... | 85 | Typhaceae | 127 |
| Theophrastaceae FTO 103..... | 85 | Xanthorrhoeaceae | 127 |
| Thymelaeaceae FTO 89 | 85 | Xyridaceae | 127 |
| Tropaeolaceae FTO 82 | 86 | Zosteraceae..... | 127 |
| Ulmaceae FTO 63..... | 86 | Gymnosperms | 127 |
| Urticaceae FTO 64..... | 86 | Cupressaceae | 127 |
| Valerianaceae FTO 135..... | 86 | Pinaceae | 128 |
| Verbenaceae..... | 86 | Podocarpaceae..... | 128 |
| Violaceae FTO 79..... | 86 | Pteridophytes..... | 128 |
| Zygophyllaceae FTO 58 | 87 | Adiantaceae..... | 128 |
| Monocots..... | 87 | Aspleniaceae | 128 |
| Agapanthaceae | 87 | Athyriaceae | 129 |
| Alismataceae..... | 87 | Azollaceae | 129 |
| Alliaceae | 87 | Blechnaceae | 129 |
| Alstroemeriaceae..... | 87 | Culcitaceae..... | 130 |
| Amaryllidaceae..... | 88 | Cyatheaceae | 130 |
| Aponogetonaceae | 88 | Dennstaedtiaceae | 130 |
| Araceae..... | 88 | Dicksoniaceae | 131 |
| Asparagaceae..... | 88 | Dryopteridaceae..... | 131 |
| Asphodelaceae..... | 88 | Equisetaceae | 131 |
| Asteliaceae..... | 88 | Gleicheniaceae | 131 |
| Blandfordiaceae | 89 | Grammitidaceae..... | 132 |

| | | | |
|---|------------|--|------------|
| Hemionitidiaceae..... | 132 | Ranunculaceae FTO 47..... | 137 |
| Hymenophyllaceae..... | 132 | Rosaceae..... | 137 |
| Isoetaceae | 133 | Rubiaceae | 137 |
| Lindsaeaceae..... | 133 | Monocots | 137 |
| Lycopodiaceae | 133 | Cyperaceae | 137 |
| Marsileaceae | 134 | Juncaceae | 137 |
| Ophioglossaceae | 134 | Orchidaceae | 137 |
| Osmundaceae | 134 | Poaceae (GRAMINEAE)..... | 137 |
| Polypodiaceae | 134 | Pteridophytes..... | 138 |
| Psilotaceae | 134 | Blechnaceae | 138 |
| Pteridaceae | 135 | Dryopteridaceae..... | 138 |
| Schizaeaceae | 135 | Gleicheniaceae | 138 |
| Selaginellaceae..... | 135 | Grammitidaceae..... | 138 |
| Thelypteridaceae..... | 135 | Hymenophyllaceae | 138 |
| Macquarie Island Vascular Plant Census | 136 | Lycopodiaceae..... | 138 |
| Eudicots | 136 | APPENDIX 1: CHANGES IN THE CENSUS | |
| Apiaceae (UMBELLIFERAE)..... | 136 | BETWEEN THE 2018 AND 2019 EDITIONS..... | 139 |
| Araliaceae..... | 136 | Ia: Taxa new to the Tasmanian Census..... | 139 |
| Asteraceae (COMPOSITAE) | 136 | Ib: Taxa that have changed names between 2018 | |
| Brassicaceae (CRUCIFERAE)..... | 136 | and 2019..... | 139 |
| Caryophyllaceae | 136 | Ic: Taxa that have changed status between 2018 | |
| Crassulaceae..... | 136 | and 2019..... | 140 |
| Haloragaceae..... | 136 | Id: Taxa now not considered to be part of the | |
| Onagraceae..... | 136 | Tasmanian Flora..... | 140 |
| Plantaginaceae | 137 | APPENDIX 2. FAMILY PLACEMENT OF | |
| Polygonaceae FTO 95..... | 137 | GENERA IN THIS CENSUS..... | 141 |
| Portulacaceae FTO 100..... | 137 | | |

Tasmanian Vascular Plant Census

BASAL ANGIOSPERMS

HYDATELLACEAE

FTO 1

- Hydatella filamentosa (Rodway) W.M.Curtis = **Trithuria filamentosa** 4b:361
 e **Trithuria filamentosa** Rodway, Pap. & Proc. Roy. Soc. Tasmania 1897: 48 (1898) 4b:361
Trithuria submersa Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 79 (1858) 4b:360

MAGNOLIIDS

ATHEROSPERMATACEAE

FTO 3

- Atherosperma moschatum** Labill. subsp. **moschatum**, Nov. Holl. Pl. 2: 74, t.224 (1806) 3:594

LAURACEAE

FTO 5

- Cassytha glabella** R.Br. f. **dispar** (Schltdl.) J.Z.Weber, J. Adelaide Bot. Gard. 3: 209 (1981) 3:597
Cassytha glabella R.Br. f. **glabella**, Prodr. Fl. Nov. Holland. 404 (1810) 3:597
Cassytha melantha R.Br., Prodr. Fl. Nov. Holland. 404 (1810) 3:597
 e **Cassytha pedicellosa** J.Z.Weber, J. Adelaide Bot. Gard. 3: 214 (1981)
Cassytha pubescens R.Br., Prodr. Fl. Nov. Holland. 404 (1810) 3:597

MONIMIACEAE

FTO 4

- Hedycarya angustifolia** A.Cunn., Ann. Nat. Hist. Ser. I 1: 215 (1838) 3:595

WINTERACEAE

FTO 2

- Drimys aromatica** (R.Br. ex DC.) F.Muell. = **Tasmannia lanceolata**
Drimys lanceolata (Poir.) Baill. = **Tasmannia lanceolata** 1:25
Tasmannia aromatica R.Br. ex DC. = **Tasmannia lanceolata**
Tasmannia lanceolata (Poir.) A.C.Sm., Taxon 18: 287 (1969) 1:25

EUDICOTS

ACANTHACEAE

FTO 120

- i **Acanthus mollis** L., Sp. Pl. 2: 639 (1753) 3:539

ADOXACEAE

- Sambucus gaudichaudiana** DC., Prodr. [A. P. de Candolle] 4: 322 (1830) 2:265
 i **Sambucus nigra** L., Sp. Pl. 1: 269 (1753) 2:265
 i # **Viburnum tinus** L., Sp. Pl. 1: 267 (1753)

AIZOACEAE

FTO 101

- Aptenia cordifolia** (L.f.) Schwantes = **Mesembryanthemum cordifolium**
 i # **Carpobrotus aequilaterus** (Haw.) N.E.Br., J. Bot. 66: 324 (1928)
 n i **Carpobrotus edulis** (L.) N.E.Br. subsp. **edulis**, Gen. S. Afr. Fl. Pl. 249 (1926) 2:238
Carpobrotus rossii (Haw.) Schwantes, Gartenflora 77: 68 (1928) 2:238

| | | |
|-----|--|---------------|
| | Disphyma australe (Aiton) N.E.Br. sensu Curtis (1963) = Disphyma crassifolium subsp. clavellatum (misapplied in Tasmania) | 2:239 |
| | Disphyma crassifolium (L.) L.Bolus subsp. clavellatum (Haw.) Chinnock, Fl. S. Austral., ed. 4, 1: 194 (1986) | 2:239 |
| i | Drosanthemum candens (Haw.) Schwantes, Z. Sukkulantenk. 3: 29 (1927) | |
| i | Galenia pubescens (Eckl. & Zeyh.) Druce var. pubescens , Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 624 (1917) | 2:241 |
| | Galenia secunda (L.f.) Sond. sensu Curtis (1963) = Galenia pubescens var. pubescens (misapplied in Tasmania) | 2:241 |
| i | Lampranthus glaucus (L.) N.E.Br., Gard. Chron. 87: 212 (1930) | 2:240 |
| | Mesembryanthemum aequilaterale Haw. sensu Rodway (1903) = Carpobrotus rossii (misapplied in Tasmania) | 2:238 |
| | Mesembryanthemum australe Sol. sensu Rodway (1903) = Disphyma crassifolium subsp. clavellatum (misapplied in Tasmania) | 2:239 |
| i # | Mesembryanthemum cordifolium L.f., Suppl. Pl. 260 (1782) | |
| i | Mesembryanthemum crystallinum L., Sp. Pl. 1: 480 (1753) | |
| | Tetragonia expansa Murray = Tetragonia tetragonoides | 2:240 |
| | Tetragonia implexicoma (Miq.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 148 (1856) | 2:241 |
| | Tetragonia tetragonoides (Pall.) Kuntze, Revis. Gen. Pl. 1: 264 (1891) | 2:240 |
| n | Zaleyia galericulata (Melville) H.Eichler sensu Walsh (1996) attributed to Tasmania in error | |
| | AMARANTHACEAE | FTO 98 |
| | Alternanthera denticulata R.Br., Prodr. Fl. Nov. Holland. 417 (1810) | 3:568 |
| | Alternanthera sessilis (L.) R.Br. ex DC. sensu Hooker (1860) = Alternanthera denticulata (misapplied in Tasmania) | |
| i | Amaranthus albus L., Syst. Nat., ed. 10: 1268 (1759) | |
| i | Amaranthus deflexus L., Mant. Pl. 2: 295 (1771) | 3:567 |
| | Amaranthus graecizans L. subsp. silvestris (Vill.) Brenan previously listed as naturalised but insufficient evidence exists to support this | |
| | Amaranthus hybridus L. sensu Curtis (1967) = Amaranthus powellii (misapplied in Tasmania) | 3:566 |
| i | Amaranthus powellii S.Watson, Proc. Amer. Acad. Arts. 10: 347 (1875) | 3:566 |
| | Amaranthus retroflexus L. sensu Curtis (1967) = Amaranthus powellii (misapplied in Tasmania) | 3:566 |
| | Amaranthus spinosus L. previously listed as naturalised but insufficient evidence exists to support this | |
| | Arthrocnemum arbuscula (R.Br.) Moq. = Tecticornia arbuscula | 3:577 |
| ?i | Atriplex australasica Moq., Chenop. Monogr. Enum. 59 (1840) | |
| t | Atriplex billardiieri (Moq.) Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 215 (1853) | 3:575 |
| | Atriplex cinerea Poir., Encycl. (Lamarck) Suppl. 1: 471 (1811) | 3:574 |
| | Atriplex halimus L. sensu Brown (1810) = Atriplex cinerea (misapplied in Tasmania) | |
| | Atriplex hastata L. sensu Curtis (1967) = Atriplex prostrata (misapplied in Tasmania) | 3:575 |
| i | Atriplex hortensis L., Sp. Pl. 2: 1053 (1753) | 3:576 |
| | Atriplex paludosa R.Br. subsp. paludosa , Prodr. Fl. Nov. Holland. 406 (1810) | 3:574 |
| i | Atriplex patula L., Sp. Pl. 2: 1053 (1753) | 3:575 |
| | Atriplex patula L. var. angustifolia Rodway a name of uncertain application | |
| | Atriplex patula L. var. littoralis Rodway a name of uncertain application | |
| i | Atriplex prostrata Boucher ex DC., Fl. Franc. (DC. & Lamarck), ed. 3, 3: 387 (1805) | 3:575 |

| | | | |
|---|-----|--|-------|
| ? | i # | Atriplex semibaccata R.Br., Prodr. Fl. Nov. Holland. 406 (1810) | |
| | | Atriplex suberecta I.Verd., Bothalia 6: 418 (1954) | |
| | i * | Bassia scoparia (L.) A.J.Scott, Feddes Repert. 89: 108 (1978) | |
| | i | Beta vulgaris L. subsp. maritima (L.) Arcang., Comp. Fl. Ital. [Arcangeli]: 593 (1882) | 3:573 |
| | i | Chenopodium album L., Sp. Pl. 1: 219 (1753) | 3:571 |
| | | Chenopodium ambiguum R.Br. = Chenopodium glaucum | |
| | | Chenopodium baccatum Labill. sensu Labillardiere (1806) = Rhagodia candolleana (misapplied in Tasmania) | |
| | | Chenopodium capitatum (L.) Ambrosi sensu Buchanan (2009) = Chenopodium foliosum (misapplied in Tasmania) | |
| | x | Chenopodium erosum R.Br., Prodr. Fl. Nov. Holland. 407 (1810) | |
| | | Chenopodium foliosum (Moench) Asch. previously listed as naturalised but insufficient evidence exists to support this | |
| | | Chenopodium furfuraceum Moq. = Rhagodia candolleana subsp. candolleana | |
| | ? | Chenopodium glaucum L., Sp. Pl. 1: 220 (1753) | 3:572 |
| | | Chenopodium glaucum L. subsp. ambiguum (R.Br.) Murr & Thell. ex Thell. = Chenopodium glaucum | |
| | i | Chenopodium murale L., Sp. Pl. 1: 219 (1753) | 3:572 |
| | | Chenopodium pumilio R.Br. = Dysphania pumilio | 3:573 |
| | i | Chenopodium vulvaria L., Sp. Pl. 1: 220 (1753) | |
| | i | Dysphania glomulifera (Nees) Paul G.Wilson subsp. glomulifera , Nuytsia 4: 183 (1983) | |
| | i | Dysphania pumilio (R.Br.) Mosyakin & Clemants, Ukrayins'k. Bot. Zhurn. 59: 382 (2002) | 3:573 |
| | | Einadia nutans (R.Br.) A.J.Scott subsp. nutans , Feddes Repert. 89: 3 (1978) | 3:570 |
| | | Hemichroa pentandra R.Br., Prodr. Fl. Nov. Holland. 409 (1810) | 3:576 |
| | | Ptilotus spathulatus (R.Br.) Poir., Encycl. (Lamarck) Suppl. 4: 620 (1816) | 3:567 |
| | | Rhagodia baccata (Labill.) Moq. sensu Curtis (1967) = Rhagodia candolleana (misapplied in Tasmania) | 3:570 |
| | | Rhagodia billardierei R.Br. sensu Brown (1810) = Rhagodia candolleana (misapplied in Tasmania) | |
| | | Rhagodia billardierei R.Br. var. congesta (Hook.f.) Benth. = Chenopodium murale | |
| | | Rhagodia candolleana Moq. subsp. candolleana , Chenop. Monogr. Enum. 10 (1840) | 3:570 |
| | | Rhagodia nutans R.Br. = Einadia nutans | 3:570 |
| | | Salicornia arbuscula R.Br. = Tecticornia arbuscula | |
| | | Salicornia australis Sol. ex Benth. nom. illeg. = Sarcocornia quinqueflora subsp. quinqueflora | |
| | | Salicornia blackiana Ulbr. = Sarcocornia blackiana | 3:579 |
| | | Salicornia indica R.Br. sensu Brown (1810) = Sarcocornia quinqueflora (misapplied in Tasmania) | |
| | | Salicornia quinqueflora Bunge ex Ung.-Sternb. = Sarcocornia quinqueflora | 3:578 |
| | | Salsola australis R.Br., Prodr. Fl. Nov. Holland. 411 (1810) | |
| | | Salsola kali L. sensu Buchanan (2005) = Salsola australis (misapplied in Tasmania) | |
| | | Salsola tragus L. sensu Buchanan (2009) = Salsola australis (misapplied in Tasmania) | |
| | | Sarcocornia blackiana (Ulbr.) A.J.Scott, Bot. J. Linn. Soc. 75: 369 (1978) | 3:579 |
| | | Sarcocornia quinqueflora (Bunge ex Ung.-Sternb.) A.J.Scott subsp. quinqueflora , Bot. J. Linn. Soc. 75: 368 (1978) | 3:578 |
| | | Sarcocornia quinqueflora (Bunge ex Ung.-Sternb.) A.J.Scott subsp. tasmanica Paul G.Wilson, Nuytsia 3: 74 (1980) | 3:578 |
| | | Sclerostegia arbuscula (R.Br.) Paul G.Wilson = Tecticornia arbuscula | 3:577 |

| | | |
|--------------------------------|--|-------|
| | Suaeda australis (R.Br.) Moq., Ann. Sci. Nat. (Paris) 23: 318 (1831) | 3:580 |
| i | Suaeda maritima (L.) Dumort. subsp. maritima , Fl. Belg. (Dumortier): 22 (1827) | |
| | Tecticornia arbuscula (R.Br.) K.A.Sheph. & Paul G.Wilson, Austral. Syst. Bot. 20: 325 (2007) | 3:577 |
| | Threlkeldia diffusa R.Br., Prodr. Fl. Nov. Holland. 410 (1810) | 3:577 |
| | Trichinium spathulatum R.Br. = Ptilotus spathulatus | |
| APIACEAE (UMBELLIFERAE) | | |
| | Aciphylla procumbens (F.Muell.) Benth. = Anisotome procumbens | 2:261 |
| | Actinotus bellidioides (Hook.f.) Benth., Fl. Austral. 3: 369 (1867) | 2:253 |
| e | Actinotus moorei F.Muell. ex Rodway, Pap. & Proc. Roy. Soc. Tasmania 1894: 52 (1896) | 2:254 |
| e | Actinotus suffocatus (Hook.f.) Rodway, Pap. & Proc. Roy. Soc. Tasmania 1893: 180 (1894) | 2:254 |
| | Aegopodium podagraria L. previously listed as naturalised but insufficient evidence exists to support this | |
| i | Ammi majus L., Sp. Pl. 1: 243 (1753) | |
| e | Anisotome procumbens (F.Muell.) C.J.Webb, Fl. & Fauna Alpine Australasia: 395 (1986) | 2:261 |
| i | Anthriscus caucalis M.Bieb., Fl. Taur.-Caucas. 1: 230 (1808) | |
| | Apium annuum P.S.Short, J. Adelaide Bot. Gard. 1: 230 (1979) | |
| | Apium australe Thouars sensu Bentham (1867) misapplied to A. annuum , A. insulare & A. prostratum | |
| | Apium insulare P.S.Short, J. Adelaide Bot. Gard. 1: 228 (1979) | |
| | Apium prostratum Labill. ex Vent. subsp. prostratum var. filiforme (A.Rich.) Kirk, Stud. Fl. New Zealand 196 (1899) | 2:255 |
| | Apium prostratum Labill. ex Vent. subsp. prostratum var. prostratum , Jard. Malmaison 2: t.81 (1805) | 2:255 |
| | Azorella dichopetala Benth. nom. illeg. = Dichosciadium ranunculaceum | |
| | Azorella saxifraga (Hook.f.) Benth. = Oschatzia saxifraga | 2:251 |
| | Caldasia brachycarpa Hook.f. = Oreomyrrhis eriopoda | |
| | Caucalis infesta Curtis = Torilis arvensis | |
| | Centella cordifolia (Hook.f.) Nannf., Svensk Bot. Tidskr. 18: 418 (1924) | 2:247 |
| i | Conium maculatum L., Sp. Pl. 1: 243 (1753) | 2:254 |
| | Crantzia lineata Nutt. sensu Bentham (1867) = Lilaeopsis polyantha (misapplied in Tasmania) | |
| i | Cyclosporum leptophyllum (Pers.) Sprague ex Britton & P.Wilson, Bot. Porto Rico 6: 52 (1925) | |
| | Daucus brachiatus Sieber ex DC. = Daucus glochidiatus | |
| i | Daucus carota L., Sp. Pl. 1: 242 (1753) | 2:262 |
| | Daucus glochidiatus (Labill.) Fisch., C.A.Mey. & Avé-Lall., Index Seminum [St. Petersburg] 9, Suppl. 11 (1844) | 2:262 |
| | Daucus pusillus Michx. sensu Hooker (1860) = Daucus glochidiatus (misapplied in Tasmania) | |
| e | Dichosciadium ranunculaceum (F.Muell.) Domin var. tasmanicum (Hook.f.) Domin, Repert. Spec. Nov. Regni Veg. 5: 105 (1908) | 2:250 |
| e | Diplaspis cordifolia (Hook.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 157 (1856) | 2:252 |
| e | Diplaspis hydrocotyle Hook.f., London J. Bot. 6: 469 bis (1847) | 2:251 |
| | Eryngium ovinum A.Cunn., Geogr. Mem. New South Wales [Field] 358 (1825) | 2:252 |
| | Eryngium rostratum Cav. sensu Curtis (1963) = Eryngium ovinum (misapplied in Tasmania) | 2:252 |
| | Eryngium vesiculosum Labill., Nov. Holl. Pl. 1: 73 t.98 (1805) | 2:252 |
| i | Foeniculum vulgare Mill., Gard. Dict., ed. 8, no.1 (1768) | 2:259 |

| | | |
|----------------------|--|-------|
| | Lilaeopsis brownii A.W.Hill = Lilaeopsis polyantha | 2:260 |
| | Lilaeopsis gunnii A.W.Hill = Lilaeopsis polyantha | |
| t | Lilaeopsis novae-zelandiae (Gand.) A.W.Hill, J. Linn. Soc. Bot.47: 549 (1927) | |
| | Lilaeopsis polyantha (Gand.) H.Eichler, Taxon 12: 296 (1963) | 2:260 |
| | Lilaeopsis sp. West Coast (A.Moscal 5655) Tas Herbarium = Lilaeopsis novae-zelandiae | |
| | Oreomyrrhis andicola (Kunth) Endl. ex Hook.f. sensu Bentham (1867) misapplied to O. ciliata , O. eriopoda & O. sessiliflora | |
| | Oreomyrrhis argentea (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 162 (1856) | 2:257 |
| | Oreomyrrhis brachycarpa (Hook.f.) Hook.f. = Oreomyrrhis eriopoda | |
| | Oreomyrrhis ciliata Hook.f., London J. Bot. 6: 471 bis (1847) | 2:258 |
| | Oreomyrrhis eriopoda (DC.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 162 (1856) | 2:257 |
| e | Oreomyrrhis gunnii Mathias & Constance, Univ. Calif. Publ. Bot. 27: 395 (1955) | 2:258 |
| e | Oreomyrrhis sessiliflora Hook.f., London J. Bot. 6: 471 bis (1847) | 2:258 |
| e | Oschatzia saxifraga (Hook.f.) Walp., Ann. Bot. Syst. (Walpers) 1: 340 (1848) | 2:251 |
| i | Pastinaca sativa L., Sp. Pl. 1: 262 (1753) | 2:261 |
| i | Petroselinum crispum (Mill.) Fuss, Fl. Transsilv. 254 (1866) | 2:256 |
| | Scandix glochidiata Labill. = Daucus glochidiatus | |
| i | Scandix pecten-veneris L., Sp. Pl. 1: 256 (1753) | 2:259 |
| | Torilis arvensis (Huds.) Link possibly recorded in error | |
| i | Torilis nodosa (L.) Gaertn., Fruct. Sem. Pl. 1: 82, t.20 (1788) | 2:262 |
| | Xanthosia dissecta Hook.f., Icon. Pl. 4: t.302 (1841) | 2:250 |
| | Xanthosia montana Sieber ex DC. = Xanthosia pilosa | |
| | Xanthosia pilosa Rudge, Trans. Linn. Soc. London 10: 301 (1811) | 2:249 |
| | Xanthosia pusilla Bunge sensu Curtis (1963) = Xanthosia tasmanica (misapplied in Tasmania) | 2:250 |
| | Xanthosia tasmanica Domin, Repert. Spec. Nov. Regni Veg. 4: 298 (1907) | 2:250 |
| | Xanthosia ternifolia J.M.Hart & Henwood, Austral. Syst. Bot. 13: 256 (2000) | |
| | Xanthosia tridentata DC., Prodr. [A. P. de Candolle] 4: 75 (1830) | 2:249 |
| APOCYNACEAE | | |
| | Alyxia buxifolia R.Br., Prodr. Fl. Nov. Holland. 470 (1810) | 3:472 |
| | Lyonsia straminea R.Br. sensu Rodway (1903) = Parsonsia brownii (misapplied in Tasmania) | |
| | Parsonsia brownii (Britten) Pichon, Notul. Syst. (Paris) 14: 10 (1950) | 3:473 |
| | Parsonsia straminea (R.Br.) F.Muell. sensu Curtis (1967) = Parsonsia brownii (misapplied in Tasmania) | 2:473 |
| i | Vinca major L., Sp. Pl. 1: 209 (1753) | 3:472 |
| AQUIFOLIACEAE | | |
| i | Ilex aquifolium L., Sp. Pl. 1: 125 (1753) | |
| ARALIACEAE | | |
| | Didiscus humilis Hook.f. f. breviscapis Domin = Trachymene humilis subsp. breviscapa | |
| | Didiscus humilis Hook.f. f. humilis = Trachymene humilis subsp. humilis | |
| | Didiscus pilosus Hook.f. nom. illeg. = Trachymene composita | |
| i | Hedera helix L., Sp. Pl. 1: 202 (1753) | |
| | Hydrocotyle asiatica L. sensu Rodway (1903) = Centella cordifolia (misapplied in Tasmania) | |
| | Hydrocotyle callicarpa Bunge, Pl. Preiss. [J.G.C.Lehman] 1: 283 (1845) | 2:246 |
| | Hydrocotyle capillaris F.Muell. ex Klatt, Linnaea 29: 707 (1859) | 2:247 |

| | | |
|--------------------------------|--|-------|
| | Hydrocotyle comocarpa F.Muell., Vict. Naturalist 3: 127 (1887) | |
| | Hydrocotyle foveolata H.Eichler, Suppl. Black's Fl. S. Austral., ed. 2, 248 (1965) | |
| | Hydrocotyle hirta R.Br. ex A.Rich., Ann. Gen. Sci. Phys. 4: 204 (1820) | 2:245 |
| | Hydrocotyle javanica Thunb. sensu Curtis (1963) = Hydrocotyle hirta (misapplied in Tasmania) | 2:245 |
| | Hydrocotyle laxiflora DC., Prodr. [A. P. de Candolle] 4: 61 (1830) | 2:246 |
| | Hydrocotyle muscosa R.Br. ex A.Rich., Ann. Gen. Sci. Phys. 4: 208 (1820) | 2:246 |
| | Hydrocotyle peduncularis R.Br. ex A.Rich. = Hydrocotyle sibthorpioides | |
| | Hydrocotyle pterocarpa F.Muell., Defin. Austral. Pl. 46 (1855) | 2:246 |
| | Hydrocotyle sibthorpioides Lam., Encycl. (Lamarck) 3: 153 (1789) | 2:245 |
| | Hydrocotyle tasmanica Hook.f. a name of uncertain application | |
| | Hydrocotyle tripartita R.Br. ex A.Rich., Ann. Gen. Sci. Phys. 4: 209 (1820) | |
| | Hydrocotyle vagans Hook.f. a name of uncertain application | |
| | Nothopanax gunnii (Hook.f.) Seem. = Pseudopanax gunnii | 2:264 |
| | Panax gunnii Hook.f. = Pseudopanax gunnii | |
| | Panax sambucifolius Sieber ex DC. sensu Bentham (1867) = Polyscias sp. Douglas-Denison (R.Schahinger HO526133) Tas Herbarium | |
| | Panax sambucifolius Sieber ex DC. = Polyscias sambucifolia | |
| | Polyscias sp. Douglas-Denison (R.Schahinger HO526133) Tas Herbarium | |
| | Polyscias aff. sambucifolia 'Douglas-Denison' (R.B.Schahinger HO526133) = Polyscias sp. Douglas-Denison (R.Schahinger HO526133) Tas Herbarium | |
| i | Polyscias sambucifolia (Sieber ex DC.) Harms, Nat. Pflanzenfam. [Engler & Prantl] III 8: 45 (1894) | |
| e | Pseudopanax gunnii (Hook.f.) K.Koch, Wochenschr. Gärtnerei Pflanzenk. 2: 366 (1859) | 2:264 |
| | Trachymene anisocarpa (Turcz.) B.L.Burtt sensu Curtis (1963) = Trachymene composita (misapplied in Tasmania) | 2:248 |
| | Trachymene australis Benth. nom. illeg. = Trachymene composita | |
| | Trachymene composita (Domin) B.L.Burtt var. composita , J. Bot. 79: 45 (1941) | 2:248 |
| | Trachymene composita (Domin) B.L.Burtt var. robertsonii (Domin) J.M.Hart, Austral. Syst. Bot. 19: 40 (2006) | |
| | Trachymene humilis (Hook.f.) Benth. subsp. breviscapa (Domin) P.S.Short, Muellera 6: 166 (1986) | 2:248 |
| | Trachymene humilis (Hook.f.) Benth. subsp. humilis , Fl. Austral. 3: 351 (1867) | 2:248 |
| ASCLEPIDACEAE | | |
| i | Gomphocarpus fruticosus (L.) W.T.Aiton subsp. fruticosus , Hortus Kew. (W.T.Aiton), ed. 2, 2: 80 (1811) | |
| ASTERACEAE (COMPOSITAE) | | |
| e | Abrotanella forsteroides (Hook.f.) Benth., Fl. Austral. 3: 554 (1867) | 2:359 |
| e | Abrotanella scapigera (F.Muell.) Benth., Fl. Austral. 3: 554 (1867) | 2:359 |
| i | Achillea distans Waldst. & Kit. ex Willd., Sp. Pl., ed. 4 [Willdenow] 3: 2207 (1800) | 2:351 |
| i | Achillea millefolium L., Sp. Pl. 2: 899 (1753) | 2:351 |
| | Achillea tanacetifolia All. = Achillea distans | 2:351 |
| | Actites megalocarpus (Hook.f.) Lander, Telopea 1: 130 (1976) | 2:390 |
| | Allittia cardiocarpa (F.Muell. ex Benth.) P.S.Short, Muellera 20: 55 (2004) | 2:296 |
| i | Ammobium alatum R.Br., Bot. Mag. 51: t.2459 (1824) | 2:343 |
| | Ammobium calyceroides (Cass.) Anderb. = Nablonium calyceroides | 2:349 |

| | | |
|---|---|-------|
| | Angianthus eriocephalus (A.Gray) Benth. = Angianthus preissianus | 2:344 |
| | Angianthus preissianus (Steetz) Benth., Fl. Austral. 3: 566 (1867) | 2:344 |
| i | Anthemis arvensis L., Sp. Pl. 2: 894 (1753) | 2:352 |
| i | Anthemis cotula L., Sp. Pl. 2: 894 (1753) | 2:352 |
| | Anthemis nobilis L. = Chamaemelum nobile | 2:352 |
| | Anthemis tinctoria L. = Cota tinctoria | 2:352 |
| | Apalochlamys billardierei DC. = Apalochlamys spectabilis | |
| | Apalochlamys spectabilis (Labill.) Steud., Nomencl. Bot. [Steudel], ed. 2, 1: 255 (1840) | 2:340 |
| i | Arctium minus (Hill) Bernh., Syst. Verz. (Bernhardi) 154 (1800) | 2:375 |
| i | Arctotheca calendula (L.) K.Lewin, J. S. African Bot. 8: 284 (1942) | 2:374 |
| i | Arctotheca populifolia (P.J.Bergius) Norl., Aquilo, Ser. Bot. 6: 84 (1967) | |
| i | Arctotis stoechadifolia P.J.Bergius, Descr. Pl. Cap. 324 (1767) | 2:374 |
| | Argentipallium dealbatum (Labill.) Paul G.Wilson, Nuytsia 8: 458 (1992) | 2:330 |
| | Argentipallium obtusifolium (Sond.) Paul G.Wilson, Nuytsia 8: 458 (1992) | 2:329 |
| e | Argentipallium × spiceri (F.Muell.) Paul G.Wilson, Nuytsia 8: 458 (1992) | 2:329 |
| | Argyrotegium fordianum (M.Gray) J.M.Ward & Breitw., New Zealand J. Bot. 41: 609 (2003) | |
| | Argyrotegium mackayi (Buchanan) J.M.Ward & Breitw., New Zealand J. Bot. 41: 609 (2003) | 2:320 |
| | Argyrotegium nitidulum (Hook.f.) J.M.Ward & Breitw., New Zealand J. Bot. 41: 609 (2003) | |
| | Argyrotegium poliochlorum (N.G.Walsh) J.M.Ward & Breitw., New Zealand J. Bot. 41: 609 (2003) | 2:320 |
| | Aster aculeatus Labill. = Olearia ramulosa | |
| | Aster argophyllus Labill. = Olearia argophylla | |
| | Aster microphyllus Labill. nom. illeg. = Olearia lepidophylla | |
| | Aster myrsinoides Labill. = Olearia myrsinoides | |
| | Aster phlogopappus Labill. = Olearia phlogopappa | |
| | Aster ramulosus Labill. = Olearia ramulosa | |
| | Aster stellulatus Labill. = Olearia stellulata | |
| | Aster subulatus Michx. = Symphotrichum subulatum | 2:299 |
| | Aster viscosus Labill. = Olearia viscosa | |
| | Bedfordia arborescens Hochr., Candollea 5: 332 (1934) | |
| e | Bedfordia linearis (Labill.) DC. subsp. linearis , Prodr. [A. P. de Candolle] 6: 441 (1838) | 2:371 |
| e | Bedfordia linearis (Labill.) DC. subsp. oblongifolia Orchard var. curvifolia Orchard, Muellera 19: 93 (2004) | 2:371 |
| e | Bedfordia linearis (Labill.) DC. subsp. oblongifolia Orchard var. oblongifolia , Muellera 19: 90 (2004) | 2:371 |
| e | Bedfordia linearis (Labill.) DC. × B. salicina (Labill.) DC. | |
| e | Bedfordia salicina (Labill.) DC., Prodr. [A. P. de Candolle] 6: 441 (1838) | 2:371 |
| | Bellis aculeata Labill. = Brachyscome aculeata | |
| | Bellis graminea Labill. = Brachyscome graminea | |
| i | Bellis perennis L., Sp. Pl. 2: 886 (1753) | 2:299 |
| | Bellis stipitata Labill. = Lagenophora stipitata | |
| i | Berkheya rigida (Thunb.) Bolus & Wolley-Dod, Trans. S.African. Philos. Soc. 14: 289 (1904) | |
| | Brachyglottis brunonis (Hook.f.) B.Nord. = Centropappus brunonis | 2:365 |
| | Brachyscome aculeata (Labill.) Less., Syn. Gen. Compos. 192 (1832) | 2:296 |

| | | |
|---|---|-------|
| | Brachyscome angustifolia A.Cunn. ex DC. = Brachyscome graminea | 2:295 |
| | Brachyscome cardiocarpa F.Muell. ex Benth. = Allittia cardiocarpa | 2:296 |
| | Brachyscome ciliaris (Labill.) Less., Syn. gen. Compos. 192 (1832) | 2:298 |
| | Brachyscome ciliaris (Labill.) Less. var. robusta Benth. = Brachyscome rigidula | |
| | Brachyscome decipiens Hook.f., London J. Bot. 6: 114 (1847) | 2:295 |
| | Brachyscome diversifolia (Graham ex Hook.) Fisch. & C.A.Mey., Index Seminum [St. Petersburg (Petropolitanus)] 2: 31 (1835) | 2:297 |
| | Brachyscome diversifolia (Graham ex Hook.) Fisch. & C.A.Mey. var. maritima Benth. = Brachyscome diversifolia | 2:297 |
| | Brachyscome graminea (Labill.) F.Muell., Fragm. (Mueller) 1(3): 49 (1858) | 2:295 |
| | Brachyscome linearifolia DC. sensu Hooker (1860) = Allittia cardiocarpa (misapplied in Tasmania) | |
| | Brachyscome nivalis F.Muell. var. alpina sensu Curtis (1963) = Brachyscome radicans (misapplied in Tasmania) | 2:297 |
| | Brachyscome parvula Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 185 (1856) | 2:298 |
| | Brachyscome perpusilla (Steetz) J.M.Black, Fl. S. Austral. [J.M. Black] 4: 587 (1929) | |
| | Brachyscome pumila Walp. sensu Hooker (1860) a name of uncertain application | |
| | Brachyscome radicans Steetz, Pl. Preiss. [J.G.C.Lehman] 1: 429 (1845) | 2:295 |
| t | Brachyscome radicata Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 127 (1852) | 2:298 |
| | Brachyscome rigidula (DC.) G.L.Davis, Proc. Linn. Soc. New South Wales 73: 219 (1948) | 2:298 |
| | Brachyscome scapiformis DC. = Brachyscome spathulata | 2:296 |
| | Brachyscome scapiformis DC. var. tenuiscapa (Hook.f.) Benth. = Brachyscome tenuiscapa | |
| | Brachyscome sieberi DC. var. gunnii DC. = Brachyscome aculeata | |
| | Brachyscome spathulata Gaudich., Voy. Uranie, Bot. 468 (1830) | |
| | Brachyscome spathulata Gaudich. subsp. glabra (DC.) H.M.Stace = Brachyscome spathulata | 2:296 |
| | Brachyscome stricta DC. = Brachyscome aculeata | 2:296 |
| | Brachyscome tadgellii Tovey & P.Morris sensu Buchanan 1995 previously recorded in error | |
| e | Brachyscome tasmanica P.S.Short, J. Adelaide Bot. Gard. 28: 160 (2014) | |
| e | Brachyscome tenuiscapa Hook.f., London J. Bot. 6: 114 (1847) | 2:294 |
| | Brachyscome tenuiscapa Hook.f. var. pubescens (Benth.) G.L.Davis sensu Buchanan (1989) = Brachyscome staceae (N.S.W., previously recorded in error) | 2:294 |
| | Bracteantha bicolor (Lindl.) Anderb. & Haegi = Xerochrysum bicolor | 2:331 |
| | Bracteantha bracteata (Vent.) Anderb. & Haegi = Xerochrysum bracteatum | |
| | Bracteantha palustris Flann = Xerochrysum palustre | |
| | Bracteantha papillosa (Labill.) Anderb. & Haegi = Xerochrysum papillosum | 2:331 |
| | Bracteantha subundulata (Sch.Bip.) Paul G.Wilson = Xerochrysum subundulatum | 2:330 |
| | Cacalia linearis Labill. = Bedfordia linearis | |
| | Cacalia salicina Labill. = Bedfordia salicina | |
| | Calea aculeata Labill. = Cassinia aculeata | |
| | Calea spectabilis Labill. = Apalochlamys spectabilis | |
| i | Calendula arvensis L., Sp. Pl., ed. 2, 2: 1303 (1763) | 2:372 |
| i | Calendula officinalis L., Sp. Pl. 2: 921 (1753) | 2:373 |
| | Calocephalus brownii (Cass.) F.Muell. = Leucophyta brownii | 2:345 |
| | Calocephalus citreus Less., Syn. Gen. Compos. 271 (1832) | 2:346 |

| | | |
|-----|--|-------|
| | Calocephalus lacteus Less., Syn. Gen. Compos. 271 (1832) | 2:345 |
| | <i>Carduus arvensis</i> (L.) Robson = Cirsium arvense | |
| | <i>Carduus lanceolatus</i> L. = Cirsium vulgare | |
| | <i>Carduus marianus</i> L. = Silybum marianum | |
| i | Carduus nutans L., Sp. Pl. 2: 821 (1753) | |
| | <i>Carduus pratensis</i> Huds. sensu Rodway (1903) = Cirsium vulgare (misapplied in Tasmania) | |
| i | Carduus pycnocephalus L., Sp. Pl., ed. 2, 2: 1151 (1763) | |
| i | Carduus tenuiflorus Curtis, Fl. Londin. (Curtis) 2(6): t.169 (1793) | 2:376 |
| i | Carthamus lanatus L., Sp. Pl. 2: 830 (1753) | 2:380 |
| i | Carthamus tinctorius L., Sp. Pl. 2: 830 (1753) | |
| | Cassinia aculeata (Labill.) R.Br. subsp. aculeata , Trans. Linn. Soc. London 12: 127 (1817) | 2:339 |
| | <i>Cassinia longifolia</i> R.Br. sensu Curtis (1963) = Cassinia trinerva (misapplied in Tasmania) | 2:340 |
| | Cassinia rugata N.G.Walsh, Muelleria 7: 141 (1990) | |
| | <i>Cassinia spectabilis</i> (Labill.) R.Br. = Apalochlamys spectabilis | 2:340 |
| | Cassinia trinerva N.A.Wakef., Vict. Naturalist 68: 69 (1951) | 2:340 |
| e | Celmisia asteliifolia Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 35 (1844) | 2:310 |
| | <i>Celmisia longifolia</i> Cass. sensu Curtis (1963) = Celmisia asteliifolia (misapplied in Tasmania) | 2:310 |
| | <i>Celmisia longifolia</i> Cass. var. saxifraga Benth. = Celmisia saxifraga | |
| e | Celmisia saxifraga (Benth.) W.M.Curtis, Taxon 17: 467 (1968) | 2:310 |
| i * | <i>Centaurea calcitrapa</i> L., Sp. Pl. 2: 917 (1753) | 2:379 |
| i * | <i>Centaurea cyanus</i> L., Sp. Pl. 2: 911 (1753) | |
| | <i>Centaurea jacea</i> L. sensu Buchanan (2007) = Centaurea × moncktonii (misapplied in Tasmania) | 2:378 |
| i | Centaurea melitensis L., Sp. Pl. 2: 917 (1753) | 2:380 |
| i | Centaurea × moncktonii C.E.Britton, Rep. Bot. Soc. Exch. Club Brit. Isles 1920, 6: 172 (1921) | 2:378 |
| | <i>Centaurea nigra</i> L. sensu Buchanan (2007) = Centaurea × moncktonii (misapplied in Tasmania) | 2:379 |
| | <i>Centaurea solstitialis</i> L. previously listed as naturalised but insufficient evidence exists to support this | 2:379 |
| | Centipeda cunninghamii (DC.) A.Braun & Asch., Index Seminum Hort. Bot. Berol. App. 6 (1867) | |
| | Centipeda elatinoides (Less.) Benth. & Hook.f. ex O.Hoffm., Nat. Pflanzenfam. [Engler & Prantl] 4(5): 280 (1892) | 2:358 |
| | <i>Centipeda minima</i> (L.) A.Braun & Asch. sensu Curtis (1963) = Centipeda elatinoides (misapplied in Tasmania) | 2:358 |
| e | Centropappus brunonis Hook.f., London J. Bot. 6: 124 (1847) | 2:365 |
| i | Chamaemelum nobile (L.) All., Fl. Pedem. 1: 185 (1785) | 2:352 |
| | <i>Chondrilla juncea</i> L. sensu Curtis (1963) anecdotal, possibly recorded in error | 2:388 |
| i | Chrysanthemoides monilifera (L.) Norl. subsp. monilifera , Stud. Calendulae 1: 374 (1943) | 2:373 |
| | <i>Chrysanthemum leucanthemum</i> L. = Leucanthemum vulgare | 2:353 |
| | <i>Chrysanthemum parthenium</i> (L.) Bernh. = Tanacetum parthenium | 2:353 |
| | Chrysocephalum apiculatum (Labill.) Steetz subsp. apiculatum , Pl. Preiss. [J.G.C.Lehman] 1: 474 (1845) | 2:332 |
| | Chrysocephalum baxteri (A.Cunn. ex DC.) Anderb., Compositae Newslett. 19: 22 (1991) | 2:328 |
| | Chrysocephalum semipapposum (Labill.) Steetz subsp. asperum (Steetz) Paul G.Wilson, Nuytsia 27: 70 (2016) | |

| | | |
|-----|---|-------|
| | Chrysocephalum semipapposum (Labill.) Steetz subsp. lineare Paul G.Wilson, Nuytsia 27: 67 (2016) | |
| | Chrysocephalum semipapposum (Labill.) Steetz subsp. semipapposum , Pl. Preiss. [J.G.C.Lehman] 1: 474 (1845) | 2:332 |
| | Chrysocoma cinerea Labill. = Ozothamnus turbinatus | |
| | Chrysocoma reticulata Labill. = Ozothamnus reticulatus | |
| | Chrysocoma squamata Labill. = Leptorhynchus squamatus | |
| i | Cichorium intybus L., Sp. Pl. 2: 813 (1753) | 2:380 |
| i | Cirsium arvense (L.) Scop. var. arvense , Fl. Carniol., ed. 2, 2: 126 (1772) | 2:377 |
| i | Cirsium vulgare (Savi) Ten., Fl. Napol. 5: 209 (1836) | 2:376 |
| | Cnicus arvensis (L.) Roth = Cirsium arvense | 2:377 |
| | Conyza albida Willd. ex Spreng. = Conyza sumatrensis | 2:315 |
| i | Conyza bonariensis (L.) Cronquist, Bull. Torrey Bot. Club 70: 632 (1943) | 2:315 |
| i | Conyza canadensis (L.) Cronquist, Bull. Torrey Bot. Club 70: 632 (1943) | 2:314 |
| | Conyza chilensis Spreng. = Conyza primulifolia | 2:314 |
| | Conyza floribunda Kunth sensu Curtis (1963) = Conyza sumatrensis (misapplied in Tasmania) | 2:315 |
| i x | Conyza primulifolia (Lam.) Cuatrec. & Lourteig, Phytologia 58: 475 (1985) | 2:314 |
| | Conyza scabiosifolia J.Rémy = Conyza primulifolia | 2:314 |
| i | Conyza sumatrensis (Retz.) E.Walker, J. Jap. Bot. 46(3): 72 (1971) | 2:315 |
| | Coronidium gunnianum (Hook.) N.G.Walsh, Muellera 32: 20 (2014) | |
| | Coronidium sp. Lowland Swamps (V.Stajsic 4226) Vic Herbarium = Coronidium gunnianum | |
| | Coronidium monticola N.G.Walsh, Muellera 32: 21 (2014) | |
| | Coronidium scorpioides (Labill.) Paul G.Wilson, Nuytsia 18: 326 (2008) | 2:328 |
| | Cota tinctoria (L.) J.Gay formerly considered naturalised though no evidence exists to support this | 2:352 |
| | Cotula alpina (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 192 t.51A (1856) | 2:356 |
| | Cotula australis (Sieber ex Spreng.) Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 128 (1852) | 2:356 |
| i | Cotula coronopifolia L., Sp. Pl. 2: 892 (1753) | 2:356 |
| | Cotula coronopifolia L. var. integrifolia (Hook.f.) Rodway = Cotula coronopifolia | |
| | Cotula filicula (Hook.f.) Benth. = Leptinella filicula | 2:357 |
| | Cotula filifolia Thunb. sensu Rodway (1903) = Cotula vulgaris var. australasica (misapplied in Tasmania) | |
| | Cotula integrifolia Hook.f. nom. illeg. = Cotula coronopifolia | |
| | Cotula longipes (Hook.f.) W.M.Curtis = Leptinella longipes | 2:357 |
| | Cotula reptans (Benth.) Benth. = Leptinella reptans | 2:357 |
| | Cotula reptans (Benth.) Benth. var. major Benth. = Leptinella longipes | |
| | Cotula vulgaris Levyns var. australasica J.H.Willis, Vict. Naturalist 73: 201 (1957) | 2:355 |
| | Craspedia alpina Backh. ex Hook.f. = Craspedia macrocephala | 2:347 |
| | Craspedia coolaminica J.Everett & Joy Thomps. = Craspedia gracilis | |
| e | Craspedia cynurica Rozefelds & A.M.Buchanan, Kanunna 4: 106 (2011) | |
| e | Craspedia glabrata (Hook.f.) Rozefelds, Telopea 9: 814 (2002) | 2:347 |
| e | Craspedia glauca (Labill.) Spreng., Syst. Veg. (ed. 16) [Sprengel] 3: 441 (1826) | 2:347 |
| | Craspedia glauca (Labill.) Spreng. var. glabrata Hook.f. ex W.M.Curtis nom. inval. = Craspedia glabrata | 2:347 |

| | | |
|-----|---|-------|
| | <i>Craspedia glauca</i> (Labill.) Spreng. var. <i>gracilis</i> W.M.Curtis nom. inval. = <i>Craspedia gracilis</i> | 2:347 |
| | <i>Craspedia glauca</i> (Labill.) Spreng. var. <i>macrocephala</i> W.M.Curtis nom. inval. = <i>Craspedia macrocephala</i> | 2:347 |
| | <i>Craspedia gracilis</i> Hook.f., London J. Bot. 6: 118 (1847) | |
| e | <i>Craspedia macrocephala</i> Hook., Bot. Mag. 62: t. 3415 (1835) | 2:347 |
| | <i>Craspedia paludicola</i> J.Everett & Doust, Telopea 5: 35 (1992) | |
| e | <i>Craspedia preminghana</i> Rozefelds, Telopea 9: 816 (2002) | |
| | <i>Craspedia richea</i> Cass. nom. illeg. = <i>Craspedia glauca</i> | |
| | <i>Craspedia richea</i> Cass. var. <i>alpina</i> (Backh. ex Hook.f.) Benth. = <i>Craspedia macrocephala</i> | |
| | <i>Craspedia richea</i> Cass. var. <i>glabrata</i> Hook.f. = <i>Craspedia glabrata</i> | |
| | <i>Craspedia richea</i> Cass. var. <i>gracilis</i> (Hook.f.) Hook.f. = <i>Craspedia gracilis</i> | |
| | <i>Craspedia richea</i> Cass. var. <i>linearis</i> Hook.f. = <i>Craspedia gracilis</i> | |
| | <i>Craspedia richea</i> Cass. var. <i>macrocephala</i> (Hook.) Benth. = <i>Craspedia macrocephala</i> | |
| e | <i>Craspedia rosulata</i> Rozefelds & A.M.Buchanan, Kanunnah 4: 111 (2011) | |
| i | <i>Crepis capillaris</i> (L.) Wallr., Erst. Beitr. Fl. Hercyn. 287 (1840) | 2:384 |
| | <i>Crepis nicaeensis</i> Baldinger & Pers. sensu Curtis (1963) recorded in error | 2:384 |
| i | <i>Crepis setosa</i> Haller f., Arch. Bot. (Leipzig) 1(2): 1 (1797) | 2:384 |
| | <i>Cryptostemma calendula</i> (L.) R.Br. = <i>Arctotheca calendula</i> | |
| | <i>Cymbonotus lawsonianus</i> Gaudich. sensu Curtis (1963) = <i>Cymbonotus preissianus</i> (misapplied in Tasmania) | 2:373 |
| | <i>Cymbonotus preissianus</i> Steetz, Pl. Preiss 1: 486 (1845) | 2:373 |
| i x | <i>Cynara cardunculus</i> L. subsp. <i>flavescens</i> Wiklund, Bot. J. Linn. Soc. 109: 120 (1992) | |
| i | <i>Delairea odorata</i> Lem., Ann. Sci. Nat., Bot., sér. 3, 1: 380 (1844) | 2:367 |
| i | <i>Dimorphotheca fruticosa</i> (L.) DC., Prodr. [A. P. de Candolle] 6: 71 (1838) | |
| i | <i>Dittrichia graveolens</i> (L.) Greuter, Exsicc. Genav. Conserv. Bot. Distrib. Fasc. 4: 71 (1973) | 2:348 |
| | <i>Erechtites argutus</i> (A.Rich.) DC. = <i>Senecio glomeratus</i> subsp. <i>glomeratus</i> | |
| | <i>Erechtites gunnii</i> Hook.f. = <i>Senecio gunnii</i> | |
| | <i>Erechtites hispidula</i> (A.Rich.) DC. = <i>Senecio hispidulus</i> | |
| | <i>Erechtites prenathoides</i> (A.Rich.) DC. = <i>Senecio prenanthoides</i> | |
| | <i>Erechtites quadridentata</i> (Labill.) DC. = <i>Senecio quadridentatus</i> | |
| | <i>Erechtites quadridentata</i> (Labill.) DC. var. <i>glabrescens</i> (DC.) Benth. sensu Bentham (1867) = <i>Senecio quadridentatus</i> (misapplied in Tasmania) | |
| | <i>Erechtites quadridentata</i> (Labill.) DC. var. <i>gunnii</i> (Hook.f.) Benth. = <i>Senecio gunnii</i> | |
| | <i>Erigeron bellidioides</i> (Hook.f.) S.J.Forbes & D.I.Morris = <i>Pappochroma bellidioides</i> | |
| | <i>Erigeron gunnii</i> (Hook.f.) F.Muell. ex Hook.f. = <i>Pappochroma gunnii</i> | |
| | <i>Erigeron gunnii</i> (Hook.f.) F.Muell. ex Hook.f. var. <i>bellidioides</i> (Hook.f.) Hook.f. = <i>Pappochroma bellidioides</i> | |
| i | <i>Erigeron karvinskianus</i> DC., Prodr. [A. P. de Candolle] 5: 285 (1836) | 2:312 |
| | <i>Erigeron linifolius</i> Willd. = <i>Conyza bonariensis</i> | |
| | <i>Erigeron mucronatus</i> DC. = <i>Erigeron karvinskianus</i> | 2:312 |
| | <i>Erigeron pappocromus</i> Labill. = <i>Pappochroma pappocromum</i> | 2:311 |
| | <i>Erigeron pappocromus</i> Labill. var. <i>billardieri</i> Benth. = <i>Pappochroma pappocromum</i> | 2:311 |
| | <i>Erigeron pappocromus</i> Labill. var. <i>gunnii</i> Benth. = <i>Pappochroma gunnii</i> | 2:312 |

| | | |
|---|---|-------|
| | Erigeron pappocromus Labill. var. oblongatus Benth. = Pappochroma tasmanicum | |
| | Erigeron pappocromus Labill. var. stellatus (Hook.f.) Benth. = Pappochroma stellatum | |
| | Erigeron stellatus (Hook.f.) W.M.Curtis = Pappochroma stellatum | 2:312 |
| | Erigeron tasmanicus (Hook.f.) Hook.f. = Pappochroma tasmanicum | 2:312 |
| | Erigeron trigonus S.J.Forbes & D.I.Morris = Pappochroma trigonum | |
| | Euchiton argentifolius (N.A.Wakef.) Anderb. = Argyrotegium mackayi | 2:320 |
| | Euchiton collinus Cass. = Euchiton japonicus | 2:319 |
| | Euchiton fordianus (M.Gray) P.S.Short = Argyrotegium fordianum | |
| | Euchiton involucratus (G.Forst.) Holub, Folia Geobot. Phytotax. 9: 271 (1974) | 2:318 |
| | Euchiton japonicus (Thunb.) Holub, Folia Geobot. Phytotax. 9: 271 (1974) | 2:319 |
| t | Euchiton lateralis (C.J.Webb) Breitw. & J.M.Ward, New Zealand J. Bot. 36: 303 (1998) | |
| | Euchiton limosus (D.G.Drury) Holub, Folia Geobot. Phytotax. 9: 271 (1974) | |
| e | Euchiton litticola A.M.Buchanan, Pap. & Proc. Roy. Soc. Tasmania 133: 115 (1999) | |
| | Euchiton poliochlorus N.G.Walsh = Argyrotegium poliochlorum | 2:320 |
| | Euchiton sphaericus (Willd.) Holub, Folia Geobot. Phytotax. 9: 271 (1974) | |
| | Euchiton traversii (Hook.f.) Holub, Folia Geobot. Phytotax. 9: 271 (1974) | 2:320 |
| | Euchiton umbricola (J.H.Willis) Anderb., Opera Bot. 104: 167 (1991) | 2:319 |
| | Eupatorium ferrugineum Labill. = Ozothamnus ferrugineus | |
| | Eupatorium rosmarinifolium Labill. = Ozothamnus rosmarinifolius | |
| | Eurybia alpina Hook.f. = Olearia tasmanica | |
| | Eurybia argophylla (Labill.) Cass. = Olearia argophylla | |
| | Eurybia ciliata Benth. = Olearia ciliata | |
| | Eurybia ericoides Steetz = Olearia ericoides | |
| | Eurybia erubescens Sieber ex DC. = Olearia erubescens | |
| | Eurybia floribunda Hook.f. = Olearia floribunda | |
| | Eurybia fulvida Cass. nom. illeg. = Olearia stellulata | |
| | Eurybia glandulosa (Labill.) DC. = Olearia glandulosa | |
| | Eurybia gunniana DC. = Olearia phlogopappa subsp. gunniana | |
| | Eurybia ledifolia DC. = Olearia ledifolia | |
| | Eurybia linearifolia DC. = Olearia axillaris | |
| | Eurybia linifolia Hook.f. = Olearia glutinosa | |
| | Eurybia lirata (Sims) DC. = Olearia lirata | |
| | Eurybia myrsinoides (Labill.) Nees = Olearia myrsinoides | |
| | Eurybia obcordata Hook.f. = Olearia obcordata | |
| | Eurybia persoonioides DC. = Olearia persoonioides | |
| | Eurybia pinifolia Hook.f. = Olearia pinifolia | |
| | Eurybia ramulosa (Labill.) DC. = Olearia ramulosa | |
| | Eurybia viscosa (Labill.) Cass. = Olearia viscosa | |
| i | Euryops abrotanifolius (L.) DC., Prodr. [A. P. de Candolle] 6: 443 (1838) | 2:372 |
| e | Ewartia catipes (DC.) Beauverd, Bull. Soc. Bot. Genève, Ser. 2, 2: 238 (1910) | 2:322 |
| e | Ewartia meredithiae (F.Muell.) Beauverd, Bull. Soc. Bot. Genève, Ser. 2, 2: 240 (1910) | 2:322 |
| e | Ewartia planchonii (Hook.f.) Beauverd, Bull. Soc. Bot. Genève, Ser. 2, 3: 253 (1911) | 2:321 |
| | Felicia erigeroides DC. sensu Raphael (1955) = Erigeron karvinskianus (misapplied in Tasmania) | |

| | | |
|---|---|-------|
| | Filago gallica L. = Logfia gallica | 2:315 |
| i | Galinsoga parviflora Cav., Icon. [Cavanilles] 3: 41 t.281 (1795) | 2:349 |
| i | Gamochaeta calviceps (Fernald) Cabrera, Bol. Soc. Argent. Bot. 9: 368 (1961) | |
| i | Gamochaeta purpurea (L.) Cabrera, Bol. Soc. Argent. Bot. 9: 377 (1961) | 2:321 |
| i | Gazania linearis (Thunb.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 624 (1917) | |
| | Gnaphalium alpinum F.Muell. ex Hook.f. nom. illeg. = Euchiton umbricola | |
| | Gnaphalium apiculatum Labill. = Chrysocephalum apiculatum | |
| | Gnaphalium argentifolium N.A.Wakef. = Argyrotegium mackayi | 2:320 |
| | Gnaphalium candidissimum Lam. = Vellereophyton dealbatum | 2:318 |
| | Gnaphalium collinum Labill. var. collinum = Euchiton japonicus | 2:319 |
| | Gnaphalium collinum Labill. var. monocephalum Hook.f. = Euchiton lateralis | 2:319 |
| | Gnaphalium fordianum M.Gray = Argyrotegium fordianum | |
| | Gnaphalium indicum L. recorded in error | |
| | Gnaphalium indutum Hook.f. subsp. indutum , London J. Bot. 6: 121 (1847) | 2:320 |
| | Gnaphalium involucratum G.Forst. = Euchiton involucratus | 2:318 |
| | Gnaphalium japonicum Thunb. = Euchiton japonicus | |
| | Gnaphalium japonicum Thunb. var. radicans = Argyrotegium mackayi | |
| | Gnaphalium luteo-album L. = Helichrysum luteoalbum | 2:318 |
| | Gnaphalium planchonii Hook.f. = Ewartia planchonii | |
| | Gnaphalium polycaulon Pers. sensu Buchanan (2009) recorded in error | |
| | Gnaphalium purpureum L. = Gamochaeta purpurea | 2:321 |
| | Gnaphalium semipapposum Labill. = Chrysocephalum semipapposum | |
| | Gnaphalium supinum L. sensu Curtis (1963) = Argyrotegium poliochlorum (misapplied in Tasmania) | 2:320 |
| | Gnaphalium traversii Hook.f. = Euchiton traversii | 2:320 |
| | Gnaphalium umbricola J.H.Willis = Euchiton umbricola | 2:319 |
| | Hedypnois cretica (L.) Dum.Cours. = Leontodon rhagadioloides | 2:382 |
| | Hedypnois rhagadioloides (L.) F.W.Schmidt = Leontodon rhagadioloides | 2:382 |
| | Helichrysum acuminatum DC. = Xerochrysum subundulatum | 2:330 |
| | Helichrysum adenophorum F.Muell. recorded in error | |
| | Helichrysum antennarium (DC.) F.Muell. ex Benth. = Ozothamnus antennaria | 2:336 |
| | Helichrysum apiculatum (Labill.) D.Don = Chrysocephalum apiculatum | 2:332 |
| | Helichrysum argophyllum (A.Cunn. ex DC.) N.A.Wakef. = Ozothamnus argophyllum | 2:335 |
| | Helichrysum baccharioides F.Muell. nom. illeg. = Ozothamnus hookeri | |
| | Helichrysum backhousei (Hook.f.) F.Muell. ex Benth. = Ozothamnus rodwayi | 2:337 |
| | Helichrysum backhousei (Hook.f.) F.Muell. ex Benth. var. backhausii = Ozothamnus rodwayi var. rodwayi | 2:337 |
| | Helichrysum backhousei (Hook.f.) F.Muell. ex Benth. var. kingii W.M.Curtis = Ozothamnus rodwayi var. kingii | 2:337 |
| | Helichrysum backhousei (Hook.f.) F.Muell. ex Benth. var. oreophilum W.M.Curtis = Ozothamnus rodwayi var. oreophilus | 2:337 |
| | Helichrysum baxteri A.Cunn. ex DC. = Chrysocephalum baxteri | 2:328 |
| | Helichrysum bicolor Lindl. = Xerochrysum bicolor | 2:331 |
| | Helichrysum bracteatum (Vent.) Andrews = Xerochrysum bracteatum | |

| | | |
|---|--|-------|
| | Helichrysum bracteatum (Vent.) Andrews var. albidum DC. = Xerochrysum papillosum (Tasmanian material) | |
| | Helichrysum bracteolatum (Hook.f.) Benth. = Ozothamnus gunnii | |
| | Helichrysum cinereum (Labill.) Benth. = Ozothamnus turbinatus | |
| | Helichrysum costatifructum R.V.Sm. = Ozothamnus costatifructus | 2:332 |
| | Helichrysum dealbatum Labill. = Argentipallium dealbatum | 2:330 |
| | Helichrysum dendroideum N.A.Wakef. = Ozothamnus ferrugineus | 2:335 |
| | Helichrysum ericeteum W.M.Curtis = Ozothamnus ericifolius | 2:334 |
| | Helichrysum expansifolium (P.Morris & J.H.Willis) N.T.Burb. = Ozothamnus expansifolius | 2:337 |
| | Helichrysum gunnii (Hook.f.) Benth. = Ozothamnus gunnii | 2:333 |
| | Helichrysum hookeri (Sond.) Druce = Ozothamnus hookeri | 2:337 |
| | Helichrysum ledifolium (A.Cunn. ex DC.) Benth. = Ozothamnus ledifolius | 2:334 |
| | Helichrysum leucopsideum DC., Prodr. [A. P. de Candolle] 6: 193 (1838) | 2:331 |
| | Helichrysum luteoalbum (L.) Rchb., Handb. Gewächsk., ed. 2, 2: 1460 (1829) | 2:318 |
| | Helichrysum lycopodioides (Hook.f.) Benth. = Ozothamnus lycopodioides | 2:338 |
| | Helichrysum milliganii Hook.f. = Xerochrysum milliganii | 2:330 |
| | Helichrysum obcordatum (DC.) Benth. = Ozothamnus obcordatus | 2:336 |
| | Helichrysum obtusifolium F.Muell. & Sond. ex Sond. = Argentipallium obtusifolium | 2:329 |
| | Helichrysum papillosum Labill. = Xerochrysum papillosum | 2:331 |
| | Helichrysum paraliium (N.T.Burb.) W.M.Curtis = Ozothamnus turbinatus | 2:333 |
| | Helichrysum pleurandroides F.Muell. sensu Buchanan et al. (1989) = Ozothamnus ericifolius | 2:334 |
| e | Helichrysum pumilum Hook.f. var. pumilum , Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 213 (1856) | 2:330 |
| e | Helichrysum pumilum Hook.f. var. spathulatum A.M.Buchanan, Pap. & Proc. Roy. Soc. Tasmania 121: 54 (1987) | 2:330 |
| | Helichrysum purpurascens (DC.) W.M.Curtis = Ozothamnus purpurascens | 2:334 |
| | Helichrysum reticulatum (Labill.) Benth. = Ozothamnus reticulatus | 2:332 |
| | Helichrysum rosmarinifolium (Labill.) Benth. = Ozothamnus rosmarinifolius | 2:335 |
| | Helichrysum rutidolepis DC. sensu Buchanan et al (1989) = Coronidium gunnianum & C. monticola (misapplied in Tasmania) | |
| | Helichrysum scorpioides Labill. = Coronidium scorpioides | 2:328 |
| | Helichrysum scutellifolium (Hook.f.) F.Muell. = Ozothamnus scutellifolius | 2:338 |
| | Helichrysum selaginoides (Sond. & F.Muell.) Benth. = Ozothamnus selaginoides | 2:339 |
| | Helichrysum semipapposum (Labill.) DC. = Chrysocephalum semipapposum | 2:332 |
| | Helichrysum spiceri F.Muell. = Argentipallium × spiceri | 2:329 |
| | Helichrysum thyrsoides (DC.) P.Morris & J.H.Willis = Ozothamnus thyrsoides | 2:335 |
| | Helipterum albicans (A.Cunn.) DC. var. incanum (Hook.) Paul G.Wilson = Leucochrysum albicans var. tricolor | 2:324 |
| | Helipterum anthemoides (Sieber ex Spreng.) DC. = Rhodanthe anthemoides | 2:324 |
| | Helipterum australe (A.Gray) Druce = Triptilodiscus pygmaeus | |
| | Helipterum demissum (A.Gray) Druce = Hyalosperma demissum | 2:325 |
| | Helipterum exiguum F.Muell. = Hyalosperma demissum | |
| | Helipterum incanum DC. nom. illeg., nom. superfl. = Leucochrysum albicans var. tricolor | |
| | Helminthia echioides (L.) Gaertn. = Helminthotheca echioides | |

| | | |
|-----|--|-------|
| i | Helminthotheca echioides (L.) Holub, Folia Geobot. Phytotax. 8: 176 (1973) | 2:383 |
| | Hieracium aurantiacum L. subsp. carpathicola Nägeli & Peter = Pilosella aurantiaca subsp. aurantiaca | 2:385 |
| | Hieracium brunneocroceum Pugsley = Pilosella aurantiaca subsp. aurantiaca | 2:385 |
| | Hieracium pilosella L. = Pilosella officinarum subsp. officinarum | |
| | Hyalosperma demissum (A.Gray) Paul G.Wilson, Nuytsia 7: 85 (1989) | 2:325 |
| i | Hypochaeris glabra L., Sp. Pl. 2: 811 (1753) | 2:386 |
| i | Hypochaeris radicata L., Sp. Pl. 2: 811 (1753) | 2:385 |
| | Inula graveolens (L.) Desf. = Dittrichia graveolens | 2:348 |
| | Isoetopsis graminifolia Turcz., Bull. Soc. Imp. Naturalistes Moscou 24: 175 t.3 (1851) | 2:360 |
| | Ixiolaena supina F.Muell. = Leiocarpa supina | 2:341 |
| | Ixodia achlaena D.I.Morris = Odixia achlaena | |
| | Ixodia angusta (N.A.Wakef.) N.T.Burb. = Odixia angusta | 2:343 |
| i | Lactuca saligna L., Sp. Pl. 2: 796 (1753) | 2:388 |
| i * | Lactuca serriola L. f. integrifolia (Gray) S.D.Prince & R.N.Carter, Cent. Pl. 2: 29 (1756) | 2:389 |
| i | Lactuca serriola L. f. serriola , Cent. Pl. 2: 29 (1756) | |
| | Lagenophora billardierei Cass. = Lagenophora stipitata | |
| | Lagenophora emphysopus Hook.f. nom. illeg. = Solenogyne gunnii | |
| | Lagenophora gracilis Steetz, Pl. Preiss. [J.G.C.Lehman] 1: 431 (1845) | |
| | Lagenophora gunniana Steetz = Lagenophora huegelii | |
| | Lagenophora huegelii Benth., Enum. Pl. [Endlicher]: 59 (1837) | 2:290 |
| | Lagenophora latifolia Hook.f. = Lagenophora stipitata | |
| | Lagenophora montana Hook.f., London J. Bot. 6: 113 (1847) | |
| | Lagenophora stipitata (Labill.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 630 (1917) | 2:290 |
| i | Lapsana communis L. subsp. communis , Sp. Pl. 2: 811 (1753) | |
| i t | Lasiospermum bipinnatum (Thunb.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 631 (1917) | 2:350 |
| | Leiocarpa supina (F.Muell.) Paul G.Wilson, Nuytsia 13: 604 (2001) | 2:341 |
| | Leontodon hirtus L. sensu Rodway (1903) = Leontodon saxatilis (misapplied in Tasmania) | |
| | Leontodon leysseri (Wallr.) Beck sensu Curtis (1963) = Leontodon saxatilis (misapplied in Tasmania) | 2:386 |
| i | Leontodon rhagadioloides (L.) Enke & Zidorn, Organisms Diversity Evol. 12: 14 (2012) | 2:382 |
| i | Leontodon saxatilis Lam., Fl. Franç. (Lamarck) 2: 115 (1779) (as "1778") | 2:386 |
| | Leontodon taraxacoides (Vill.) Mérat = Leontodon saxatilis | 2:386 |
| | Leptinella filicula (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 194 (1856) | 2:357 |
| | Leptinella intricata Hook.f. = Leptinella reptans | |
| | Leptinella longipes Hook.f., London J. Bot. 6: 117 (1847) | 2:357 |
| | Leptinella reptans (Benth.) D.G.Lloyd & C.J.Webb, New Zealand J. Bot. 25: 103 (1987) | 2:357 |
| | Leptorhynchos elongatus DC., Prodr. [A. P. de Candolle] 6: 160 (1838) | 2:323 |
| | Leptorhynchos linearis Less. sensu Curtis (1963) = Leptorhynchos nitidulus (misapplied in Tasmania) | 2:323 |
| | Leptorhynchos nitidulus DC., Prodr. [A. P. de Candolle] 6: 160 (1838) | 2:323 |
| | Leptorhynchos squamatus (Labill.) Less. subsp. alpinus Flann, Austral. Syst. Bot. 15: 217 (2002) | 2:323 |
| | Leptorhynchos squamatus (Labill.) Less. subsp. squamatus , Syn. Gen. Compos. 273 (1832) | 2:323 |
| i | Leucanthemum × superbum (Bergmans ex J.W.Ingram) D.H.Kent, Watsonia 18: 89 (1990) | |
| i | Leucanthemum vulgare Lam., Fl. Franç. (Lamarck) 2: 137 (1779) (as "1778") | 2:353 |

| | | |
|-----|--|-------|
| | Leucochrysum albicans (A.Cunn.) Paul G.Wilson subsp. tricolor (DC.) N.G.Walsh, Muellera 34: 13 (2015) | 2:324 |
| | Leucophyta brownii Cass., Dict. Sci. Nat., ed. 2 [F.Cuvier] 26: 159 (1823) | 2:345 |
| i | Logfia gallica (L.) Coss. & Germ., Ann. Sci. Nat., Bot., sér. 2, 20: 291 (1843) | 2:315 |
| i * | <i>Matricaria chamomilla</i> L., Sp. Pl. 2: 891 (1753) | |
| | <i>Matricaria chamomilla</i> L. sensu de Salas & Baker 2014 = Matricaria discoidea (misapplied in Tasmania) | |
| i | Matricaria discoidea DC., Prodr. [A. P. de Candolle] 6: 50 (1838) | 2:354 |
| | <i>Matricaria matricarioides</i> (Less.) Porter = Matricaria discoidea | 2:354 |
| | <i>Matricaria perforata</i> Mérat = Tripleurospermum maritimum subsp. inodorum | |
| | <i>Matricaria recutita</i> L. sensu de Salas & Baker 2014 = Matricaria chamomilla (misapplied in Tasmania) | |
| | <i>Microseris forsteri</i> Hook.f. nom. illeg. = Microseris lanceolata | |
| | Microseris lanceolata (Walp.) Sch.Bip., Jahresber. Pollichia 22-24: 310 (1866) | 2:381 |
| | <i>Microseris scapigera</i> (Sol. ex A.Cunn.) Sch.Bip. sensu Curtis (1963) = Microseris lanceolata (misapplied in Tasmania) | 2:381 |
| | Microseris walteri Gand., Bull. Soc. Bot. France 65: 52 (1918) | |
| | Millotia muelleri (Sond.) P.S.Short, Muellera 7: 246 (1990) | |
| | Millotia tenuifolia Cass. var. tenuifolia , Ann. Sci. Nat. (Paris) 17: 417 (1829) | 2:342 |
| | <i>Myriogyne minuta</i> (G.Forst.) Less. sensu Bentham (1867) = Centipeda elatinoidea (misapplied in Tasmania) | |
| e | Nablonium calyceroides Cass., Dict. Sci. Nat., ed. 2 [F.Cuvier] 34: 101 (1825) | 2:349 |
| e | Odivia achlaena (D.I.Morris) Orchard, Brunonia 4: 194 (1982) | |
| e | Odivia angusta (N.A.Wakef.) Orchard, Brunonia 4: 194 (1982) | 2:343 |
| | Olearia algida N.A.Wakef., Vict. Naturalist 73: 97 (1956) | 2:308 |
| | <i>Olearia alpina</i> (Hook.f.) W.M.Curtis nom. illeg. = Olearia tasmanica | 2:304 |
| e | Olearia archeri Lander, Muellera 7: 117 (1989) | |
| | Olearia argophylla (Labill.) F.Muell. ex Benth., Fl. Austral. 3: 470 (1867) | 2:303 |
| | Olearia axillaris (DC.) F.Muell. ex Benth., Fl. Austral. 3: 475 (1867) | 2:307 |
| | Olearia ciliata (Benth.) F.Muell. ex Benth., Fl. Austral. 3: 488 (1867) | 2:309 |
| e | Olearia ericoides (Steetz) N.A.Wakef., Vict. Naturalist. 73: 97 (1956) | 2:307 |
| | Olearia erubescens (DC.) Dippel, Handb. Laubholzk 1: 290 (1889) | 2:303 |
| | Olearia floribunda (Hook.f.) Benth., Fl. Austral. 3: 477 (1867) | 2:308 |
| | Olearia glandulosa (Labill.) Benth., Fl. Austral. 3: 483 (1867) | 2:309 |
| | Olearia glutinosa (Lindl.) Benth., Fl. Austral. 3: 482 (1867) | 2:309 |
| | <i>Olearia gunniana</i> (DC.) Hook.f. ex Hook. = Olearia phlogopappa subsp. gunniana | |
| e | Olearia hookeri (Sond.) Benth., Fl. Austral. 3: 483 (1867) | 2:309 |
| | <i>Olearia lanceolata</i> (Benth.) D.I.Morris sensu Morris (1977) = Olearia archeri (misapplied in Tasmania) | |
| e | Olearia ledifolia (DC.) Benth., Fl. Austral. 3: 472 (1867) | 2:305 |
| | Olearia lepidophylla (Pers.) Benth., Fl. Austral. 3: 477 (1867) | 2:308 |
| | Olearia lirata (Sims) Hutch., Gard. Chron., Ser. 3, 61: 14 fig.4 (1917) | 2:306 |
| | Olearia myrsinoides (Labill.) F.Muell. ex Benth., Fl. Austral. 3: 470 (1867) | 2:303 |
| | <i>Olearia myrsinoides</i> (Labill.) F.Muell. ex Benth. var. <i>erubescens</i> (DC.) Benth. = Olearia erubescens | |
| e | Olearia obcordata (Hook.f.) Benth., Fl. Austral. 3: 471 (1867) | 2:304 |
| e | Olearia persoonioides (DC.) Benth., Fl. Austral. 3: 471 (1867) | 2:304 |

| | | |
|-----|---|-------|
| | <i>Olearia persoonioides</i> (DC.) Benth. var. <i>alpina</i> (Hook.f.) Benth. = <i>Olearia tasmanica</i> | |
| | <i>Olearia persoonioides</i> (DC.) Benth. var. <i>lanceolata</i> Benth. sensu Morris (1977) = <i>Olearia archeri</i> | |
| e | <i>Olearia phlogopappa</i> (Labill.) DC. subsp. <i>angustifolia</i> (Hook.f.) Messina, Austral. Syst. Bot. 26: 57 (2013) | 2:306 |
| e | <i>Olearia phlogopappa</i> (Labill.) DC. subsp. <i>gunniana</i> (DC.) Messina, Austral. Syst. Bot. 26: 60 (2013) | 2:306 |
| | <i>Olearia phlogopappa</i> (Labill.) DC. subsp. <i>insularis</i> Messina, Austral. Syst. Bot. 26: 63 (2013) | |
| e | <i>Olearia phlogopappa</i> (Labill.) DC. subsp. <i>phlogopappa</i> , Prodr. [A. P. de Candolle] 5: 272 (1836) | 2:306 |
| | <i>Olearia phlogopappa</i> (Labill.) DC. subsp. <i>salicina</i> (Hook.f.) Messina, Austral. Syst. Bot. 26: 63 (2013) | 2:306 |
| e | <i>Olearia phlogopappa</i> (Labill.) DC. subsp. <i>subrepanda</i> (DC.) Messina, Austral. Syst. Bot. 26: 67 (2013) | 2:306 |
| | <i>Olearia phlogopappa</i> (Labill.) DC. var. <i>angustifolia</i> (Hutch.) W.M.Curtis = <i>Olearia phlogopappa</i> subsp. <i>angustifolia</i> | 2:306 |
| | <i>Olearia phlogopappa</i> (Labill.) DC. var. <i>brevipes</i> (Hutch.) W.M.Curtis = <i>Olearia phlogopappa</i> subsp. <i>gunniana</i> | 2:306 |
| | <i>Olearia phlogopappa</i> (Labill.) DC. var. <i>microcephala</i> (Hutch.) W.M.Curtis = <i>Olearia phlogopappa</i> subsp. <i>gunniana</i> | 2:306 |
| | <i>Olearia phlogopappa</i> (Labill.) DC. var. <i>salicifolia</i> (Hutch.) W.M.Curtis = <i>Olearia phlogopappa</i> subsp. <i>salicina</i> | 2:306 |
| | <i>Olearia phlogopappa</i> (Labill.) DC. var. <i>subrepanda</i> (DC.) J.H.Willis = <i>Olearia phlogopappa</i> subsp. <i>subrepanda</i> | 2:306 |
| e | <i>Olearia pinifolia</i> (Hook.f.) Benth., Fl. Austral. 3: 471 (1867) | 2:305 |
| | <i>Olearia ramulosa</i> (Labill.) Benth., Fl. Austral. 3: 476 (1867) | 2:307 |
| e | <i>Olearia rugosa</i> (F.Muell. ex W.Archer bis) Hutch. subsp. <i>intermedia</i> Messina, Austral. Syst. Bot. 26: 76 (2013) | |
| | <i>Olearia stellulata</i> (Labill.) DC., Prodr. [A. P. de Candolle] 5: 272 (1836) | 2:305 |
| | <i>Olearia stellulata</i> (Labill.) DC. var. <i>lirata</i> (Sims) Benth. = <i>Olearia lirata</i> | |
| | <i>Olearia stellulata</i> (Labill.) DC. var. <i>quercifolia</i> (Sieber ex DC.) Benth. sensu Bentham (1867) = <i>Olearia rugosa</i> subsp. <i>intermedia</i> (misapplied in Tasmania) | |
| e | <i>Olearia tasmanica</i> W.M.Curtis, Vict. Naturalist 87: 251 (1970) | 2:304 |
| | <i>Olearia viscosa</i> (Labill.) Benth., Fl. Austral. 3: 468 (1867) | 2:302 |
| i | <i>Onopordum acanthium</i> L., Sp. Pl. 2: 827 (1753) | 2:377 |
| i x | <i>Onopordum acaulon</i> L., Sp. Pl., ed. 2, 2: 1159 (1763) | |
| | <i>Osteospermum fruticosum</i> (L.) Norl. = <i>Dimorphotheca fruticosa</i> | |
| e | <i>Ozothamnus antennaria</i> (DC.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 203 (1856) | 2:336 |
| | <i>Ozothamnus argophyllus</i> (A.Cunn. ex DC.) Anderb., Opera Bot. 104: 89 (1991) | 2:335 |
| | <i>Ozothamnus backhousei</i> Hook.f. nom. illeg. = <i>Ozothamnus rodwayi</i> | |
| n | <i>Ozothamnus bracteolatus</i> Hook.f. = <i>Ozothamnus gunnii</i> | 2:333 |
| e | <i>Ozothamnus buchananii</i> Puttock ex de Salas & Schmidt-Leb., Phytotaxa 358: 134–135 (2018) | |
| | <i>Ozothamnus cinereus</i> (Labill.) Sw. = <i>Ozothamnus turbinatus</i> | 2:333 |
| | <i>Ozothamnus cinereus</i> (Labill.) Sw. var. <i>bracteolatus</i> (Hook.f.) Rodway = <i>Ozothamnus gunnii</i> | |
| e | <i>Ozothamnus costatifructus</i> (R.V.Sm.) Anderb., Opera Bot. 104: 89 (1991) | 2:332 |
| e | <i>Ozothamnus ericifolius</i> Hook.f., London J. Bot. 6: 119 (1847) | 2:334 |
| e | <i>Ozothamnus × expansifolius</i> (P.Morris & J.H.Willis) Anderb., Opera Bot. 104: 89 (1991) | 2:337 |
| | <i>Ozothamnus ferrugineus</i> (Labill.) Sweet, Hort. Brit. [Sweet] 1: 221 (1826) | 2:335 |

| | | |
|-----|---|-------|
| | Ozothamnus ferrugineus DC. var. gravesii Rodway = Ozothamnus argophyllus | |
| e | Ozothamnus floribundus de Salas & Schmidt-Leb., Phytotaxa 358: 135 (2018) | |
| n e | Ozothamnus gunnii Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 205 (1856) | 2:333 |
| e | Ozothamnus hookeri Sond., Linnaea 25: 509 (1853) | 2:337 |
| e | Ozothamnus ledifolius (A.Cunn. ex DC.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 204 (1856) | 2:334 |
| e | Ozothamnus lycopodioides Hook.f., London J. Bot. 6: 119 (1847) | 2:338 |
| | Ozothamnus obcordatus DC., Prodr. [A. P. de Candolle] 6: 165 (1838) | 2:336 |
| e | Ozothamnus purpurascens DC., Prodr. [A. P. de Candolle] 6: 165 (1838) | |
| | Ozothamnus purpurascens DC. sensu de Salas & Baker (2017) and earlier authors = Ozothamnus reflexus (misapplied in Tasmania) | 2:334 |
| e | Ozothamnus reflexifolius Leeson & Rozefelds, Austral Syst. Bot. 16: 319 (2003) | |
| e | Ozothamnus reflexus (N.T.Burb.) de Salas & Schmidt-Leb., Phytotaxa 358: 133 (2018) | 2:334 |
| e | Ozothamnus reticulatus (Labill.) DC., Prodr. [A. P. de Candolle] 6: 164 (1838) | 2:332 |
| e | Ozothamnus rodwayi Orchard var. kingii (W.M.Curtis) P.S.Short, Muelleria 7: 522 (1992) | 2:337 |
| e | Ozothamnus rodwayi Orchard var. oreophilus (W.M.Curtis) P.S.Short, Muelleria 7: 522 (1992) | 2:337 |
| e | Ozothamnus rodwayi Orchard var. rodwayi , Muelleria 7: 522 (1992) | 2:337 |
| | Ozothamnus rosmarinifolius (Labill.) Sweet, Hort. Brit. [Sweet] 1: 221 (1826) | 2:335 |
| | Ozothamnus rosmarinifolius (Labill.) Sweet var. ericifolius (Hook.f.) Rodway = Ozothamnus ericifolius | |
| e | Ozothamnus scutellifolius Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 202 (1856) | 2:338 |
| e x | Ozothamnus selaginoides Sond. & F.Muell., Linnaea 25: 510 (1853) | 2:339 |
| | Ozothamnus thyrsoides DC., Prodr. [A. P. de Candolle] 6: 165 (1838) | 2:335 |
| | Ozothamnus turbinatus DC., Prodr. [A. P. de Candolle] 6: 164 (1838) | 2:333 |
| | Pappochroma bellidioides (Hook.f.) G.L.Nesom, Phytologia 85: 277 (1998) | |
| e | Pappochroma gunnii (Hook.f.) G.L.Nesom, Phytologia 76: 426 (1994) | 2:312 |
| e | Pappochroma pappocromum (Labill.) G.L.Nesom, Phytologia 85: 278 (1998) | 2:311 |
| e | Pappochroma stellatum (Hook.f.) G.L.Nesom, Phytologia 85: 278 (1998) | 2:312 |
| | Pappochroma tasmanicum (Hook.f.) G.L.Nesom, Phytologia 76: 426 (1994) | 2:312 |
| e | Pappochroma trigonum (S.J.Forbes & D.I.Morris) G.L.Nesom, Phytologia 85: 279 (1998) | |
| | Paquerina graminea (Labill.) Cass. ex Less. = Brachyscome graminea | 2:295 |
| | Petasites fragrans (Vill.) C.Presl = Petasites pyrenaicus | 2:360 |
| i | Petasites pyrenaicus (L.) G.López, Anales Jard. Bot. Madrid 42: 343 (1986) | 2:360 |
| | Picris angustifolia DC. subsp. angustifolia , Prodr. [A. P. de Candolle] 7: 130 (1838) | 2:383 |
| | Picris angustifolia DC. subsp. merxmulleri Lack & S.Holzappel, Willdenowia 23: 190 (1993) | 2:383 |
| | Picris echioides L. = Helminthotheca echioides | 2:383 |
| | Picris hieracioides L. sensu Curtis (1963) = Picris angustifolia (misapplied in Tasmania) | 2:383 |
| i | Pilosella aurantiaca (L.) F.W.Schultz & Sch.Bip. subsp. aurantiaca , Flora 42: 426 (1862) | 2:385 |
| i x | Pilosella officinarum Vaill. subsp. officinarum , Königl. Akad. Wiss. Paris Anat. Abh. 5: 703 (1754) | |
| | Podolepis acuminata R.Br. = Podolepis jaceoides | |
| | Podolepis decipiens Jeanes, Muelleria 33: 26 (2015) | |
| | Podolepis jaceoides (Sims) Voss, Vilm. Blumengärtn., ed. 3, 1: 537 (1894) | 2:348 |
| | Podosperma angustifolia Labill. = Podotheca angustifolia | 2:341 |
| ?x | Podotheca angustifolia (Labill.) Less., Syn. Gen. Compos. 272 (1832) | 2:341 |

| | | |
|---|---|-------|
| | Pseudognaphalium luteoalbum (L.) Hilliard & B.L.Burtt = Helichrysum luteoalbum | 2:318 |
| e | Pterygopappus lawrencei Hook.f., London J. Bot. 6: 120 (1847) | 2:316 |
| | Raoulia catipes (DC.) Hook.f. = Ewartia catipes | |
| | Raoulia meredithae (F.Muell.) Rodway = Ewartia meredithae | |
| | Raoulia planchonii (Hook.f.) Benth. = Ewartia planchonii | |
| | Rhagadiolus hedypnois (L.) All. = Leontodon rhagadioloides | |
| | Rhodanthe anthemoides (Sieber ex Spreng.) Paul G.Wilson, Nuytsia 8: 386 (1992) | 2:324 |
| | Richea glauca Labill. = Craspedia glauca | |
| | Rutidosis multiflora (Nees) B.L.Rob. = Siloxerus multiflorus | 2:342 |
| | Rutidosis pumilo Benth. = Siloxerus multiflorus | |
| | Scleroleima forsteroides Hook.f. = Abrotanella forsteroides | |
| i | Scorzonera laciniata L. var. laciniata , Sp. Pl. 2: 791 (1753) | |
| e | Senecio albogilvus I.Thomps., Muelleria 20: 130 (2004) | 2:364 |
| i | Senecio angulatus L.f., Suppl. Pl. 369 (1782) | |
| | Senecio australis Willd. sensu Rodway (1903) = Senecio linearifolius (misapplied in Tasmania) | |
| | Senecio biserratus Belcher, Ann. Missouri Bot. Gard. 43: 43 (1956) | 2:368 |
| | Senecio campylocarpus I.Thomps., Muelleria 20: 139 (2004) | |
| | Senecio capillifolius Hook.f. = Senecio pinnatifolius var. capillifolius | 2:365 |
| | Senecio centropappus F.Muell. = Centropappus brunonis | 2:365 |
| i | Senecio elegans L., Sp. Pl. 2: 869 (1753) | 2:366 |
| | Senecio extensus I.Thomps., Muelleria 19: 150 (2004) | |
| x | Senecio georgianus DC., Prodr. [A. P. de Candolle] 6: 371 (1838) | |
| | Senecio glandulosus (A.Cunn. ex DC.) Sch.Bip. = Senecio campylocarpus | |
| | Senecio glomeratus Desf. ex Poir. subsp. glomeratus , Encycl. (Lamarck) Suppl. 5: 130 (1817) | 2:370 |
| | Senecio glomeratus Desf. ex Poir. subsp. longifructus I.Thomps., Muelleria 19: 148 (2004) | 2:370 |
| | Senecio gunnii (Hook.f.) Belcher, Ann. Missouri Bot. Gard. 43: 60 (1956) | 2:370 |
| | Senecio hispidissimus I.Thomps., Muelleria 19: 138 (2004) | |
| | Senecio hispidulus A.Rich., Voy. Astrolabe 2: 94 t.34 (1834) | 2:370 |
| i | Senecio jacobaea L., Sp. Pl. 2: 870 (1753) | 2:366 |
| | Senecio lautus G.Forst. ex Willd. sensu Curtis (1963) = Senecio pinnatifolius (sensu lato, misapplied in Tasmania) | 2:365 |
| e | Senecio leptocarpus DC., Prodr. [A. P. de Candolle] 6: 372 (1838) | 2:364 |
| | Senecio linearifolius A.Rich. var. arachnoideus I.Thomps., Muelleria 20: 98 (2004) | 2:366 |
| | Senecio linearifolius A.Rich. var. denticulatus I.Thomps., Muelleria 20: 93 (2004) | 2:366 |
| | Senecio linearifolius A.Rich. var. latifolius I.Thomps., Muelleria 20: 96 (2004) | 2:366 |
| | Senecio linearifolius A.Rich. var. linearifolius , Voy. Astrolabe 2: 129 (1834) | 2:366 |
| | Senecio longipilus I.Thomps., Muelleria 19: 193 (2004) | |
| x | Senecio macrocarpus F.Muell. ex Belcher, Muelleria 5: 119 (1983) | |
| | Senecio microbasis I.Thomps., Muelleria 19: 175 (2004) | |
| | Senecio mikanioides Otto ex Harv. = Delairea odorata | 2:367 |
| | Senecio minimus Poir., Encycl. (Lamarck) Suppl. 5: 130 (1817) | 2:368 |
| | Senecio odoratus Hornem., Enum. Pl. Hort. Hafn. Suppl. [1809] (1809) | 2:367 |
| | Senecio × orarius J.M.Black, Trans. & Proc. Roy. Soc. South Australia 52: 230 (1928) | |

| | | |
|------|--|-------|
| e | Senecio papillosus F.Muell., Trans. & Proc. Philos. Inst. Victoria 2: 69 (1857) | 2:363 |
| | Senecio pectinatus DC. var. leptocarpus = Senecio leptocarpus | |
| | Senecio pectinatus DC. var. major F.Muell. ex Belcher recorded in error | |
| | Senecio pectinatus DC. var. ochroleucus F.Muell. = Senecio albogilvus | 2:364 |
| e | Senecio pectinatus DC. var. pectinatus , Prodr. [A. P. de Candolle] 6: 372 (1838) | 2:364 |
| | Senecio pectinatus DC. var. pleiocephalus Benth. sensu Rodway (1903) = Senecio pinnatifolius var. alpinus (misapplied in Tasmania) | |
| | Senecio phelleus I.Thomps., Muelleria 19: 171 (2004) | |
| | Senecio pinnatifolius A.Rich. var. alpinus (Ali) I.Thomps., Muelleria 21: 53 (2005) | 2:365 |
| e | Senecio pinnatifolius A.Rich. var. capillifolius (Hook.f.) I.Thomps., Muelleria 21: 51 (2005) | 2:365 |
| | Senecio pinnatifolius A.Rich. var. lanceolatus (Benth.) I.Thomps., Muelleria 21: 49 (2005) | 2:365 |
| | Senecio pinnatifolius A.Rich. var. maritimus (Ali) I.Thomps., Muelleria 21: 54 (2005) | 2:365 |
| | Senecio pinnatifolius A.Rich. var. pinnatifolius , Voy. Astrolabe 2: 117 (1834) | 2:365 |
| | Senecio pinnatifolius A.Rich. var. pleiocephalus (Rodway) Belcher = Senecio pinnatifolius var. alpinus | |
| | Senecio prenanthoides A.Rich., Voy. Astrolabe 2: 96 (1834) | |
| e | Senecio primulaefolius F.Muell., Trans. & Proc. Philos. Inst. Victoria 2: 69 (1857) | 2:364 |
| | Senecio psilocarpus Belcher & Albr., Muelleria 8: 113 (1994) | |
| | Senecio quadridentatus Labill., Nov. Holl. Pl. 2: 48 t.194 (1806) | 2:369 |
| e | Senecio spathulatus A.Rich. var. spathulatus , Voy. Astrolabe 2: 125 (1834) | 2:364 |
| | Senecio squarrosus A.Rich., Voy. Astrolabe 2: 107 t.35 (1834) | 2:369 |
| e x | Senecio tasmanicus I.Thomps., Muelleria 19: 158 (2004) | |
| | Senecio vagus F.Muell. subsp. vagus , Trans. & Proc. Philos. Soc. Victoria 1: 46 (1855) | |
| | Senecio velleioides A.Cunn. ex DC., Prodr. [A. P. de Candolle] 6: 374 (1838) | 2:365 |
| i | Senecio vulgaris L., Sp. Pl. 2: 867 (1753) | 2:367 |
| ?i # | Sigesbeckia orientalis L., Sp. Pl. 2: 900 (1753) | |
| | Siloxerus multiflorus Nees, Pl. Preiss. [J.G.C.Lehman] 2: 244 (1845) | 2:342 |
| i | Silybum marianum (L.) Gaertn., Fruct. Sem. Pl. 2: 378 t.162 (1791) | 2:378 |
| | Skirrhophorus eriocephalus Hook.f. ex A.Gray = Angianthus preissianus | |
| | Solenogyne bellioides Cass. var. gunnii (Hook.f.) G.L.Davis = Solenogyne gunnii | 2:292 |
| | Solenogyne dominii L.G.Adams, Brunonia 2: 58 (1979) | |
| | Solenogyne gunnii (Hook.f.) Cabrera, Blumea 14: 307 (1966) | 2:292 |
| i | Soliva sessilis Ruiz & Pav., Syst. Veg. Fl. Peruv. Chil. 225 (1798) | 2:358 |
| i | Soliva valdiviana Phil., Linnaea 33: 168 (1864) | |
| i | Sonchus asper (L.) Hill, Herb. Brit. 1: 47 t.34 (1769) | 2:390 |
| | Sonchus asper (L.) Hill subsp. glaucescens (Jord.) Ball = Sonchus asper | |
| | Sonchus hydrophilus Boulos, Suppl. Black's Fl. S. Austral. 331 (1965) | |
| | Sonchus megalocarpus (Hook.f.) J.M.Black = Actites megalocarpus | 2:390 |
| i | Sonchus oleraceus L., Sp. Pl. 2: 794 (1753) | 2:389 |
| i | Symphotrichum novi-belgii (L.) G.L.Nesom, Phytologia 77: 287 (1995) (as "1994") | |
| i | Symphotrichum subulatum (Michx.) G.L.Nesom, Phytologia 77: 293 (1995) (as "1994") | 2:299 |
| i t | Tanacetum cinerariifolium (Trevir.) Sch.Bip., Tanaceteeen: 58 (1844) | |
| i | Tanacetum parthenium (L.) Sch.Bip., Tanaceteeen: 55 (1844) | 2:353 |
| i | Tanacetum vulgare L., Sp. Pl. 2: 844 (1753) | 2:360 |

| | | |
|-----|---|---------------|
| | Taraxacum aristum G.E.Haglund & Markl., Bot. Not. 117: 197 (1964) | |
| | Taraxacum cygnorum Hand.-Mazz., Monogr. Taraxacum: 55 (1907) | |
| | Taraxacum dens-leonis Desf. sensu Rodway (1903) = Taraxacum officinale (misapplied in Tasmania) | |
| | Taraxacum kok-saghyz L.E.Rodin previously listed as naturalised but insufficient evidence exists to support this | 2:387 |
| i | Taraxacum officinale F.H.Wigg., Prim. Fl. Holsat. 56 (1780) | 2:387 |
| i | Tolpis barbata (L.) Gaertn., Fruct. Sem. Pl. 2: 372 (1791) | 2:381 |
| | Tolpis umbellata Bertol. = Tolpis barbata | 2:381 |
| i | Tragopogon porrifolius L. subsp. porrifolius , Sp. Pl. 2: 789 (1753) | 2:390 |
| | Tripleurospermum inodorum (L.) Sch.Bip. = Tripleurospermum maritimum subsp. inodorum | |
| i | Tripleurospermum maritimum (L.) W.D.J.Koch subsp. inodorum (L.) Appleg., Taxon 51: 760 (2002) | |
| | Triptilodiscus pygmaeus Turcz., Bull. Soc. Imp. Naturalistes Moscou 24: 66 (1851) | |
| i t | Urospermum dalechampii (L.) F.W.Schmidt, Samml. Phys.-Oekon. Aufsätze 1: 275 (1795) | 2:391 |
| i | Vellereophyton dealbatum (Thunb.) Hilliard & B.L.Burt, Bot. J. Linn. Soc. 82: 210 (1981) | 2:318 |
| | Vittadinia australasica (Turcz.) N.T.Burb. var. oricola N.T.Burb., Brunonia 5: 44 (1982) | |
| | Vittadinia australis A.Rich. sensu Benth (1867) misapplied to V. australasica and V. cuneata | |
| | Vittadinia australis A.Rich. var. megacephala F.Muell. ex Benth. = Vittadinia megacephala | |
| e | Vittadinia burbridgeae A.M.Gray & Rozefelds, Kanunnah 1: 9 (2005) | |
| | Vittadinia cuneata DC. var. cuneata , Prodr. [A. P. de Candolle] 5: 281 (1836) | 2:313 |
| | Vittadinia gracilis (Hook.f.) N.T.Burb., Brunonia 5: 54 (1982) | 2:313 |
| | Vittadinia megacephala (F.Muell. ex Benth.) J.M.Black recorded in error | 2:314 |
| | Vittadinia muelleri N.T.Burb., Proc. Linn. Soc. New South Wales 93: 440 fig.1D (1969) | |
| | Vittadinia scabra DC. sensu Hooker (1860) = Vittadinia cuneata (misapplied in Tasmania) | |
| | Vittadinia triloba (Gaudich.) DC. sensu Curtis (1963) = Vittadinia cuneata (misapplied in Tasmania) | 2:313 |
| | Vittadinia triloba (Gaudich.) DC. var. lanuginosa J.M.Black sensu Curtis (1963) = Vittadinia gracilis (misapplied in Tasmania) | 2:313 |
| | Vittadinia triloba (Gaudich.) DC. var. megacephala (F.Muell. ex Benth.) Ewart = Vittadinia megacephala | 2:314 |
| i | Xanthium spinosum L., Sp. Pl. 2: 987 (1753) | 2:349 |
| e | Xerochrysum alpinum Paul G.Wilson, Nuytsia 28: 36 (2017) | |
| | Xerochrysum bicolor (Lindl.) R.J.Bayer, Kew Bull. 56: 1014 (2001) | 2:331 |
| | Xerochrysum bracteatum (Vent.) Tzvelev, Novosti Sist. Vyssh. Rast. 27: 151 (1990) | |
| e | Xerochrysum collierianum A.M.Buchanan, Muelleria 20: 49 (2004) | |
| e | Xerochrysum milliganii (Hook.f.) Paul G.Wilson, Taxon 64: 106 (2015) | 2:330 |
| | Xerochrysum palustre (Flann) R.J.Bayer, Kew Bull. 56: 1014 (2001) | |
| | Xerochrysum papillosum (Labill.) R.J.Bayer, Kew Bull. 56: 1015 (2001) | 2:331 |
| | Xerochrysum subundulatum (Sch.Bip.) R.J.Bayer, Kew Bull. 56: 1015 (2001) | 2:330 |
| i | Xerochrysum viscosum (Sieber ex DC.) R.J.Bayer, Kew Bulletin 56: 1015 (2001) | |
| | BASELLACEAE | FTO 99 |
| i * | Anredera cordifolia (Ten.) Steenis, Fl. Malesiana 5: 303 fig.2 a-j (1957) | 3:580 |
| | BERBERIDACEAE | FTO 46 |
| i * | Berberis aquifolium Pursh, Fl. Amer. Sept. (Pursh) 1: 219 (1813) | |
| i | Berberis darwinii Hook., Icon. Pl. 7: t.672 (1844) | |

BETULACEAE

Alnus cordata (Loisel.) Duby previously listed as naturalised but insufficient evidence exists to support this 3:645

i * *Alnus glutinosa* (L.) Gaertn., Fruct. Sem. Pl. 2: 54 (1791)

i ***Betula pendula*** Roth, Tent. Fl. Germ. 1: 405 (1788)

BIGNONIACEAE

FTO 121

Pandorea pandorana (Andrews) Steenis, Bull. Jard. Bot. Buitenzorg 10: 198 (1928) 3:538

BORAGINACEAE

i ***Amsinckia calycina*** (Moris) Chater, Bot. J. Linn. Soc. 64: 380 (1971) 3:489

Amsinckia hispida (Ruiz & Pav.) I.M.Johnst. = ***Amsinckia calycina*** 3:489

i ***Anchusa arvensis*** (L.) M.Bieb., Fl. Taur.-Caucas. 1: 123 (1808)

Anchusa sempervirens L. = ***Pentaglottis sempervirens*** 3:491

Austrocynoglossum latifolium (R.Br.) Popov ex R.R.Mill = ***Hackelia latifolia*** 3:488

i ***Borago officinalis*** L., Sp. Pl. 1: 137 (1753) 3:490

i ***Buglossoides arvensis*** (L.) I.M.Johnst., J. Arnold Arbor. 35: 42 (1954) 3:494

Cynoglossum australe R.Br., Prodr. Fl. Nov. Holland. 495 (1810) 3:489

Cynoglossum latifolium R.Br. = ***Hackelia latifolia*** 3:488

Cynoglossum suaveolens R.Br. = ***Hackelia suaveolens*** 3:488

i ***Echium candicans*** L.f., Suppl. Pl. 131 (1782)

Echium lycopsis L. sensu Curtis (1967) = ***Echium plantagineum*** (misapplied in Tasmania) 3:495

i ***Echium plantagineum*** L., Mant. Pl. 2: 202 (1771) 3:495

i ***Echium vulgare*** L., Sp. Pl. 1: 139 (1753) 3:495

Exarrhena suaveolens R.Br. = ***Myosotis exarrhena***

Hackelia latifolia (R.Br.) Dimon & M.A.M.Renner, Aust. Syst. Bot. 30: 121–122 (2017) 3:488

Hackelia suaveolens (R.Br.) Dimon & M.A.M.Renner, Aust. Syst. Bot. 30: 122 (2017) 3:488

Lithospermum arvense L. = ***Buglossoides arvensis*** 3:494

i t ***Lithospermum officinale*** L., Sp. Pl. 1: 132 (1753) 3:494

Lycopsis arvensis L. = ***Anchusa arvensis***

i ***Myosotis arvensis*** (L.) Hill, Veg. Syst. 7: 55 (1764)

Myosotis australis R.Br., Prodr. Fl. Nov. Holland. 495 (1810) 3:492

Myosotis caespitosa Schultz = ***Myosotis laxa*** subsp. ***caespitosa*** 3:493

i ***Myosotis discolor*** Pers., Syst. Veg., ed. 15 (J.A.Murray) 190 (1798) 3:493

Myosotis exarrhena F.Muell., Syst. Census Austral. Pl. Suppl. 4: 7 (1889) 3:492

i ***Myosotis laxa*** Lehm. subsp. ***caespitosa*** (Schultz) Hyl. ex Nordh., Norsk. Fl. (Nordhagen): 529 (1940) 3:493

i ***Myosotis scorpioides*** L., Sp. Pl. 1: 131 (1753) 3:492

Myosotis suaveolens (R.Br.) Poir. = ***Myosotis exarrhena*** 3:492

i ***Myosotis sylvatica*** Hoffm., Deutschl. Fl. (Hoffm.): 61 (1791) 3:493

i ***Nonea lutea*** (Desr.) Rchb. ex A.DC., Prodr. [A. P. de Candolle] 10: 28 (1846)

i ***Pentaglottis sempervirens*** (L.) Tausch ex L.H.Bailey, Man. Cult. Pl., ed. 2: 837 (1949) 3:491

i ***Phacelia tanacetifolia*** Benth., Edwards's Bot. Reg. 20: t.1696 (1835) 3:486

Symphytum peregrinum Ledeb. = *Symphytum* × *uplandicum* 3:490

i * *Symphytum* × *uplandicum* Nyman, Syll. Fl. Eur. 80 (1855) 3:490

BRASSICACEAE (CRUCIFERAE)

| | | |
|-----|---|------|
| | <i>Alyssum maritimum</i> (L.) Lam. = Lobularia maritima | 1:45 |
| i | Arabidopsis thaliana (L.) Heynh., Fl. Sachsen 1: 538 (1842) | 1:50 |
| x | Ballantinia antipoda (F.Muell.) E.A.Shaw, Contr. Gray Herb. 205: 161 (1974) | 1:42 |
| e | Barbarea australis Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 14 (1852) | 1:47 |
| i | Barbarea intermedia Boreau, Fl. Centre France [Boreau] 2: 48 (1840) | 1:48 |
| i | Barbarea verna (Mill.) Asch., Fl. Brandenburg 1: 36 (1860) | 1:48 |
| | <i>Barbarea vulgaris</i> W.T.Aiton sensu Bentham (1863) = Barbarea australis (partly misapplied in Tasmania) | |
| i # | Brassica × juncea (L.) Czern., Conspect. Pl. Chark. 8 (1859) | |
| i | Brassica × napus L., Sp. Pl. 2: 666 (1753) | 1:34 |
| i | Brassica nigra (L.) W.D.J.Koch, Deutschl. Fl. (Röhling), ed. 3, 4: 713 (1833) | 1:35 |
| i * | <i>Brassica oleracea</i> L., Sp. Pl. 2: 667 (1753) | |
| i | Brassica rapa L., Sp. Pl. 2: 666 (1753) | 1:34 |
| | <i>Brassica sinapistrum</i> Boiss. = Sinapis arvensis | |
| i | Brassica tournefortii Gouan, Ill. Observ. Bot. 3: 44, t.20A (1773) | |
| i | Cakile edentula (Bigelow) Hook., Fl. Bor.-Amer. (Hooker) 1: 59 (1830) | 1:38 |
| i | Cakile maritima Scop. subsp. maritima , Fl. Carniol., ed. 2, 2: 35 (1772) | 1:38 |
| | <i>Camelina alyssum</i> (Mill.) Thell. sensu Curtis (1975) = Camelina sativa (misapplied in Tasmania) | 1:51 |
| i | Camelina sativa (L.) Crantz, Stirp. Austr. Fasc.1: 17 (1762) | 1:51 |
| | <i>Capsella antipoda</i> F.Muell. = Ballantinia antipoda | |
| | <i>Capsella australis</i> (Hook.f.) Benth. = Ballantinia antipoda | |
| i | Capsella bursa-pastoris (L.) Medik., Pfl.-Gatt. 1: 85 (1792) | 1:43 |
| | <i>Capsella elliptica</i> C.A.Mey. nom. illeg. = Hornungia procumbens | |
| | <i>Capsella procumbens</i> (L.) Fr. = Hornungia procumbens | |
| | <i>Capsella tasmanica</i> (Hook.) F.Muell. ex B.D.Jacks. = Hutchinsia tasmanica | 1:43 |
| | Cardamine astoniae I.Thomps., Muelleria 9: 156 (1996) | |
| | <i>Cardamine</i> aff. <i>corymbosa</i> Hook.f. sensu Hewson (1982) = Cardamine papillata | |
| | <i>Cardamine dictyosperma</i> Hook. = Rorippa dictyosperma | |
| i | Cardamine flexuosa With., Arr. Brit. Pl., ed. 3, 3: 578 (1796) | |
| | Cardamine franklinensis I.Thomps., Muelleria 9: 152 (1996) | |
| | Cardamine gunnii Hewson, Fl. Australia 8: 390 (1982) | 1:46 |
| | <i>Cardamine heterophylla</i> Hook. = Cardamine gunnii | 1:46 |
| i | Cardamine hirsuta L., Sp. Pl. 2: 655 (1753) | 1:46 |
| | <i>Cardamine hirsuta</i> L. var. <i>parvifolia</i> Rodway a name of uncertain application | |
| | <i>Cardamine hirsuta</i> L. var. <i>tenuifolia</i> (Hook.) Rodway = Cardamine tenuifolia | |
| | <i>Cardamine intermedia</i> Hook. sensu Curtis (1956) = Cardamine lilacina | |
| | Cardamine lilacina Hook., Companion Bot. Mag. 1: 273 (1836) | |
| | Cardamine papillata I.Thomps., Muelleria 9: 161 (1996) | |
| | Cardamine paucijuga Turcz., Bull. Soc. Imp. Naturalistes Moscou 27(2): 295 (1854) | |
| | <i>Cardamine pratensis</i> L. sensu Hooker (1860) = Cardamine tenuifolia (misapplied in Tasmania) | |
| | <i>Cardamine pratensis</i> L. var. <i>lilacina</i> (Hook.) Hook.f. = Cardamine lilacina | |
| | <i>Cardamine radicata</i> Hook.f. = Pachycladon radicans | |

| | | |
|-----|--|------|
| | Cardamine stylosa DC. = Rorippa gigantea | |
| | Cardamine tenuifolia Hook., J. Bot. (Hooker) 1: 247 (1837) | 1:46 |
| | Cardamine tryssa I.Thomps., Muelleria 18: 27 (2004) | |
| | Cardaria draba (L.) Desv. = Lepidium draba | 1:42 |
| | Carrichtera annua (L.) DC. previously listed as naturalised but insufficient evidence exists to support this | |
| | Cheesemaniania radicata (Hook.f.) O.E.Schulz = Pachycladon radicans | 1:47 |
| | Coronopus didymus (L.) Sm. = Lepidium didymum | 1:41 |
| | Coronopus squamatus (Forssk.) Asch. = Lepidium coronopus | 1:41 |
| | Cuphonotus antipodus (F.Muell.) J.M.Black = Ballantinia antipoda | 1:42 |
| i | Diplotaxis muralis (L.) DC., Syst. Nat. [Candolle] 2: 634 (1821) | 1:36 |
| i | Diplotaxis tenuifolia (L.) DC., Syst. Nat. [Candolle] 2: 632 (1821) | 1:36 |
| | Draba muralis L. sensu Curtis & Morris (1975) and earlier = Draba nemorosa (misapplied in Tasmania) | 1:45 |
| i | Draba muralis L., Sp. Pl. 2: 642 (1753) | 1:45 |
| i t | Draba nemorosa L., Sp. Pl. 2: 643 (1753) | 1:45 |
| i | Erophila verna (L.) Chevall. subsp. praecox (Steven) Walters, Feddes Repert. Spec. Nov. Regni Veg. 69: 57 (1964) | 1:45 |
| i | Erophila verna (L.) Chevall. subsp. verna , Fl. Gén. Env. Paris 2: 898 (1827) | 1:45 |
| | Eruca sativa Mill. previously listed as naturalised but insufficient evidence exists to support this | |
| | Eruca vesicaria (L.) Cav. subsp. sativa (Mill.) Thell. = Eruca sativa | |
| | Geococcus pusillus J.Drumm. ex Harv., Hooker's J. Bot. Kew Gard. Misc. 7: 52 (1855) | 1:50 |
| i | Hirschfeldia incana (L.) Lagr.-Foss., Fl. Tarn Garonne: 19 (1847) | 1:35 |
| i | Hornungia procumbens (L.) Hayek, Repert. Spec. Nov. Regni Veg. Beih. 30: 480 (1925) | 1:44 |
| | Hutchinsia australis Hook.f. = Ballantinia antipoda | |
| | Hutchinsia tasmanica Hook. a name of uncertain application | |
| | Hymenolobus procumbens (L.) Nutt. = Hornungia procumbens | 1:44 |
| i | Lepidium africanum (Burm.f.) DC., Syst. Nat. [Candolle] 2: 522 (1821) | |
| i | Lepidium bonariense L., Sp. Pl. 2: 645 (1753) | |
| i | Lepidium campestre (L.) W.T.Aiton, Hortus Kew. (W.T.Aiton), ed. 2, 4: 88 (1812) | 1:40 |
| i | Lepidium coronopus (L.) Al-Shehbaz, Novon 14: 156 (2004) | 1:41 |
| | Lepidium cuneifolium DC. = Lepidium foliosum | |
| | Lepidium desvauxii Thell., Neue Denkschr. Allg. Schweiz. Ges. Gesamten Naturwiss. 41: 307 (1906) | 1:39 |
| i | Lepidium didymum L., Mant. Pl. 1: 92 (1767) | 1:41 |
| i | Lepidium draba L., Sp. Pl. 2: 645 (1753) | 1:42 |
| t | Lepidium flexicaule Kirk, Trans. & Proc. New Zealand Inst. 14: 380 (1882) | |
| | Lepidium foliosum Desv., J. Bot. (Desvaux) 3: 164, 180 (1815) | 1:39 |
| | Lepidium halmaturinum J.M.Black = Lepidium desvauxii | 1:39 |
| | Lepidium heterophyllum Benth. previously listed as naturalised but insufficient evidence exists to support this | 1:40 |
| | Lepidium hyssopifolium Desv., J. Bot. (Desvaux) 3: 179 (1815) | |
| i | Lepidium latifolium L., Sp. Pl. 2: 644 (1753) | 1:41 |
| | Lepidium praetervisum Domin = Lepidium desvauxii | 1:39 |

| | | |
|----------------------|---|-------|
| | Lepidium pseudotasmanicum Thell., Neue Denkschr. Allg. Schweiz. Ges. Gesamten Naturwiss. 41: 307 (1906) | 1:39 |
| | Lepidium ruderales L. sensu Bentham (1863) = Lepidium foliosum (misapplied in Tasmania) | |
| i | Lepidium sativum L., Sp. Pl. 2: 644 (1753) | 1:40 |
| | Lepidium squamatum Forssk. = Lepidium coronopus | 1:41 |
| | Lepidium tasmanicum Thell. a name of uncertain application | |
| i | Lobularia maritima (L.) Desv., J. Bot. (Desvaux) 3: 162 (1815) | 1:45 |
| i | Lunaria annua L., Sp. Pl. 2: 653 (1753) | |
| i | Matthiola incana (L.) W.T.Aiton, Hortus Kew. (W.T.Aiton), ed. 2, 4: 119 (1812) | |
| i | Nasturtium microphyllum Boenn. ex Rchb., Fl. Germ. Excurs. 2: 683 (1830) | 1:49 |
| i | Nasturtium officinale W.T.Aiton, Hortus Kew., (W.T.Aiton), ed. 2, 4: 110 (1812) | 1:49 |
| | Nasturtium palustre (L.) DC. = Rorippa palustris | |
| | Nasturtium terrestre (With.) Aiton = Rorippa palustris | |
| e | Pachycladon radicans (Hook.f.) Heenan & A.D.Mitch., New Zealand J. Bot. 40: 558 (2002) | 1:47 |
| i * | Raphanus maritimus Sm., Engl. Bot. 23: 1643 (1806) | 1:37 |
| i | Raphanus raphanistrum L., Sp. Pl. 2: 669 (1753) | 1:37 |
| i | Rapistrum rugosum (L.) All., Fl. Pedem. 1: 257 t.78 (1785) | 1:37 |
| | Rorippa dictyosperma (Hook.) L.A.S.Johnson, Contr. New South Wales Natl. Herb. 3: 97 (1962) | 1:49 |
| | Rorippa gigantea (Hook.) Garn.-Jones, New Zealand J. Bot. 16: 119 (1978) | 1:48 |
| | Rorippa islandica (Oeder) Borbás sensu Curtis & Morris (1975) = Rorippa palustris (misapplied in Tasmania) | 1:48 |
| | Rorippa microphylla (Boenn. ex Rchb.) H.Hyl. = Nasturtium microphyllum | 1:49 |
| | Rorippa nasturtium-aquaticum (L.) Hayek = Nasturtium officinale | 1:49 |
| i | Rorippa palustris (L.) Besser, Enum. Pl. [Besser]: 27 (1822) | 1:48 |
| | Rorippa stylosa (DC.) Allan = Rorippa gigantea | 1:48 |
| i # | Rorippa sylvestris (L.) Besser, Enum. Pl. [Besser]: 27 (1822) | |
| | Senebiera coronopus (L.) Poir. = Lepidium coronopus | |
| | Senebiera didyma (L.) Pers. = Lepidium didymum | |
| i | Sinapis alba L., Sp. Pl. 2: 668 (1753) | |
| i | Sinapis arvensis L., Sp. Pl. 2: 668 (1753) | 1:35 |
| i | Sisymbrium irio L., Sp. Pl. 2: 659 (1753) | |
| i | Sisymbrium officinale (L.) Scop., Fl. Carniol., ed. 2, 2: 26 (1772) | 1:50 |
| i | Sisymbrium orientale L., Cent. Pl. 2: 24 (1756) | 1:50 |
| | Stenopetalum lineare R.Br. ex DC., Syst. Nat. [Candolle] 2: 513 (1821) | 1:44 |
| i t | Teesdalia nudicaulis (L.) W.T.Aiton, Hortus Kew. (W.T.Aiton), ed. 2, 4: 83 (1812) | 1:43 |
| i * | Thlaspi arvense L., Sp. Pl. 2: 646 (1753) | |
| | Thlaspi tasmanicum (Hook.) Hook.f. = <i>Hutchinsia tasmanica</i> | |
| CAMPANULACEAE | | |
| | Campanula gracilis G.Forst. var. littoralis (Labill.) R.Br. = Wahlenbergia gymnoclada | |
| | Campanula gracilis G.Forst. var. vincaeflora R.Br. = Wahlenbergia gracilis | |
| | Campanula littoralis Labill. = Wahlenbergia gymnoclada | |
| i * | Campanula rapunculoides L., Sp. Pl. 1: 165 (1753) | 2:410 |

| | | |
|------------------------|--|-------|
| | Isotoma fluviatilis (R.Br.) F.Muell. ex Benth. subsp. australis McComb, Contr. New South Wales Natl. Herb. 4: 109 (1970) | 2:411 |
| | Lobelia alata Labill. = Lobelia anceps | 2:414 |
| | Lobelia anceps L.f., Suppl. Pl. 395 (1782) | 2:414 |
| | Lobelia browniana Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 5: 71 (1819) | |
| | Lobelia cuneiformis Labill. = Lobelia anceps | |
| | Lobelia dentata Cav., Icon. Pl. 6: 14, t. 522 (1800) | |
| i # | Lobelia erinus L., Sp. Pl. 2: 932 (1753) | |
| | Lobelia fluviatilis R.Br. = Isotoma fluviatilis | |
| | Lobelia gibbosa Labill., Nov. Holl. Pl. 1: 50, t.71 (1805) | 2:414 |
| | Lobelia gibbosa Labill. var. browniana (Schult.) F.M.Bailey = Lobelia browniana | |
| | Lobelia irrigua R.Br., Prodr. Fl. Nov. Holland. 563 (1810) | 2:412 |
| | Lobelia pedunculata R.Br., Prodr. Fl. Nov. Holland. 563 (1810) | 2:413 |
| | Lobelia platycalyx (F.Muell.) F.Muell. = Lobelia irrigua | |
| | Lobelia pratioides Benth., Fl. Austral. 4: 131 (1868) | 2:415 |
| | Lobelia rhombifolia de Vriese, Pl. Preiss. [J.G.C.Lehman] 1: 397 (1845) | 2:414 |
| | Lobelia simplicicaulis R.Br., Prodr. Fl. Nov. Holland. 564 (1810) | |
| | Lobelia surrepens Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 237 (1856) | 2:412 |
| | Pratia irrigua (R.Br.) Benth. = Lobelia irrigua | 2:412 |
| | Pratia pedunculata (R.Br.) Benth. = Lobelia pedunculata | 2:413 |
| | Pratia platycalyx (F.Muell.) Benth. = Lobelia irrigua | 2:412 |
| | Pratia surrepens (Hook.f.) E.Wimm. = Lobelia surrepens | 2:412 |
| | Wahlenbergia ceracea Lothian, Victorian Nat. 72: 166 (1956) | 2:407 |
| | Wahlenbergia consimilis Lothian = Wahlenbergia stricta | 2:408 |
| | Wahlenbergia gracilentia Lothian, Proc. Linn. Soc. New South Wales 71: 217 (1947) | 2:407 |
| | Wahlenbergia gracilis (G.Forst.) A.DC., Monogr. Campan. 142 (1830) | 2:409 |
| | Wahlenbergia gymnoclada Lothian, Proc. Linn. Soc. New South Wales 71: 227 (1947) | 2:408 |
| | Wahlenbergia littoricola P.J.Sm. subsp. littoricola , Fl. S. Austral. [J.M.Black], ed. 4, 3: 1380 (1986) | |
| | Wahlenbergia multicaulis Benth., Enum. Pl. [Endlicher]: 75 (1837) | 2:409 |
| | Wahlenbergia quadrifida (R.Br.) A.DC. = Wahlenbergia gracilis | 2:409 |
| e | Wahlenbergia saxicola (R.Br.) A.DC., Monogr. Campan. 144 (1830) | 2:410 |
| | Wahlenbergia stricta (R.Br.) Sweet subsp. stricta , Hort. Brit. [Sweet], ed. 2: 593 (1830) | 2:408 |
| | Wahlenbergia tadgellii Lothian = Wahlenbergia multicaulis | 2:409 |
| CANNABACEAE | | |
| i | Humulus lupulus L., Sp. Pl. 2: 1028 (1753) | |
| CAPRIFOLIACEAE | | |
| i | Leycesteria formosa Wall., Fl. Ind. (Carey & Wallich ed.) 2: 182 (1824) | 2:266 |
| i | Lonicera japonica Thunb., Syst. Veg., ed. 14 (J.A.Murray) 216 (1784) | |
| i # | Lonicera periclymenum L., Sp. Pl. 1: 173 (1753) | 2:266 |
| CARYOPHYLLACEAE | | |
| i | Agrostemma githago L., Sp. Pl. 1: 435 (1753) | 1:69 |
| i | Arenaria leptoclados (Rchb.) Guss., Fl. Sicul. Syn. 2: 824 (1845) | |
| i | Arenaria serpyllifolia L., Sp. Pl. 1: 423 (1753) | 1:72 |

- i **Cerastium balearicum** F.Herm., Verh. Bot. Vereins Prov. Brandenburg 54: 247 (1913)
Cerastium fontanum Baumg. subsp. *vulgare* (Hartm.) Greuter & Burdet = **Cerastium vulgare** 1:69
- i **Cerastium glomeratum** Thuill., Fl. Env. Paris, ed. 2: 226 (1799) 1:69
- i **Cerastium semidecandrum** L., Sp. Pl. 1: 438 (1753)
Cerastium viscosum L. = **Cerastium vulgare**
- i **Cerastium vulgare** Hartm., Hand. Skand. Fl. 182 (1820) 1:69
Cerastium vulgatum L. sensu Curtis (1956) = **Cerastium vulgare** (misapplied in Tasmania)
- Colobanthus affinis** (Hook.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 45 (1855) 1:74
- Colobanthus apetalus** (Labill.) Druce var. *apetalus*, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 616 (1917) 74
Colobanthus billardierei Fenzl nom. illeg. = **Colobanthus apetalus** var. *apetalus*
- Colobanthus curtisiae** J.G.West, Aspects of Tasmanian Botany: 75 (1991)
- Colobanthus pulvinatus** F.Muell., Trans. Philos. Soc. Victoria 1: 101 (1855)
- i **Dianthus armeria** L., Sp. Pl. 1: 410 (1753)
Githago segetum Link = **Agrostemma githago**
- Gypsophila australis* (Schltld.) A.Gray = **Gypsophila tubulosa** 1:66
- i **Gypsophila tubulosa** (Jaub. & Spach) Boiss., Diagn. Pl. Orient., Ser. 1, 1: 11 (1842) 1:66
Kohlruschia prolifera (L.) Kunth sensu Curtis (1956) = **Petrorhagia nanteuillii** (misapplied in Tasmania)
- Kohlruschia velutina* (Guss.) Rchb. = **Petrorhagia dubia**
- Lychnis coronaria* (L.) Desr. = **Silene coronaria**
- Melandrium album* (Mill.) Garcke = **Silene latifolia** subsp. *alba*
- Minuartia hybrida* (Vill.) Schischk. sensu Buchanan et al. (1989) = **Minuartia mediterranea** (misapplied in Tasmania)
- i **Minuartia mediterranea** (Ledeb. ex Link) K.Maly, Glasn. Zemaljsk. Muz. Bosni Hercegovini 20: 563 (1908)
Mnium biflorum J.R.Forst. & G.Forst. = **Scleranthus biflorus**
- Mnium fasciculatum* R.Br. = **Scleranthus fasciculatus**
- i **Moenchia erecta** (L.) G.Gaertn., B.Mey. & Scherb., Oekon. Fl. Wetterau 1: 219 (1799) 1:70
- i **Paronychia brasiliiana** DC., Encycl. (Lamarck) 5: 23 (1804)
- i **Petrorhagia dubia** (Raf.) G.López & Romo, Anales Jard. Bot. Madrid 45: 363 (1988) 1:66
- i **Petrorhagia nanteuillii** (Burnat) P.W.Ball & Heywood, Bull. Brit. Mus. (Nat. Hist.) Bot. 3: 164 (1964) 1:65
Petrorhagia prolifera (L.) P.W.Ball & Heywood sensu Curtis & Morris (1975) = **Petrorhagia nanteuillii** (misapplied in Tasmania) 1:65
- Petrorhagia velutina* (Guss.) P.W.Ball & Heywood = **Petrorhagia dubia** 1:66
- i **Polycarpon tetraphyllum** (L.) L., Syst. Nat., ed. 10, 2: 881 (1759) 1:76
- i **Sagina apetala** Ard., Animadv. Bot. Spec. Alt. 2: 22 (1764) 1:73
- e **Sagina diemensis** L.G.Adams, Muellera 9: 64 (1996)
- i **Sagina maritima** Don, Herb. Brit. [Don] 7: 155 (1806) 1:73
Sagina namadgi L.G.Adams, Muellera 9: 63 (1996)
- i **Sagina procumbens** L., Sp. Pl. 1: 128 (1753) 1:73
- i **Saponaria officinalis** L., Sp. Pl. 1: 408 (1753) 1:67
Saponaria tubulosa (Jaub. & Spach) F.Muell. = **Gypsophila tubulosa**
- i **Scleranthus annuus** L., Sp. Pl. 1: 406 (1753) 1:77

| | | |
|-----|---|------|
| | Scleranthus biflorus (J.R.Forst. & G.Forst.) Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 74 (1852) | 1:76 |
| | Scleranthus brockiei P.A.Will., Rec. Domin. Mus. 3: 16 (1956) | |
| | Scleranthus diander R.Br., Prodr. Fl. Nov. Holland. 412 (1810) | 1:77 |
| | Scleranthus fasciculatus (R.Br.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 42 (1855) | |
| | <i>Silene alba</i> (Mill.) E.H.L.Krause = Silene latifolia subsp. alba | 1:67 |
| | <i>Silene colorata</i> Poir. previously listed as naturalised but insufficient evidence exists to support this | |
| | <i>Silene conica</i> L. previously listed as naturalised but insufficient evidence exists to support this | |
| i | Silene coronaria (L.) Clairv., Man. Herbor. Suisse 145 (1811) | |
| i # | Silene dichotoma Ehrh., Beitr. Naturk. [Ehrhart] 7: 144 (1792) | |
| i | Silene gallica L. var. gallica , Sp. Pl. 1: 417 (1753) | 1:68 |
| i | Silene gallica L. var. quinquevulnera (L.) W.D.J.Koch, Syn. Fl. Germ. Helv. 1: 100 (1835) | 1:68 |
| i | Silene latifolia Poir. subsp. alba (Mill.) Greuter & Burdet, Willdenowia 12: 189 (1982) | 1:67 |
| | <i>Silene longicaulis</i> Pourr. ex Lag. sensu Buchanan (1995) = <i>Silene colorata</i> (misapplied in Tasmania) | |
| i | Silene nocturna L., Sp. Pl. 1: 416 (1753) | 1:68 |
| | <i>Silene pratensis</i> (Rafn) Godr. & Gren. = Silene latifolia subsp. alba | 1:67 |
| i | Silene vulgaris (Moench) Garcke, Fl. N. Mitt.-Deutschland, ed. 9: 64 (1869) | 1:68 |
| | <i>Spergula apetala</i> Labill. = Colobanthus apetalus | |
| i | Spergula arvensis L. var. arvensis , Sp. Pl. 1: 440 (1753) | |
| i | Spergula arvensis L. var. sativa (Boenn.) Mert. & W.D.J.Koch, Deutschl. Fl. (Mertens & W. D. J. Koch), ed. 3. 3: 360 (1831) | |
| i | Spergularia bocconeii (Scheele) Graebn., Syn. Mitteleur. Fl. [Ascherson & Graebner] 5: 849 (1919) | |
| | <i>Spergularia marginata</i> Kitt. sensu Curtis (1956) = Spergularia tasmanica (misapplied in Tasmania) | |
| i | Spergularia marina (L.) Griseb., Spic. Fl. Rumel. 1: 213 (1843) | 1:75 |
| | <i>Spergularia media</i> (L.) C.Presl sensu Curtis & Morris (1975) = Spergularia tasmanica (misapplied in Tasmania) | 1:75 |
| i | Spergularia rubra (L.) J.Presl & C.Presl, Fl. Cech. 94 (1819) | 1:75 |
| | <i>Spergularia rubra</i> (L.) J.Presl & C.Presl var. marina (Roth) A.Gray = Spergularia marina | |
| | <i>Spergularia salina</i> J.Presl & C.Presl = Spergularia marina | |
| | Spergularia tasmanica (Kindb.) L.G.Adams, Austral. Syst. Bot. 21: 258 (2008) | 1:75 |
| | Stellaria angustifolia Hook. subsp. angustifolia , J. Bot. (Hooker) 1: 250 (1834) | |
| | Stellaria angustifolia Hook. subsp. tenella (Benth.) C.H.Mill. & J.G.West, J. Adelaide Bot. Gard. 25: 36 (2012) | |
| | <i>Stellaria caespitosa</i> Hook.f. = Stellaria angustifolia subsp. angustifolia | |
| | Stellaria flaccida Hook., Companion Bot. Mag. 1: 275 (1836) | 1:71 |
| | <i>Stellaria glauca</i> With. sensu Hooker (1860) = Stellaria angustifolia (misapplied in Tasmania) | |
| i * | <i>Stellaria graminea</i> L., Sp. Pl. 1: 422 (1753) | 1:72 |
| i | Stellaria media (L.) Vill., Hist. Pl. Dauphiné (Villars) 3: 615 (1789) | 1:71 |
| | Stellaria multiflora Hook. subsp. multiflora , Companion Bot. Mag. 1: 275 (1836) | |
| | Stellaria multiflora Hook. subsp. nebulosa C.H.Mill. & J.G.West, J. Adelaide Bot. Gard. 25: 48 (2012) | |
| i | Stellaria pallida (Dumort.) Crép., Man. Fl. Belgique, ed. 2: 19 (1866) | 1:72 |
| | <i>Stellaria palustris</i> Retz. sensu Curtis (1956), Curtis & Morris (1975) = Stellaria angustifolia (misapplied in Tasmania) | 1:71 |
| | Stellaria pungens Brongn., Voy. Monde, Atlas: t.78 (1826) | 1:71 |

- Vaccaria hispanica* (Mill.) Rauschert previously listed as naturalised but insufficient evidence exists to support this 1:66
- Vaccaria pyramidata* Medik. = *Vaccaria hispanica* 1:66
- Vaccaria segetalis* (Neck.) Garcke ex Asch. = *Vaccaria hispanica*

CASUARINACEAE**FTO 67**

- e ***Allocasuarina crassa*** L.A.S.Johnson, Fl. Australia 3: 194 (1989)
- e ***Allocasuarina duncanii*** L.A.S.Johnson & D.I.Morris, Telopea 5: 793 (1994)
- Allocasuarina littoralis*** (Salisb.) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 76 (1982) 3:644
- e ***Allocasuarina monilifera*** (L.A.S.Johnson) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 76 (1982) 3:644
- Allocasuarina paludosa*** (Sieber ex Spreng.) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 77 (1982) 3:644
- Allocasuarina verticillata*** (Lam.) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 79 (1982) 3:644
- e ***Allocasuarina zephyrea*** L.A.S.Johnson, Fl. Australia 3: 199 (1989)
- Casuarina bicuspidata* Benth. sensu Rodway (1903) = ***Allocasuarina paludosa*** (misapplied in Tasmania) 3:644
- Casuarina distyla* Vent. sensu Rodway (1903) = ***Allocasuarina monilifera*** (misapplied in Tasmania) 3:644
- Casuarina littoralis* Salisb. = ***Allocasuarina littoralis*** 3:644
- Casuarina monilifera* L.A.S.Johnson = ***Allocasuarina monilifera*** 3:644
- Casuarina paludosa* Sieber ex Spreng. = ***Allocasuarina paludosa*** 3:644
- Casuarina quadrivalvis* Labill. = ***Allocasuarina verticillata*** 3:644
- Casuarina stricta* Dryand. = ***Allocasuarina verticillata*** 3:644
- Casuarina suberosa* Otto & A.Dietr. = ***Allocasuarina littoralis*** 3:644

CELASTRACEAE**FTO 69**

- i ***Euonymus europaeus*** L., Sp. Pl. 1: 197 (1753)
- i ***Euonymus japonicus*** Thunb., Nova Acta Regiae Soc. Sci. Upsal. 3: 208 (1780)
- i ***Maytenus magellanica*** (Lam.) Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.). 2: 254 (1845)
- Stackhousia flava* Hook. = ***Stackhousia viminea*** 1:112
- Stackhousia gunnii* Hook.f. = ***Stackhousia subterranea*** 1:112
- Stackhousia liniarifolia* A.Cunn. = ***Stackhousia monogyna***
- Stackhousia monogyna*** Labill., Nov. Holl. Pl. 1: 77 t.104 (1805) 1:112
- Stackhousia pulvinaris*** F.Muell., Trans. Philos. Soc. Victoria 1: 101 (1855) 1:112
- Stackhousia spathulata*** Sieber ex Spreng., Syst. Veg. (ed. 16) [Sprengel] 4: 124 (1827) 1:112
- Stackhousia subterranea*** W.R.Barker, J. Adelaide Bot. Gard. 21: 90 (2007) 1:112
- Stackhousia viminea*** Sm., Cycl. (Rees) 33 (1816) 1:112

CISTACEAE

- i ***Cistus creticus*** L., Sp. Pl., ed. 2: 738 (1762)
- i # ***Cistus inflatus*** Pourr. ex Demoly, Acta Bot. Gallica 144: 42 (1998)
- Cistus psilosepalus* Sweet sensu Buchanan (2004) = ***Cistus inflatus*** (misapplied in Tasmania)

CONVOLVULACEAE**FTO 111**

- Calystegia marginata*** R.Br., Prodr. Fl. Nov. Holland. 483 (1810)
- n ***Calystegia sepium*** (L.) R.Br. subsp. ***sepium***, Prodr. Fl. Nov. Holland. 483 (1810) 3:498
- n i ***Calystegia silvatica*** (Kit.) Griseb. subsp. ***silvatica***, Spic. Fl. Rumel. 1: 74 (1843) 3:498
- Calystegia soldanella*** (L.) Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 4: 184 (1819) 3:499
- Convolvulus angustissimus*** R.Br. subsp. ***angustissimus***, Prodr. Fl. Nov. Holland. 482 (1810) 3:497

| | | |
|---------------------|---|-------|
| i | Convolvulus arvensis L., Sp. Pl. 1: 153 (1753) | 3:497 |
| | Convolvulus erubescens Sims sensu Curtis (1967) = Convolvulus angustissimus subsp. angustissimus (misapplied in Tasmania) | 3:497 |
| | Convolvulus sepium L. sensu Bentham (1868) = Calystegia sepium subsp. sepium | |
| | Convolvulus soldanella L. sensu Bentham (1868) = Calystegia soldanella | |
| i | Cuscuta epithymum (L.) L., Syst. Veg., ed. 13: 140 (1774) | 3:501 |
| i x | Cuscuta suaveolens Ser., Ann. Sci. Phys. Nat. Lyon 3: 519 (1840) | |
| | Cuscuta tasmanica Engelm., Trans. Acad. Sci. St. Louis 1: 512 (1859) | 3:501 |
| | Dichondra repens J.R.Forst. & G.Forst., Char. Gen. Pl. 40: t.20 (1775) | 3:499 |
| i | Ipomoea indica (Burm.) Merr., Interpr. Herb. Amboin. 445 (1917) | |
| | Wilsonia backhousei Hook.f., London J. Bot. 6: 275 (1847) | 3:500 |
| | Wilsonia humilis R.Br., Prodr. Fl. Nov. Holland. 490 (1810) | 3:500 |
| | Wilsonia rotundifolia Hook., Icon. Pl. 5: t.160 (1842) | |
| CRASSULACEAE | | |
| i | Aeonium arboreum (L.) Webb & Berthel., Hist. Nat. Iles Canaries (Phytogr.) 1: 185 (1840) | |
| n i | Aeonium haworthii Webb & Berthel., Hist. Nat. Iles Canaries (Phytogr.) 1: 193 (1840) | |
| i | Cotyledon orbiculata L., Sp. Pl. 1: 429 (1753) | |
| | Crassula closiana (Gay) Reiche, Fl. Chile [Reiche] 2: 369 (1897) | |
| | Crassula decumbens Thunb. var. decumbens , Prodr. Pl. Cap. 1: 54 (1794) | 1:185 |
| | Crassula exserta (Reader) Ostenf., Dansk Bot. Ark. 2: 47 (1918) | 1:184 |
| | Crassula helmsii (Kirk) Cockayne, Trans. New Zealand Inst. 39: 349 (1907) | 1:185 |
| | Crassula macrantha (Hook.f.) Diels & E.Pritz. = Crassula decumbens | 1:185 |
| t | Crassula moschata G.Forst., Commentat. Soc. Regiae Sci. Gott. 9: 26 (1789) | |
| i | Crassula multicava Lem. subsp. multicava , Ill. Hort. 9: 40 (1862) | |
| i # | Crassula muscosa L. var. muscosa , Pl. Rar. Afr. 10 (1760) | |
| i | Crassula natans Thunb. var. minus (Eckl. & Zeyh.) G.D.Rowley, Cact. Succ. J. Gr. Brit. 40: 53 (1978) | |
| | Crassula pedicellosa (F.Muell.) Ostenf. = Crassula closiana | |
| | Crassula peduncularis (Sm.) F.Meigen, Bot. Jahrb Syst. 17: 239 (1893) | 1:184 |
| | Crassula sieberiana (Schult. & Schult.f.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 618 (1917) | 1:184 |
| | Crassula sieberiana (Schult. & Schult.f.) Druce subsp. tetramera Toelken = Crassula tetramera | |
| i # | Crassula tetragona L. subsp. robusta (Toelken) Toelken, J. S. African Bot. 41: 122 (1975) | |
| | Crassula tetramera (Toelken) A.P.Druce & Sykes, New Zealand J. Bot. 25: 128 (1987) | |
| i | Sedum acre L., Sp. Pl. 1: 432 (1753) | 1:186 |
| i | Sedum album L., Sp. Pl. 1: 432 (1753) | |
| i | Sedum forsterianum Sm., Engl. Bot. 26: t.1802 (1808) | |
| i | Sedum praealtum A. DC., Mém. Soc. Phys. Genève 11: 445 (1847) | |
| | Sedum reflexum L. sensu Buchanan (2007) = Sedum rupestre (misapplied in Tasmania) | |
| i | Sedum rupestre L., Sp. Pl. 1: 431 (1753) | |
| i | Sedum sediforme (Jacq.) Pau, Acta y Memorias Prim. Congr. Nat. Esp. 246 (1909) | |
| i t | Sedum sexangulare L., Sp. Pl. 1: 432 (1753) | 1:186 |
| | Tillaea exserta Reader = Crassula exserta | |
| | Tillaea macrantha Hook.f. = Crassula decumbens | |
| | Tillaea purpurata Hook.f. = Crassula peduncularis | |

- Tillaea recurva (Hook.f.) Hook.f. = **Crassula helmsii**
 Tillaea sieberiana Schult. & Schult.f. = **Crassula sieberiana**
 Tillaea verticillaris DC. = **Crassula sieberiana**

CUCURBITACEAE

FTO 65

- i **Cucumis myriocarpus** Naudin, Ann. Sci. Nat., Bot., sér. 4, 11: 22 (1859)
 i # **Ecballium elaterium** (L.) A.Rich., Dict. Class. Hist. Nat. [Bory] 6: 19 (1824)
 Sicyos angulatus L. sensu Curtis (1963) = **Sicyos australis** (misapplied in Tasmania) 2:237
Sicyos australis Endl., Prodr. Fl. Norfolk.: 67 (1833) 2:237

CUNONIACEAE

FTO 72

- e **Anodopetalum biglandulosum** (A.Cunn. ex Hook.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 148 (1856) 1:180
Bauera rubioides Andrews, Bot. Repos. 3: t.198 (1801) 1:180
 Carpodontos lucida Labill. = **Eucryphia lucida**
 Eucryphia billardierei Spach nom. illeg., nom. superfl. = **Eucryphia lucida**
 Eucryphia billardierei Spach var. milliganii (Hook.f.) Benth. nom. illeg. = **Eucryphia milliganii**
 e **Eucryphia lucida** (Labill.) Baill., Hist. Pl. (Baillon) 1: 402 (1869) 1:183
 e **Eucryphia milliganii** Hook.f. subsp. **milliganii**, Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 54 (1855) 1:183
 e **Eucryphia milliganii** Hook.f. subsp. **pubescens** R.W.Barnes, G.J.Jord., R.S.Hill & McCoull, Austral. J. Bot. 48: 488 (2000)

DILLENIACEAE

FTO 92

- Dillenia procumbens Labill. = **Hibbertia procumbens**
Hibbertia acicularis (Labill.) F.Muell., Pl. Victoria 1: 17 (1862) 1:23
 Hibbertia angustifolia (R.Br. ex DC.) Benth. = **Hibbertia procumbens**
Hibbertia appressa Toelken, J. Adelaide Bot. Gard. 19: 54 (2000)
 Hibbertia aspera DC. sensu Curtis & Morris (1975) = **Hibbertia hirticalyx** (misapplied in Tasmania) 1:23
 e **Hibbertia basaltica** A.M.Buchanan & Schah., Muelleria 22: 105 (2005)
 Hibbertia billardierei F.Muell. nom. illeg. = **Hibbertia empetrifolia**
 Hibbertia billardierei F.Muell. var. monadelpha F.Muell. ex Benth. = **Hibbertia hirticalyx**
 Hibbertia billardierei F.Muell. var. obovata R.Br. ex Benth. = **Hibbertia hirticalyx**
Hibbertia calycina (DC.) N.A.Wakef., Vict. Naturalist 72: 122 (1955)
 Hibbertia densiflora (Hook.f.) F.Muell. = **Hibbertia sericea**
Hibbertia dispar Toelken, J. Adelaide Bot. Gard. 26: 31-69 (2013)
Hibbertia empetrifolia (DC.) Hoogland subsp. **empetrifolia**, Kew Bull. 29: 155 (1974) 1:23
Hibbertia ericifolia Hook.f. subsp. **ericifolia**, Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 14 (1855) (as H. ericaefolia)
 Hibbertia fasciculata R.Br. ex DC. sensu Curtis & Morris (1975) = **Hibbertia prostrata** (misapplied in Tasmania) 1:24
Hibbertia hirsuta (Hook.) Benth., Fl. Austral. 1: 26 (1863) 1:22
Hibbertia hirticalyx Toelken, J. Adelaide Bot. Gard. 18: 146 (1998) 1:23
 Hibbertia linearis R.Br. var. obtusifolia (DC.) Benth. = **Hibbertia obtusifolia**
 x **Hibbertia obtusifolia** DC., Syst. Nat. [Candolle] 1: 429 (1817) 1:24
Hibbertia procumbens (Labill.) DC., Syst. Nat. [Candolle] 1: 427 (1817) 1:24
Hibbertia prostrata Hook., J. Bot. (Hooker) 1: 246 (1834) 1:24

| | | |
|---|--|----------------|
| | Hibbertia riparia (R.Br. ex DC.) Hoogland, Kew Bull. 29: 155 (1974) | 1:22 |
| | Hibbertia rufa N.A.Wakef., Vict. Naturalist 72: 119 (1955) | 1:23 |
| | Hibbertia sericea (R.Br. ex DC.) Benth. var. sericea , Fl. Austral. 1: 26 (1863) | 1:22 |
| | Hibbertia serpyllifolia R.Br. ex DC., Syst. Nat. [Candolle] 1: 430 (1817) | 1:24 |
| | <i>Hibbertia stricta</i> (DC.) F.Muell. = Hibbertia riparia | |
| | <i>Hibbertia stricta</i> (DC.) F.Muell. var. <i>canescens</i> = Hibbertia riparia | |
| | <i>Hibbertia stricta</i> (DC.) F.Muell. var. <i>glabriuscula</i> = Hibbertia riparia | |
| | Hibbertia virgata R.Br. ex DC., Syst. Nat. [Candolle] 1: 428 (1817) | 1:24 |
| | <i>Pleurandra acicularis</i> Labill. = Hibbertia acicularis | |
| | <i>Pleurandra hirsuta</i> Hook. = Hibbertia hirsuta | |
| | <i>Pleurandra ovata</i> Labill. = Hibbertia empetrifolia and Hibbertia appressa | |
| | <i>Pleurandra ovata</i> Labill. var. <i>scabra</i> (R.Br. ex DC.) Hook.f. sensu Hooker(1860) = Hibbertia hirticalyx (misapplied in Tasmania) | |
| | <i>Pleurandra riparia</i> R.Br. ex DC. (and vars.) = Hibbertia riparia | |
| | <i>Pleurandra sericea</i> R.Br. ex DC. = Hibbertia sericea | |
| | DIPSACACEAE | FTO 134 |
| i | Dipsacus fullonum L., Sp. Pl. 1: 97 (1753) | 2:280 |
| | <i>Dipsacus sylvestris</i> Huds. = Dipsacus fullonum | 2:280 |
| i | Scabiosa atropurpurea L., Sp. Pl. 1: 100 (1753) | 2:280 |
| i | Scabiosa farinosa Coss., Ill. Fl. Atlant. 6: 58, t. 134. (1893) | |
| | DONATIACEAE | FTO 126 |
| t | Donatia novae-zelandiae Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 81 (1852) | 2:397 |
| | DROSERACEAE | FTO 96 |
| | Drosera arcturi Hook., J. Bot. (Hooker) 1: 247 (1834) | 1:187 |
| | Drosera auriculata Backh. ex Planch., Ann. Sci. Nat., Bot., sér. 3, 9: 295 (1848) | 1:188 |
| | Drosera binata Labill., Nov. Holl. Pl. 1: 78 t.105 (1805) | 1:188 |
| | <i>Drosera foliosa</i> Hook.f. ex Planch. nom. illeg. = Drosera hookeri | |
| | Drosera glanduligera Lehm., Nov. Stirp. Pug. [Lehmann] 8: 37 (1844) | 1:187 |
| | <i>Drosera gracilis</i> Hook.f. ex Planch. = Drosera peltata | 1:188 |
| | Drosera gunniana (Planch.) de Salas, Muelleria 36: 102 (2018) | |
| | Drosera hookeri R.P.Gibson, B.J.Conn & Conran, J. Adelaide Bot. Gard. 24: 41 (2010) | |
| | Drosera macrantha Endl., Enum. Pl. [Endlicher] 6 (1837) | 1:189 |
| | <i>Drosera macrantha</i> Endl. subsp. <i>planchonii</i> (Hook.f. ex Planch.) N.G.Marchant = Drosera macrantha | 1:189 |
| | <i>Drosera menziesii</i> R.Br. ex DC. sensu Rodway (1903) = Drosera macrantha (misapplied in Tasmania) | |
| | <i>Drosera menziesii</i> R.Br. ex DC. var. <i>albiflora</i> Benth. = Drosera macrantha | |
| e | Drosera murfetii Lowrie & Conran, J. Adelaide Bot. Gard. 27: 17 (2014) | |
| | Drosera peltata Thunb., Drosera 7 (1797) | 1:188 |
| | <i>Drosera peltata</i> Thunb. sensu Rodway and later (1903) authors = Drosera gunniana (misapplied in Tasmania) | |
| | <i>Drosera peltata</i> Thunb. subsp. <i>auriculata</i> (Backh. ex Planch.) Conn = Drosera auriculata | 1:188 |
| | <i>Drosera peltata</i> Thunb. var. <i>foliosa</i> (Hook.f. ex Planch.) Benth. = Drosera hookeri | |
| | <i>Drosera peltata</i> Thunb. var. <i>gracilis</i> (Hook.f. ex Planch.) Benth. = Drosera peltata | |
| | <i>Drosera planchonii</i> Hook.f. = Drosera macrantha | 1:189 |

| | | |
|-------|---|----------------|
| | Drosera pygmaea DC., Prodr. [A. P. de Candolle] 1: 317 (1824) | 1:187 |
| | Drosera spatulata Labill. var. spatulata , Nov. Holl. Pl. 1: 79 t.106 fig.1 (1805) | 1:187 |
| | ELAEOCARPACEAE | FTO 71 |
| e | Aristotelia peduncularis (Labill.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 52 (1855) | 1:91 |
| | Elaeocarpus cyaneus Sims = Elaeocarpus reticulatus | |
| | Elaeocarpus peduncularis Labill. = Aristotelia peduncularis | |
| | Elaeocarpus reticulatus Sm., Cycl. 12, no.6 (1809) | 1:91 |
| | Tetratheca ciliata Lindl., Three Exped. Australia [Mitchell] 2: 205 (1838) | 1:60 |
| e | Tetratheca sp. Flinders Is. (T.Rudman HO510551) Tas Herbarium | |
| e | Tetratheca sp. Freycinet Pen. (A.C.Rozefelds 323) Tas Herbarium | |
| | Tetratheca glandulosa Labill. nom. illeg. = Tetratheca labillardierei | 1:60 |
| e | Tetratheca gunnii Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 36, t.7B (1855) | |
| | Tetratheca labillardierei Joy Thomps., Telopea 1: 189 (1976) | 1:60 |
| | Tetratheca pilosa Labill. subsp. latifolia Joy Thomps., Telopea 1: 213 (1976) | |
| | Tetratheca pilosa Labill. subsp. pilosa , Nov. Holl. Pl. 1: 95, t.122 (1805) | 1:60 |
| | Tetratheca pilosa Labill. var. procumbens = Tetratheca procumbens | |
| | Tetratheca procumbens Gunn ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 35, t.7A (1855) | 1:61 |
| | ELATINACEAE | FTO 76 |
| | Elatine americana (Pursh) Arn. sensu Bentham (1863) = Elatine gratiolooides (misapplied in Tasmania) | |
| | Elatine gratiolooides A.Cunn., Ann. Nat. Hist. 4: 26 (1840) | 1:80 |
| | ERICACEAE | FTO 105 |
| | Acrothamnus hookeri (Sond.) Quinn, Austral. Syst. Bot. 18: 452 (2005) | 2:438 |
| | Acrothamnus montanus (R.Br.) Quinn, Austral. Syst. Bot. 18: 452 (2005) | 2:433 |
| | Acrotriche affinis DC., Prodr. [A. P. de Candolle] 7: 757 (1839) | |
| | Acrotriche cordata (Labill.) R.Br., Prodr. Fl. Nov. Holland. 548 (1810) | |
| | Acrotriche patula sensu Hooker (1860) = Acrotriche serrulata (misapplied in Tasmania) | |
| | Acrotriche serrulata (Labill.) R.Br., Prodr. Fl. Nov. Holland. 547 (1810) | 2:440 |
| e | Androstoma verticillata (Hook.f.) Quinn, Austral. Syst. Bot. 18: 450 (2005) | 2:439 |
| n i # | Arbutus unedo L., Sp. Pl. 1: 395 (1753) | |
| e | Archeria comberi Melville, Kew Bull. 12: 389 (1957) | 2:453 |
| e | Archeria eriocarpa Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 263 t.80B (1857) | 2:452 |
| e | Archeria hirtella (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 263 t.81 (1857) | 2:452 |
| e | Archeria minor Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 264 (1857) | 2:452 |
| e | Archeria serpyllifolia Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 263 t.80A (1857) | 2:452 |
| | Archeria serpyllifolia Hook.f. var. minor (Hook.f.) Benth. = Archeria minor | 2:452 |
| | Astroloma humifusum (Cav.) R.Br., Prodr. Fl. Nov. Holland. 538 (1810) | 2:424 |
| n | Astroloma pinifolium (R.Br.) Benth. = Stenantha pinifolia | 2:424 |
| | Brachyloma ciliatum (R.Br.) Benth., Fl. Austral. 4: 173 (1868) | 2:432 |
| | Brachyloma ciliatum (R.Br.) Benth. var. intermedium Rodway = Brachyloma ciliatum | |
| | Brachyloma daphnoides Benth. sensu Bentham (1868) mistakenly attributed to Tasmania | |
| | Brachyloma depressum (F.Muell.) Benth., Fl. Austral. 4: 173 (1868) | 2:432 |
| | Brossaea lanceolata (Hook.f.) Kuntze = Gaultheria lanceolata | |
| i t | Calluna vulgaris (L.) Hull, Brit. Fl., ed. 2, 1: 114 (1808) | 2:418 |

| | | |
|---|---|-------|
| | Cyathodes abietina (Labill.) R.Br. = Leptecophylla abietina | 2:427 |
| | Cyathodes acerosa R.Br. ex Roem. & Schult. sensu Rodway (1903) = Leptecophylla oxycedrus (misapplied in Tasmania) | |
| | Cyathodes adscendens Hook.f. = Planocarpa petiolaris | |
| | Cyathodes dealbata R.Br. = Montitega dealbata | 2:427 |
| | Cyathodes disticha Labill. = Trochocarpa disticha | |
| | Cyathodes divaricata Hook.f. = Leptecophylla divaricata | 2:428 |
| e | Cyathodes glauca Labill., Nov. Holl. Pl. 1: 57 t.81 (1805) | 2:426 |
| | Cyathodes juniperina (J.R.Forst. & G.Forst.) Druce sensu Curtis & Morris (1975) = Leptecophylla oxycedrus and L. pogonocalyx subsp. decipiens (misapplied in Tasmania) | 2:427 |
| | Cyathodes macrantha Hook.f. = Cyathodes straminea | |
| | Cyathodes nitida Jarman = Planocarpa nitida | |
| | Cyathodes oxycedrus (Labill.) R.Br. = Leptecophylla oxycedrus | |
| | Cyathodes parvifolia R.Br. = Leptecophylla parvifolia | 2:428 |
| | Cyathodes pendulosa Jarman = Leptecophylla pendulosa | |
| | Cyathodes petiolaris (DC.) Druce = Planocarpa petiolaris | 2:426 |
| e | Cyathodes platystoma C.M.Weiller, Austral. Syst. Bot. 9: 502 (1996) | |
| e | Cyathodes straminea R.Br., Prodr. Fl. Nov. Holland. 539 (1810) | 2:426 |
| | Cyathodes straminea var. macrantha (Hook.f.) Rodway = Cyathodes straminea | |
| | Cyathodes sulcata Mihaich = Planocarpa sulcata | |
| | Cystanthe acerosa (Lindl.) F.Muell. = Richea acerosa | |
| | Cystanthe dracophylla (Hook.f.) Kuntze = Richea dracophylla | |
| | Cystanthe gunnii (Hook.f.) Kuntze = Richea gunnii | |
| | Cystanthe milliganii (Hook.f.) F.Muell. = Richea milliganii | |
| | Cystanthe pandanifolia (Hook.f.) Kuntze = Richea pandanifolia | |
| | Cystanthe procera F.Muell. = Richea procera | |
| | Cystanthe scoparia (Hook.f.) Kuntze = Richea scoparia | |
| | Cystanthe sprengelioides R.Br. = Richea sprengelioides | |
| | Decaspora cunninghamii DC. = Trochocarpa cunninghamii | |
| | Decaspora disticha (Labill.) R.Br. = Trochocarpa disticha | |
| | Decaspora gunnii Hook.f. = Trochocarpa gunnii | |
| | Decaspora thymifolia R.Br. = Trochocarpa thymifolia | |
| e | Dracophyllum milliganii Hook.f., Icon. Pl. 9: t.845 (1852) | 2:461 |
| e | Dracophyllum minimum F.Muell., Fragm. (Mueller) 1: 39 (1859) | 2:461 |
| e | Epacris acuminata Benth., Fl. Austral. 4: 240 (1868) | 2:451 |
| e | Epacris apsleyensis Crowden, Pap. & Proc. Roy. Soc. Tasmania 120: 17 (1986) | |
| e | Epacris barbata Melville, Kew Bull. 7: 176 (1952) | 2:450 |
| | Epacris campanulata Lodd. ex DC. = Epacris impressa | |
| e | Epacris cerasicollina Crowden, Muellera 25: 124 (2007) | |
| | Epacris ceriflora Graham = Epacris impressa | |
| | Epacris cerinthoides Labill. = Prionotes cerinthoides | |
| e | Epacris corymbiflora Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 261 (1857) | 2:447 |
| e | Epacris curtisiae Jarman, Pap. & Proc. Roy. Soc. Tasmania 122: 115 (1988) | |

- e **Epacris exserta** R.Br., Prodr. Fl. Nov. Holland. 551 (1810) 2:448
Epacris exserta var. *virgata* (Hook.f.) Benth. = **Epacris virgata**
- e **Epacris franklinii** Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 261 (1857) 2:448
- e **Epacris glabella** Jarman, Aspects of Tasmanian Botany: 100 (1991)
- e **Epacris grandis** Crowden, Pap. & Proc. Roy. Soc. Tasmania 120: 19 (1986)
- e **Epacris graniticola** Crowden, Muellera 25: 126 (2007)
- Epacris gunnii** Hook.f., London J. Bot. 6: 272 (1847) 2:451
- e **Epacris heteronema** Labill. var. **gigantea** F.Muell., Fragm (Mueller) 4: 127 (1864)
- e **Epacris heteronema** Labill. var. **heteronema**, Nov. Holl. Pl. 1: 42 t.56 (1805)
Epacris heteronema var. *planifolia* = **Epacris heteronema** (Tasmanian material)
Epacris hirtella Hook.f. = **Archeria hirtella**
- Epacris impressa** Labill., Nov. Holl. Pl. 1: 43 t.58 (1805) 2:446
Epacris impressa f. *ceriflora* (Graham) Siebert & Voss = **Epacris impressa**
Epacris impressa f. *diemenica* Gand. = **Epacris impressa**
Epacris impressa f. *lucida* Gand. = **Epacris impressa**
Epacris impressa f. *milliganii* Gand. = **Epacris impressa**
Epacris impressa f. *ruscifolia* (R.Br.) Siebert & Voss = **Epacris impressa**
Epacris impressa var. *campanulata* (Lodd. ex DC.) Hook.f. = **Epacris impressa**
Epacris impressa var. *ceriflora* (Graham) Rodway = **Epacris impressa**
Epacris impressa var. *nivea* Hook.f. = **Epacris impressa**
Epacris impressa var. *ovata* Benth. = **Epacris impressa**
Epacris impressa var. *ruscifolia* (R.Br.) Rodway = **Epacris impressa**
Epacris impressa var. *variabilis* (Lodd. ex Paxton) Hook.f. = **Epacris impressa**
- Epacris lanuginosa** Labill., Nov. Holl. Pl. 1: 42 t.57 (1805) 2:448
- e **Epacris limbata** K.J.Williams & F.Duncan, Aspects of Tasmanian Botany: 95 (1991)
- e **Epacris marginata** Melville, Kew Bull. 7: 175 (1952) 2:449
Epacris micranthera F.Muell. = **Archeria serpyllifolia**
Epacris microphylla Hook.f. nom. illeg., sensu Hooker (1847) = **Epacris petrophila**
Epacris microphylla R.Br. sensu Bentham (1868), Rodway (1903) p.p. = **Epacris gunnii** (Tasmanian material)
Epacris microphylla var. *gunnii* (Hook.f.) Benth. = **Epacris gunnii**
- e **Epacris moscaliana** Crowden, Muellera 25: 127 (2007)
- e **Epacris mucronulata** R.Br., Prodr. Fl. Nov. Holland. 552 (1810) 2:448
Epacris mucronulata sensu Hooker (1860) p.p. = **Epacris acuminata** (misapplied in Tasmania)
- e **Epacris myrtifolia** Labill., Nov. Holl. Pl. 1: 41 t.55 (1805) 2:447
Epacris myrtifolia var. *corymbiflora* (Hook.f.) Rodway = **Epacris corymbiflora**
- e **Epacris navicularis** Jarman, Pap. & Proc. Roy. Soc. Tasmania 112: 3 (1978)
- Epacris obtusifolia** Sm., Exot. Bot. 1: 77 t.40 (1804) 2:447
- Epacris paludosa** R.Br., Prodr. Fl. Nov. Holland. 551 (1810) 2:449
- Epacris petrophila** Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 261 (1857) 2:446
Epacris ruscifolia R.Br. = **Epacris impressa**
- e **Epacris serpyllifolia** R.Br., Prodr. Fl. Nov. Holland. 551 (1810) 2:450
Epacris serpyllifolia var. *squarrosa* (Hook.f.) Benth. = **Epacris tasmanica**

| | | |
|-----|--|-------|
| | Epacris squarrosa Hook.f. = Epacris tasmanica | 2:450 |
| e | Epacris stuartii Stapf, Bull. Misc. Inform. Kew 1910: 217 (1910) | 2:449 |
| e | Epacris tasmanica W.M.Curtis, Taxon 18: 244 (1969) | 2:450 |
| | Epacris variabilis Lodd. ex Paxton nom. illeg., non Courtois (1833) = Epacris impressa | |
| e | Epacris virgata Hook.f., London J. Bot. 6: 271 (1847) | 2:448 |
| | Erica andromedaeiflora Andrews = Erica holosericea | 2:419 |
| i | Erica arborea L., Sp. Pl. 1: 353 (1753) | 2:420 |
| i | Erica baccans L., Mant. Pl. 2: 233 (1771) | 2:419 |
| i | Erica caffra L., Sp. Pl. 1: 353 (1753) | 2:421 |
| i t | Erica holosericea Salisb., Trans. Linn. Soc. London 6: 352 (1802) | 2:419 |
| i | Erica lusitanica Rudolphi, J. Bot. (Schrader) 2: 286 (1800) | 2:420 |
| i t | Erica scoparia L., Sp. Pl. 1: 353 (1753) | |
| | Erica aff. willmorei Knowles & Westc. sensu Curtis (1963) recorded in error | 2:420 |
| | Gaultheria antipoda G.Forst. sensu Hooker (1860), Bentham (1868), Rodway (1903) = Gaultheria depressa (misapplied in Tasmania) | |
| | Gaultheria antipoda G.Forst. var. depressa (Hook.f.) Hook.f. = Gaultheria depressa | |
| t | Gaultheria depressa Hook.f., London J. Bot. 6: 267 (1847) | 2:417 |
| | Gaultheria hispida R.Br., Prodr. Fl. Nov. Holland. 559 (1810) | 2:417 |
| e | Gaultheria lanceolata Hook.f., London J. Bot. 6: 267 (1847) | 2:416 |
| e | Gaultheria tasmanica (Hook.f.) D.J.Middleton, Edinburgh J. Bot. 47: 299 (1990) | 2:416 |
| e | Leptecophylla abietina (Labill.) C.M.Weiller, Muelleria 12: 211 (1999) | 2:427 |
| e | Leptecophylla divaricata (Hook.f.) C.M.Weiller, Muelleria 12: 207 (1999) | 2:428 |
| | Leptecophylla juniperina (J.R.Forst. & G.Forst.) C.M.Weiller subsp. juniperina sensu de Salas & Baker (2016) = Leptecophylla oxycedrus and L. pogonocalyx subsp. decipiens (misapplied in Tasmania) | 2:427 |
| | Leptecophylla juniperina (J.R.Forst. & G.Forst.) C.M.Weiller subsp. oxycedrus (Labill.) C.M.Weiller = Leptecophylla oxycedrus | |
| | Leptecophylla juniperina (J.R.Forst. & G.Forst.) C.M.Weiller subsp. parvifolia (R.Br.) C.M.Weiller = Leptecophylla parvifolia | |
| | Leptecophylla oxycedrus (Labill.) Jarman, Swainsona 31: 1 | |
| e | Leptecophylla parvifolia (R.Br.) Jarman, Swainsona 31: 7 | 2:428 |
| e | Leptecophylla pendulosa (Jarman) C.M.Weiller, Muelleria 12: 209 (1999) | |
| e | Leptecophylla pogonocalyx C.M.Weiller subsp. decipiens Jarman, Swainsona 31: 13 (2017) | |
| e | Leptecophylla pogonocalyx C.M.Weiller subsp. pogonocalyx , Muelleria 12: 206 (1999) | |
| | Leucopogon affinis R.Br., Prodr. Fl. Nov. Holland 541 (1810) | 2:435 |
| | Leucopogon australis R.Br., Prodr. Fl. Nov. Holland. 541 (1810) | 2:436 |
| | Leucopogon ciliatus A.Cunn. ex DC. = Leucopogon collinus | |
| | Leucopogon ciliatus var. [alpha] Hook.f. = Leucopogon collinus | |
| | Leucopogon ciliatus var. [beta] Hook.f. = Leucopogon oreophilus and Leucopogon pilifer | |
| | Leucopogon collinus (Labill.) R.Br., Prodr. Fl. Nov. Holland. 543 (1810) | 2:436 |
| | Leucopogon collinus var. billardierei DC. = Leucopogon collinus | |
| | Leucopogon collinus var. brownii DC. = Leucopogon collinus | |
| | Leucopogon ericoides (Sm.) R.Br., Prodr. Fl. Nov. Holland. 543 (1810) | 2:438 |

| | | |
|---|--|-------|
| | Leucopogon ericoides var. [alpha] Hook.f. = Leucopogon ericoides | |
| | Leucopogon ericoides var. [beta] Hook.f. = Leucopogon ericoides | |
| | Leucopogon esquamatus R.Br., Prodr. Fl. Nov. Holland. 546 (1810) | 2:439 |
| | Leucopogon fraseri A.Cunn., Ann. Nat. Hist. 2: 47 (1839) | 2:439 |
| | Leucopogon hookeri Sond. = Acrothamnus hookeri | 2:438 |
| | Leucopogon lanceolatus (Sm.) R.Br. = Leucopogon affinis | 2:435 |
| | Leucopogon milliganii (F.Muell.) Rodway = Androstoma verticillata | 2:439 |
| | Leucopogon montanus (R.Br.) J.H.Willis = Acrothamnus montanus | 2:433 |
| | Leucopogon obtusatus nom. illeg., sensu Hooker (1847) = Acrothamnus hookeri | 2:438 |
| e | Leucopogon oreophilus J.M.Powell, Aspects of Tasmanian Botany: 108 (1991) | |
| | Leucopogon parviflorus (Andrews) Lindl., Edwards's Bot. Reg. 18: t.1560 (1832) | 2:435 |
| | Leucopogon pilifer N.A.Wakef., Vict. Naturalist 73: 58 (1956) | |
| | Leucopogon richei (Labill.) R.Br. = Leucopogon parviflorus | |
| | Leucopogon richei var. parviflorus (Andrews) Hook.f. = Leucopogon parviflorus | |
| | Leucopogon richei var. [gamma] Hook.f. = Leucopogon affinis | |
| | Leucopogon rufus Lindl. sensu Bentham (1868) recorded in error | |
| | Leucopogon stuartii F.Muell. ex Sond. = Leucopogon fraseri | 2:439 |
| | Leucopogon trichocarpus (Labill.) R.Br. = Leucopogon ericoides | |
| | Leucopogon virgatus (Labill.) R.Br. var. brevifolius Benth., Fl. Austral. 4: 202 (1868) | 2:438 |
| | Leucopogon virgatus (Labill.) R.Br. var. virgatus , Prodr. Fl. Nov. Holland. 543 (1810) | 2:438 |
| | Lissanthe ciliata R.Br. = Brachyloma ciliatum | |
| | Lissanthe daphnoides (Sm.) R.Br. sensu Brown (1810), Hooker (1860) mistakenly attributed to Tasmania | |
| | Lissanthe montana R.Br. = Acrothamnus montanus | 2:433 |
| | Lissanthe strigosa (Sm.) R.Br. subsp. subulata (R.Br.) J.M.Powell, Telopea 5: 741 (1994) | 2:433 |
| | Monotoca elliptica (Sm.) R.Br., Prodr. Fl. Nov. Holland. 546 (1810) | 2:441 |
| e | Monotoca empetrifolia R.Br., Prodr. Fl. Nov. Holland. 547 (1810) | 2:443 |
| | Monotoca glauca (Labill.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 636 (1917) | 2:442 |
| | Monotoca lineata R.Br. = Monotoca glauca | |
| | Monotoca lineata var. linifolia Rodway = Monotoca linifolia | |
| e | Monotoca linifolia (Rodway) W.M.Curtis subsp. algida Jarman, Aspects of Tasmanian Botany: 102 (1991) | 2:442 |
| e | Monotoca linifolia (Rodway) W.M.Curtis subsp. linifolia , The Student's Flora of Tasmania 2: 464 (1963) | 2:442 |
| | Monotoca scoparia (Sm.) R.Br., Prodr. Fl. Nov. Holland. 547 (1810) | 2:442 |
| | Monotoca scoparia var. submutica Benth. = Monotoca submutica | |
| e | Monotoca submutica (Benth.) Jarman var. autumnalis Jarman, Pap. & Proc. Roy. Soc. Tasmania 112: 1 (1978) | |
| e | Monotoca submutica (Benth.) Jarman var. submutica , Pap. & Proc. Roy. Soc. Tasmania 112: 1 (1978) | |
| t | Montitega dealbata (R.Br.) C.M.Weiller, Austral. Syst. Bot. 23: 328 (2010) | 2:427 |
| e | Pentachondra ericifolia Hook.f., London J. Bot. 6: 271 (1847) | 2:430 |
| e | Pentachondra involucrata R.Br., Prodr. Fl. Nov. Holland. 549 (1810) | 2:429 |
| | Pentachondra mucronata Hook.f. = Leucopogon fraseri | |

| | | |
|---|---|-------|
| | Pentachondra pumila (J.R.Forst. & G.Forst.) R.Br., Prodr. Fl. Nov. Holland. 549 (1810) | 2:429 |
| | <i>Pentachondra verticillata</i> Hook.f. = Androstoma verticillata | |
| | <i>Pernettya lanceolata</i> (Hook.f.) B.L.Burtt & A.W.Hill = Gaultheria lanceolata | 2:416 |
| | <i>Pernettya tasmanica</i> Hook.f. = Gaultheria tasmanica | 2:416 |
| | <i>Pilitis acerosa</i> Lindl. = Richea acerosa | |
| | <i>Pilitis milliganii</i> Hook.f. = Richea milliganii | |
| e | Planocarpa nitida (Jarman) C.M.Weiller, Austral. Syst. Bot. 9: 517 (1996) | |
| e | Planocarpa petiolaris (DC.) C.M.Weiller, Austral. Syst. Bot. 9: 514 (1996) | 2:426 |
| e | Planocarpa sulcata (Mihaich) C.M.Weiller, Austral. Syst. Bot. 9: 516 (1996) | |
| e | Prionotes cerinthoides (Labill.) R.Br., Prodr. Fl. Nov. Holland. 553 (1810) | 2:453 |
| i | Rhododendron ponticum L., Sp. Pl., ed. 2: 562 (1762) | |
| e | Richea acerosa (Lindl.) F.Muell., Fragm. (Mueller) 6: 69 (1867) | 2:458 |
| e | Richea alpina Menadue, Austral. Syst. Bot. 13: 798 (2000) | |
| | <i>Richea angustifolia</i> B.L.Burtt = Richea scoparia | 2:460 |
| e | Richea × curtisiae A.M.Gray, Muelleria 2: 143 (1971) = R. pandanifolia × R. scoparia | |
| e | Richea dracophylla R.Br., Prodr. Fl. Nov. Holland. 555 (1810) | 2:460 |
| e | Richea gunnii Hook.f., London J. Bot. 6: 273 (1847) | 2:459 |
| e | Richea milliganii (Hook.f.) F.Muell., Fragm. (Mueller) 6: 69 (1867) | 2:458 |
| e | Richea pandanifolia Hook.f. subsp. pandanifolia , Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 50 (1844) | 2:460 |
| e | Richea pandanifolia Hook.f. subsp. ramulosa Menadue, Austral. Syst. Bot. 13: 795 (2000) | 2:460 |
| e | Richea procera (F.Muell.) F.Muell., Fragm. (Mueller) 6: 68 (1867) | 2:458 |
| e | Richea scoparia Hook.f., London J. Bot. 6: 273 (1847) | 2:459 |
| e | Richea sprengelioides (R.Br.) F.Muell., Fragm. (Mueller) 6: 68 (1867) | 2:457 |
| e | Sprengelia distichophylla (Rodway) W.M.Curtis, The Student's Flora of Tasmania 2: 464 (1963) | 2:456 |
| | Sprengelia incarnata Sm., Kongl. Vetensk. Acad. Nya Handl. 15: 260 (1794) | 2:454 |
| | <i>Sprengelia incarnata</i> f. <i>montana</i> (R.Br.) Siebert & Voss = Sprengelia montana | |
| | <i>Sprengelia incarnata</i> f. <i>propinqua</i> (A.Cunn. ex DC.) Siebert & Voss = Sprengelia propinqua | |
| | <i>Sprengelia incarnata</i> var. <i>distichophylla</i> Rodway = Sprengelia distichophylla | |
| | <i>Sprengelia incarnata</i> var. <i>montana</i> (R.Br.) Domin = Sprengelia montana | |
| | <i>Sprengelia macrantha</i> Hook.f. = Sprengelia propinqua | |
| e | Sprengelia minima Crowden, Telopea 15: 62 (2013) | |
| e | Sprengelia montana R.Br., Prodr. Fl. Nov. Holland. 555 (1810) | 2:455 |
| | <i>Sprengelia</i> sp. Mt Field (A.M.Gray 1513) Tas Herbarium = Sprengelia minima | |
| e | Sprengelia sp. Mt Read , Telopea 15: 63 (2013) (R.K. Crowden 0802 006) Tas Herbarium | |
| e | Sprengelia propinqua A.Cunn. ex DC., Prodr. [A. P. de Candolle] 7: 768 (1839) | |
| | <i>Sprengelia propinqua</i> var. <i>demissa</i> F.Muell. = Sprengelia minima | |
| n | Stenantha pinifolia R.Br., Prodr. Fl. Nov. Holland. 538 (1810) | 2:424 |
| | <i>Styphelia abietina</i> Labill. = Leptecophylla abietina | |
| | Styphelia adscendens R.Br., Prodr. Fl. Nov. Holland. 537 (1810) | 2:423 |
| | <i>Styphelia billardiarei</i> = Cyathodes glauca | |
| | <i>Styphelia ciliata</i> (R.Br.) F.Muell. = Brachyloma ciliatum | |
| | <i>Styphelia collina</i> Labill. = Leucopogon collinus | |
| | <i>Styphelia empetrifolia</i> (R.Br.) F.Muell. = Monotoca empetrifolia | |

| | | |
|-------|--|----------------|
| | Styphelia glauca Labill. = Monotoca glauca | |
| | Styphelia hookeri = Acrothamnus hookeri | |
| | Styphelia lanceolata Sm. = Leucopogon affinis | |
| | Styphelia montana (R.Br.) F.Muell. = Acrothamnus montanus | |
| | Styphelia mucronata (Hook.f.) J.H.Willis = Leucopogon fraseri | |
| | Styphelia oxycedrus Labill. = Leptecophylla oxycedrus | |
| | Styphelia serrulata Labill. = Acrotriche serrulata | |
| | Styphelia straminea (R.Br.) Spreng. = Cyathodes straminea | |
| | Styphelia strigosa Sm. = Lissanthe strigosa | |
| | Styphelia trichocarpa Labill. = Leucopogon ericoides | |
| | Styphelia virgata Labill. = Leucopogon virgatus | |
| e | Trochocarpa cunninghamii (DC.) W.M.Curtis, The Student's Flora of Tasmania 2: 463 (1963) | 2:431 |
| e | Trochocarpa disticha (Labill.) Spreng., Syst. Veg. (ed. 16) [Sprengel] 1: 660 (1824) | 2:430 |
| | Trochocarpa disticha var. cunninghamii (DC.) Benth. = Trochocarpa cunninghamii | |
| | Trochocarpa disticha var. microphylla F.Muell. = Trochocarpa disticha | |
| | Trochocarpa glauca (Labill.) Spreng. = Cyathodes glauca | |
| e | Trochocarpa gunnii (Hook.f.) Benth., Fl. Austral. 4: 167 (1868) | 2:431 |
| e | Trochocarpa thymifolia (R.Br.) Spreng., Syst. Veg. (ed. 16) [Sprengel] 1: 660 (1824) | 2:431 |
| | ESCALLONIACEAE | FTO 131 |
| e | Anopterus glandulosus Labill., Nov. Holl. Pl. 1: 86 t. 112 (1805) | 1:181 |
| | EUPHORBIACEAE | |
| | Adriana quadripartita (Labill.) Müll.Arg. sensu Bentham (1873) recorded in error | |
| | Amperea spartioides Brongn. = Amperea xiphoclada var. xiphoclada | |
| | Amperea xiphoclada (Sieber ex Spreng.) Druce var. xiphoclada , Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 604 (1917) | 3:637 |
| | Bertya rosmarinifolia (A.Cunn.) Planch. sensu Curtis (1967) = Bertya tasmanica subsp. tasmanica (misapplied in Tasmania) | |
| e | Bertya tasmanica (Sond. & F.Muell.) Müll.Arg. subsp. tasmanica , Linnaea 34: 63 (1865) | 3:637 |
| | Beyeria backhousei Hook.f. = Beyeria lechenaultii | |
| | Beyeria lechenaultii (DC.) Baill., Adansonia 6: 307 (1866) | 3:635 |
| | Beyeria lechenaultii (DC.) Baill. var. latifolia Grüning = Beyeria lechenaultii | |
| | Beyeria oblongifolia (Klotzsch) Hook.f. = Beyeria viscosa | |
| | Beyeria opaca F.Muell. sensu Bentham (1873) = Beyeria lechenaultii (misapplied in Tasmania) | |
| | Beyeria viscosa (Labill.) Miq., Ann. Sci. Nat., Bot., sér. 3, 1: 350 t. 15 (1844) | 3:635 |
| | Chamaesyce drummondii (Boiss.) D.C.Hassall = Euphorbia drummondii | 3:632 |
| i | Euphorbia cyparissias L., Sp. Pl. 1: 461 (1753) | 3:633 |
| | Euphorbia drummondii Boiss. sensu Curtis (1967) recorded in error | 3:632 |
| i | Euphorbia exigua L., Sp. Pl. 1: 456 (1753) | |
| i | Euphorbia helioscopia L., Sp. Pl. 1: 459 (1753) | 3:632 |
| i | Euphorbia lathyris L., Sp. Pl. 1: 457 (1753) | 3:632 |
| i | Euphorbia paralias L., Sp. Pl. 1: 458 (1753) | 3:633 |
| i | Euphorbia peplus L., Sp. Pl. 1: 456 (1753) | 3:633 |
| i # t | Euphorbia stricta L., Syst. Nat., ed. 10, 2: 1049 (1759) | |

| | | |
|-------------------------------|--|-------|
| | Ricinocarpos major F.Muell. sensu Benth (1873) a name of uncertain application | |
| | Ricinocarpos pinifolius Desf., Mém. Mus. Hist. Nat. 3: 459 t.22 (1817) | 3:636 |
| FABACEAE (LEGUMINOSAE) | | |
| | Acacia armata R.Br. = Acacia paradoxa | 1:127 |
| e | Acacia axillaris Benth., London J. Bot. 1: 341 (1842) | 1:127 |
| i | Acacia baileyana F.Muell., Trans. & Proc. Roy. Soc. Victoria 24: 168 (1888) | |
| | Acacia botrycephala (Vent.) Desf. = Acacia terminalis | 1:129 |
| | Acacia crassiuscula H.L.Wendl. sensu Hooker (1860), Benth (1864) = Acacia uncifolia (misapplied in Tasmania) | |
| | Acacia dealbata Link subsp. dealbata , Enum. Hort. Berol. Alt. 2: 445 (1822) | 1:130 |
| i | Acacia decurrens Willd., Sp. Pl., ed. 4 [Willdenow], 4: 1072 (1806) | 1:130 |
| | Acacia decurrens Willd. f. mollis (Lindl.) Benth. = Acacia dealbata subsp. dealbata | |
| | Acacia dependens A.Cunn. ex Benth. nom. illeg. = Acacia mucronata subsp. dependens | |
| e | Acacia derwentiana A.M.Gray, Muellera 21: 107 (2005) | |
| | Acacia diffusa Ker Gawl. = Acacia genistifolia | |
| | Acacia discolor (Andrews) Willd. = Acacia terminalis | |
| | Acacia dissitiflora Benth. = Acacia mucronata subsp. longifolia | |
| i | Acacia floribunda (Vent.) Willd., Sp. Pl., ed. 4 [Willdenow], 4: 1051 (1806) | |
| | Acacia genistifolia Link, Enum. Hort. Berol. Alt. 2: 442 (1822) | 1:126 |
| | Acacia gunnii Benth., London J. Bot. 1: 332 (1842) | 1:127 |
| i | Acacia howittii F.Muell., Vict. Naturalist 10: 16 (1893) | |
| | Acacia implexa Benth. sensu Buchanan (1999) = Acacia uncifolia (misapplied in Tasmania) | |
| | Acacia juniperina (Vent.) Willd. sensu Benth (1864) = Acacia ulicifolia (misapplied in Tasmania) | |
| | Acacia leprosa Sieber ex DC. var. graveolens Maslin & D.J.Murphy, Muellera 27: 201 (2009) | 1:127 |
| | Acacia linearis (J.C.Wendl.) Sims sensu Hooker (1860), Benth (1864) = Acacia mucronata subsp. mucronata (misapplied in Tasmania) | |
| | Acacia longifolia (Andrews) Willd. f. dissitiflora (Benth.) Benth. = Acacia mucronata subsp. longifolia | |
| | Acacia longifolia (Andrews) Willd. f. mucronata (Willd. ex H.L.Wendl.) Benth. = Acacia mucronata | |
| | Acacia longifolia (Andrews) Willd. subsp. longifolia , Sp. Pl., ed. 4 [Willdenow], 4: 1052 (1806) | 1:129 |
| | Acacia longifolia (Andrews) Willd. subsp. sophorae (Labill.) Court, Fl. Australia 11B: 491 (2001) | 1:129 |
| | Acacia mearnsii De Wild., Pl. Bequaert. 3: 61 (1925) | 1:130 |
| | Acacia melanoxydon R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 5: 462 (1813) | 1:128 |
| | Acacia mollissima Willd. sensu Hooker (1860) = Acacia mearnsii (misapplied in Tasmania) | |
| e | Acacia mucronata Willd. ex H.L.Wendl. subsp. dependens (Hook.f.) Court, Fl. Australia 11B: 491 (2001) | 1:129 |
| | Acacia mucronata Willd. ex H.L.Wendl. subsp. longifolia (Benth.) Court, Fl. Australia 11B: 491 (2001) | 1:129 |
| e | Acacia mucronata Willd. ex H.L.Wendl. subsp. mucronata , Comm. Acac. Aphyll. 46 t.12 (1820) | 1:129 |
| | Acacia mucronata Willd. ex H.L.Wendl. var. dissitiflora (Benth.) Hook.f. = Acacia mucronata subsp. longifolia | |
| | Acacia mucronata Willd. ex H.L.Wendl. var. linearis (Sims) Rodway = Acacia mucronata subsp. mucronata | |
| | Acacia myrtifolia (Sm.) Willd., Sp. Pl., ed. 4 [Willdenow], 4: 1054 (1806) | 1:128 |
| | Acacia ovoidea Benth. = Acacia verticillata subsp. ovoidea | |

| | | |
|---|---|-------|
| | Acacia oxycedrus Sieber ex DC. sensu Benth (1864) recorded in error | |
| i | Acacia paradoxa DC., Cat. Pl. Horti Monsp. 74 (1813) | 1:127 |
| e | Acacia pataczekii D.I.Morris, Records of the Queen Victoria Museum 50: 1 (1974) | 1:128 |
| | Acacia penninervis Sieber ex DC. sensu Benth (1864) recorded in error | |
| i | Acacia pravissima F.Muell. ex Benth., Linnaea 26: 608 (1854) | |
| i | Acacia provincialis A.Camus, Bull. Soc. Dendrol. France 64: 68 (1927) | |
| i | Acacia pycnantha Benth., London J. Bot. 1: 351 (1842) | |
| i | Acacia retinodes Schltld., Linnaea 20: 664 (1847) | |
| | Acacia retinodes Schltld. var. uncifolia J.M.Black = Acacia uncifolia | 1:128 |
| e | Acacia riceana Hensl., Botanist 3: t.135 (1839) | 1:127 |
| | Acacia siculiformis A.Cunn. ex Benth., London J. Bot. 1: 337 (1842) | 1:126 |
| | Acacia sophorae (Labill.) R.Br. = Acacia longifolia subsp. sophorae | 1:129 |
| | Acacia stricta (Andrews) Willd., Sp. Pl., ed. 4 [Willdenow], 4: 1052 (1806) | 1:127 |
| | Acacia stuartiana F.Muell. ex Benth. = Acacia siculiformis | |
| | Acacia suaveolens (Sm.) Willd., Sp. Pl., ed. 4 [Willdenow], 4: 1050 (1806) | 1:128 |
| e | Acacia sp. Sugarloaf (A.Moscal 8301) Tas Herbarium | |
| | Acacia terminalis (Salisb.) J.F.Macbr., Contr. Gray Herb. 59: 7 (1919) | 1:129 |
| | Acacia ulicifolia (Salisb.) Court, Vict. Naturalist 73: 173 (1957) | 1:126 |
| | Acacia uncifolia (J.M.Black) O'Leary, J. Adelaide Bot. Gard. 21: 100 (2007) | 1:128 |
| | Acacia verniciflua A.Cunn. sensu Curtis (1975) = Acacia leprosa var. graveolens (misapplied in Tasmania) | 1:127 |
| | Acacia verticillata (L'Hér.) Willd. subsp. ovoidea (Benth.) Court, Fl. Australia 11B: 492 (2001) | 1:126 |
| | Acacia verticillata (L'Hér.) Willd. subsp. ruscifolia (A.Cunn. ex G.Don) Court, Fl. Australia 11B: 492 (2001) | 1:126 |
| | Acacia verticillata (L'Hér.) Willd. subsp. verticillata , Sp. Pl., ed. 4 [Willdenow], 4: 1049 (1806) | 1:126 |
| | Acacia verticillata (L'Hér.) Willd. var. latifolia Benth. nom. illeg. = Acacia verticillata subsp. ruscifolia | |
| | Acacia vomeriformis A.Cunn. ex Benth. = Acacia gunnii | 1:127 |
| | Albizia lophantha (Willd.) Benth. = Paraserianthes lophantha | 1:130 |
| | Almaleea subumbellata (Hook.) Crisp & P.H.Weston, Telopea 4: 310 (1991) | 1:141 |
| | Anthyllis vulneraria L. recorded in error, only known from cultivated plants | |
| | Aotus ericoides (Vent.) G.Don, Gen. Hist. 2: 120 (1832) | 1:137 |
| | Aotus ferruginea Labill. = Aotus ericoides | |
| | Aotus villosa Sm. = Aotus ericoides | 1:137 |
| | Bossiaea cinerea R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 4: 268 (1812) | 1:147 |
| | Bossiaea cinerea R.Br. var. rigida Rodway = Bossiaea tasmanica | |
| | Bossiaea cordifolia Sweet, Fl. Australas. (Sweet) 20: 20 (1827) | |
| | Bossiaea cordigera Benth. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 95 t.16 (1856) | 1:147 |
| | Bossiaea ensata Sieber sensu Hooker (1860) non Sieber = Bossiaea riparia (misapplied in Tasmania) | |
| | Bossiaea heterophylla Vent., Descr. Pl. Nouv. 1: 7, pl.7 (1800) | |
| | Bossiaea microphylla Sm. sensu Curtis (1956) = Bossiaea tasmanica (misapplied in Tasmania) | |
| | Bossiaea obcordata (Vent.) Druce = Bossiaea tasmanica | 1:148 |
| | Bossiaea prostrata R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 4: 268 (1812) | 1:147 |
| | Bossiaea riparia A.Cunn. ex Benth., Fl. Austral. 2: 166 (1864) | 1:148 |

| | | |
|---|---|-------|
| e | Bossiaea tasmanica I.Thomps., Muelleria 30: 144 (2012) | 1:148 |
| i | Callistachys lanceolata Vent., Jard. Malmaison 2:115, t.115 (1805) | |
| i | Chamaecytisus palmensis (Christ) F.A.Bisby & K.W.Nicholls, Bot. J. Linn. Soc. 74: 114 (1977) | 1:151 |
| | Chamaecytisus prolifer Link sensu Curtis & Morris (1975) = Chamaecytisus palmensis (misapplied in Tasmania) | 1:151 |
| | Coronilla varia L. = Securigera varia | 1:165 |
| | Cullen microcephalum (Rchb. ex Kunze) J.W.Grimes, Muelleria 9: 195 (1996) | 1:163 |
| | Cytisus monspessulanus L. = Genista monspessulana | |
| i | Cytisus multiflorus (L'Hér.) Sweet, Hort. Brit. [Sweet] 112 (1826) | |
| | Cytisus palmensis (H.Christ) Hutch. = Chamaecytisus palmensis | 1:151 |
| | Cytisus prolifer L.f. sensu Curtis (1956) = Chamaecytisus palmensis (misapplied in Tasmania) | |
| i | Cytisus scoparius (L.) Link, Enum. Hort. Berol. Alt. 2: 241 (1822) | 1:151 |
| | Daviesia latifolia R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 3: 20 (1811) | 1:136 |
| | Daviesia sejugata G.Chandler & Crisp, Austral. Syst. Bot. 10: 33 (1997) | |
| | Daviesia ulicifolia Andrews subsp. ruscifolia (A.Cunn. ex Benth.) G.Chandler & Crisp, Austral. Syst. Bot. 10: 44 (1997) | 1:136 |
| | Daviesia ulicifolia Andrews subsp. ulicifolia , Bot. Repos. 5: t.304 (1803) | 1:136 |
| | Daviesia ulicina Sm. nom. illeg., nom. superfl. = Daviesia ulicifolia | |
| | Daviesia ulicina Sm. f. communis Benth. = Daviesia ulicifolia subsp. ulicifolia | |
| | Daviesia ulicina Sm. f. ruscifolia (A.Cunn. ex Benth.) Benth. = Daviesia ulicifolia subsp. ruscifolia | |
| | Daviesia ulicina Sm. f. subumbellata Benth. = Daviesia ulicifolia subsp. ulicifolia | |
| | Daviesia umbellata Sm. sensu Labill. (1805) = Daviesia ulicifolia (misapplied in Tasmania) | |
| | Desmodium gunnii Benth. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 101 (1856) | 1:166 |
| | Desmodium varians (Labill.) G.Don, Gen. Hist. 2: 298 (1832) | 1:166 |
| | Desmodium varians (Labill.) G.Don var. gunnii (Benth. ex Hook.f.) Benth. = Desmodium gunnii | |
| | Dillwynia cinerascens R.Br. ex Sims, Bot. Mag. 48: t.2247 (1821) | 1:145 |
| | Dillwynia ericifolia Sm. f. glaberrima (Sm.) Benth. = Dillwynia glaberrima | |
| | Dillwynia floribunda Sm. sensu Bentham (1864), Rodway (1903), Curtis (1956) = Dillwynia sericea (misapplied in Tasmania) | |
| | Dillwynia glaberrima Sm., Ann. Bot. [König & Sims] 1: 510 (1805) | 1:144 |
| | Dillwynia sericea A.Cunn., Geogr. Mem. New South Wales: 347 (1825) | 1:144 |
| i | Dipogon lignosus (L.) Verdc., Taxon 17: 537 (1968) | |
| | Dolichos lignosus L. = Dipogon lignosus | |
| | Eutaxia microphylla (R.Br.) C.H.Wright & Dewar, Johnson's Gard. Dict. 1052 (1894) | 1:143 |
| i | Genista linifolia L., Sp. Pl., ed. 2, 2: 997 (1763) | |
| | Genista maderensis (Webb & Berthel.) Lowe sensu Curtis & Morris (1975) = Genista stenopetala (misapplied in Tasmania) | 1:152 |
| i | Genista monspessulana (L.) L.A.S.Johnson, Contr. New South Wales Natl. Herb. 3: 98 (1962) | 1:151 |
| i | Genista × spachiana Webb, Bot. Mag. 71: t.4195 (1845) | |
| i | Genista stenopetala Webb & Berthel., Hist. Nat. Iles Canaries (Phytogr.) 3(2): 39 (1836) | 1:152 |
| | Glycine clandestina J.C.Wendl., Bot. Beob. [Wendland]: 54 (1798) | 1:168 |
| | Glycine clandestina J.C.Wendl. var. latrobeana Rodway = Glycine latrobeana | |
| | Glycine latrobeana (Meisn.) Benth., Fl. Austral. 2: 244 (1864) | 1:168 |

| | | |
|-----|--|-------|
| | Glycine microphylla (Benth.) Tindale, Brunonia 9: 181 (1987) | |
| | Glycine tabacina (Labill.) Benth., Fl. Austral. 2: 244 (1864) | |
| | Gompholobium ecostatum Kuchel, Suppl. Black's Fl. S. Austral. 182 (1965) | 1:135 |
| | Gompholobium ellipticum Labill. = Oxylobium ellipticum | |
| | Gompholobium huegelii Benth., Enum. Pl. [Endlicher]: 29 (1837) | 1:135 |
| | Gompholobium latifolium Labill. nom. illeg., non Sm. = Gompholobium huegelii | |
| | Goodia lotifolia Salisb., Parad. Lond. 1: t.41 (1806) | 1:149 |
| | Goodia lotifolia Salisb. var. pubescens (Sims) H.B. Will. = Goodia pubescens | 1:149 |
| | Goodia pubescens Sims, Bot. Mag. 32: 1310 (1810) | 1:149 |
| | Hardenbergia monophylla (Vent.) Benth. = Hardenbergia violacea | |
| | Hardenbergia violacea (Schneev.) Stearn, J. Bot. 78: 70 (1940) | 1:169 |
| | Hedysarum coronarium L. previously listed as naturalised but insufficient evidence exists to support this | 1:165 |
| | Hovea corrickiae J.H. Ross, Muelleria 7: 203 (1990) | |
| | Hovea heterophylla A. Cunn. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 93 (1856) | 1:148 |
| | Hovea lanceolata Sims sensu Buchanan (1999) = Hovea tasmanica (misapplied in Tasmania) | |
| | Hovea linearis (Sm.) R.Br. sensu Buchanan (1999) = Hovea heterophylla (misapplied in Tasmania) | |
| | Hovea longifolia R.Br. sensu Curtis (1956), Curtis & Morris (1975) misapplied to H. montana & H. tasmanica | 1:149 |
| | Hovea magnibractea I. Thomps., Austral. Syst. Bot. 14: 74 (2001) | |
| | Hovea montana (Hook.f.) J.H. Ross, Muelleria 6: 427 (1988) | 1:149 |
| | Hovea purpurea Sweet var. montana Hook.f. = Hovea montana | |
| e | Hovea tasmanica I. Thomps. & J.H. Ross, Austral. Syst. Bot. 14: 68 (2001) | 1:149 |
| | Indigofera australis Willd. subsp. australis , Sp. Pl., ed. 4 [Willdenow], 3: 1235 (1802) | 1:164 |
| | Indigofera australis Willd. var. angulata Benth. = Indigofera australis subsp. australis | |
| i | Kennedia nigricans Lindl., Edwards's Bot. Reg. 20: 1715, pl. 1715 (1835) | |
| | Kennedia prostrata R.Br., Hortus Kew. (W.T. Aiton), ed. 2, 4: 299 (1812) | 1:169 |
| i | Kennedia rubicunda (Schneev.) Vent., Jard. Malmaison 2: 104 t.104 (1804) | |
| i # | Laburnum anagyroides Medik., Vorles. Churpfälz. Phys.-Öcon. Ges. 2: 363 (1787) | |
| i # | Lathyrus nissolia L., Sp. Pl. 2: 729 (1753) | 1:168 |
| i | Lathyrus tingitanus L., Sp. Pl. 2: 732 (1753) | |
| | Leptocyamus clandestinus (J.C. Wendl.) Benth. var. clandestinus = Glycine clandestina | |
| | Leptocyamus clandestinus (J.C. Wendl.) Benth. var. microphylla (Benth.) Hook.f. = Glycine microphylla | |
| | Leptocyamus tasmanicus Benth. ex Hook.f. = Glycine latrobeana | |
| i # | Lotus angustissimus L., Sp. Pl. 2: 774 (1753) | |
| | Lotus australis Andrews, Bot. Repos. 10 t.624 (1811) | 1:162 |
| i | Lotus corniculatus L. var. corniculatus , Sp. Pl. 2: 776 (1753) | 1:162 |
| i | Lotus corniculatus L. var. tenuifolius L., Sp. Pl. 2: 776 (1753) | 1:162 |
| | Lotus hispidus Desf. ex DC. sensu Curtis & Morris (1975) = Lotus subbiflorus (misapplied in Tasmania) | 1:163 |
| | Lotus pedunculatus Cav. sensu Curtis & Morris (1975) = Lotus uliginosus (misapplied in Tasmania) | 1:163 |
| | Lotus suaveolens Pers. = Lotus subbiflorus | 1:163 |
| i | Lotus subbiflorus Lag., Varied. Ci. 2(4): 213 (1805) | 1:163 |

| | | |
|-----|---|-------|
| | Lotus tenuis Waldst. & Kit. = Lotus corniculatus var. tenuifolius | |
| i | Lotus uliginosus Schkuhr, Handb. [C.Schkuhr] 2: 412 (1796) | I:163 |
| i # | Lupinus angustifolius L., Sp. Pl. 2: 721 (1753) | I:150 |
| i | Lupinus arboreus Sims, Bot. Mag. 18: t.682 (1803) | I:150 |
| i | Lupinus × regalis Bergmans, Vaste Pl. Rotsheesters: 328 (1924) | |
| i | Medicago arabica (L.) Huds., Fl. Angl. (Hudson): 288 (1762) | I:154 |
| i # | Medicago arborea L., Sp. Pl. 2: 778 (1753) | |
| | Medicago denticulata Willd. = Medicago polymorpha | |
| | Medicago falcata L. sensu Curtis & Morris (1975) = Medicago sativa nothosubsp × varia (misapplied in Tasmania) | I:153 |
| | Medicago hispida Gaertn. var. apiculata (Willd.) Urb. = Medicago polymorpha | |
| | Medicago hispida Gaertn. var. confinis (W.D.J.Koch) Burnat = Medicago polymorpha | |
| | Medicago hispida Gaertn. var. denticulata (Willd.) Urb. = Medicago polymorpha | |
| i | Medicago lupulina L., Sp. Pl. 2: 779 (1753) | I:153 |
| | Medicago maculata Willd. nom. illeg. = Medicago arabica | |
| i | Medicago minima (L.) L. ex Bartal., Cat. Piante Siena 61 (1776) | I:154 |
| i | Medicago polymorpha L., Sp. Pl. 2: 779 (1753) | I:154 |
| i | Medicago sativa L., Sp. Pl. 2: 778 (1753) | I:153 |
| i # | Medicago sativa L. nothosubsp × varia (Martyn) Arcang., Comp. Fl. Ital. 160 (1882) = Medicago falcata × M. sativa | I:153 |
| i | Medicago scutellata (L.) Mill., Gard. Dict., ed. 8, no. 2 (1768) | I:154 |
| | Melilotus alba Desr. nom. illeg. = Melilotus albus | |
| i | Melilotus albus Medik., Vorles. Churpfälz. Phys.-Öcon. Ges. 2: 382 (1787) | I:155 |
| | Melilotus arvensis Wallr. = Melilotus officinalis | |
| i | Melilotus indicus (L.) All., Fl. Pedem. 1: 308 (1785) | I:155 |
| i | Melilotus officinalis (L.) Pall., Reise Russ. Reich. 3: 537 (1776) | I:155 |
| | Melilotus parviflorus Desf. = Melilotus indicus | |
| | Mimosa sophorae Labill. = Acacia longifolia subsp. sophorae | |
| | Mimosa suaveolens Sm. = Acacia suaveolens | |
| | Mirbelia oxylobioides F.Muell., Fragm. (Mueller) 2: 154 (1861) | |
| | Onobrychis viciifolia Scop. previously listed as naturalised but insufficient evidence exists to support this | I:165 |
| | Ononis arvensis L. sensu Rodway (1903) = Ononis spinosa (misapplied in Tasmania) | I:152 |
| | Ononis repens L. sensu Curtis (1956), Curtis & Morris (1975) = Ononis spinosa (misapplied in Tasmania) | I:152 |
| i | Ononis spinosa L., Sp. Pl. 2: 716 (1753) | I:152 |
| i | Ornithopus compressus L., Sp. Pl. 2: 744 (1753) | |
| i | Ornithopus pinnatus (Mill.) Druce, J. Bot. 45: 420 (1907) | |
| i # | Ornithopus sativus Brot., Fl. Lusit. 2: 160 (1805) | |
| | Oxylobium arborescens R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 3: 10 (1811) | I:134 |
| | Oxylobium ellipticum (Vent.) R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 3: 10 (1811) | I:134 |
| | Oxylobium ellipticum (Vent.) R.Br. var. angustifolium Benth. = Oxylobium arborescens | |
| i | Paraserianthes lophantha (Willd.) I.C.Nielsen subsp. lophantha , Bull. Mus. Natl. Hist. Nat., B, Adansonia 5: 326 (1983) - Possibly native in some Furneaux populations, see Harris et al. (2001) | I:130 |

| | | |
|---|--|-------|
| e | Phyllota diffusa (Hook.f.) F.Muell., Fragm. (Mueller) 1: 8 (1858) | 1:137 |
| | Platylobium formosum Sm. sensu Bentham (1864) = Platylobium parviflorum (misapplied in Tasmania) | 1:146 |
| | Platylobium formosum Sm. subsp. parviflorum (Sm.) A.T.Lee = Platylobium parviflorum | |
| | Platylobium murrayanum Hook.f. = Platylobium triangulare | |
| | Platylobium obtusangulum Hook., Bot. Mag. 60: t.3258 (1833) | 1:146 |
| | Platylobium parviflorum Sm., Spec. Bot. New Holland 18 (1795) | |
| | Platylobium triangulare R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 4: 266 (1812) | 1:146 |
| i | Podalyria sericea (Andrews) R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 3: 7 (1811) | |
| | Psoralea adscendens F.Muell. = Cullen microcephalum | 1:163 |
| i | Psoralea arborea Sims, Bot. Mag. 46: t. 2090 (1819) | |
| | Psoralea gunnii Hook.f. = Cullen microcephalum | |
| i | Psoralea pinnata L., Sp. Pl. 2: 762 (1753) | 1:163 |
| | Pultenaea cordata Hook. = Pultenaea juniperina | |
| | Pultenaea daphnoides J.C.Wendl., Bot. Beob. [Wendland]: 49 (1798) | 1:139 |
| | Pultenaea daphnoides J.C.Wendl. var. obcordata (Andrews) Hook.f. = Pultenaea daphnoides | |
| | Pultenaea dentata Labill., Nov. Holl. Pl. 1: 103 t.131 (1805) | 1:141 |
| | Pultenaea diffusa Hook.f. = Phyllota diffusa | |
| | Pultenaea fasciculata Benth., Commentat. Legum. Gen. 18 (1837) | 1:142 |
| | Pultenaea gunnii Benth. subsp. gunnii , Commentat. Legum. Gen. 18 (1837) | 1:140 |
| | Pultenaea gunnii Benth. var. baeckeoides (A.Cunn. ex Benth.) Rodway = Pultenaea gunnii subsp. gunnii | 1:140 |
| | Pultenaea hibbertioides Hook.f. = Pultenaea mollis | 1:141 |
| | Pultenaea humilis Benth. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 91 (1856) | 1:142 |
| | Pultenaea juniperina Labill., Nov. Holl. Pl. 1: 102 t.130 (1805) | 1:142 |
| | Pultenaea juniperina Labill. var. latifolia Benth. = Pultenaea juniperina | |
| | Pultenaea mollis Lindl., Three Exped. Australia [Mitchell] 2: 258 (1838) | 1:141 |
| | Pultenaea paleacea Willd. var. sericea Benth. = Pultenaea sericea | 1:140 |
| | Pultenaea pedunculata Hook., Bot. Mag. 55: t.2859 (1828) | 1:140 |
| | Pultenaea pimelioides Hook.f. = Pultenaea dentata | |
| | Pultenaea prostrata Benth. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 89 (1856) | 1:142 |
| | Pultenaea selaginoides Hook.f. = Stonesiella selaginoides | 1:141 |
| | Pultenaea sericea (Benth.) Corrick, Muellera 8: 392 (1995) | 1:140 |
| | Pultenaea stricta Sims, Bot. Mag. 38: t.1588 (1813) | 1:140 |
| | Pultenaea subumbellata Hook. = Almaleea subumbellata | 1:141 |
| | Pultenaea tenuifolia R.Br. ex Sims, Bot. Mag. 46: t.2086 (1819) | 1:143 |
| | Racosperma axillare (Benth.) Pedley = Acacia axillaris | 1:127 |
| | Racosperma dealbatum (Link) Pedley = Acacia dealbata | 1:130 |
| | Racosperma decurrens (Willd.) Pedley = Acacia decurrens | 1:130 |
| | Racosperma gunnii (Benth.) Pedley = Acacia gunnii | 1:127 |
| | Racosperma implexum (Benth.) Pedley = Acacia uncifolia (misapplied in Tasmania) | |
| | Racosperma mearnsii (De Wild.) Pedley = Acacia mearnsii | 1:130 |
| | Racosperma melanoxyton (R.Br.) Pedley = Acacia melanoxyton | 1:128 |

| | | |
|---|--|-------|
| | Racosperma myrtifolium (Sm.) Mart. = Acacia myrtifolia | 1:128 |
| | Racosperma paradoxum (DC.) Mart. = Acacia paradoxa | 1:127 |
| | Racosperma riceanum (Hensl.) Pedley = Acacia riceana | 1:127 |
| | Racosperma sophorae (Labill.) Mart. = Acacia longifolia subsp. sophorae | 1:129 |
| | Racosperma strictum (Andrews) Mart. = Acacia stricta | 1:127 |
| | Racosperma suaveolens (Sm.) Mart. = Acacia suaveolens | 1:128 |
| | Racosperma ulicifolium (Salisb.) Pedley = Acacia ulicifolia | 1:126 |
| | Racosperma vernicifluum (A.Cunn.) Pedley = Acacia leprosa var. graveolens | 1:127 |
| | Racosperma verticillatum (L'Hér.) Pedley = Acacia verticillata | 1:126 |
| | Sarothamnus scoparius (L.) Wimm. ex W.D.J.Koch = Cytisus scoparius | 1:151 |
| i | Securigera varia (L.) Lassen, Svensk. Bot. Tidskr. 83: 86 (1989) | 1:165 |
| i | Spartium junceum L., Sp. Pl. 2: 708 (1753) | |
| | Sphaerolobium minus Labill., Nov. Holl. Pl. 1: 108, t.138 (1805) | |
| | Sphaerolobium vimineum Sm., Ann. Bot. [König & Sims] 1: 509 (1805) | 1:135 |
| e | Stonesiella selaginoides (Hook.f.) Crisp & P.H.Weston, Taxon 48: 712 (1999) | 1:141 |
| | Swainsona lessertiifolia DC., Ann. Sci. Nat., Bot., sér. 1, 4: 99 (1825) | 1:164 |
| | Trifolium agrarium L. sensu Rodway (1903) = Trifolium campestre (misapplied in Tasmania) | |
| i | Trifolium alexandrinum L., Cent. Pl. 1: 25 (1755) | |
| i | Trifolium angustifolium L., Sp. Pl. 2: 769 (1753) | 1:161 |
| i | Trifolium arvense L., Sp. Pl. 2: 769 (1753) | 1:160 |
| i | Trifolium campestre Schreb., Die Kleearten Deutschlands 1(4): t.17.4 (1800) | 1:157 |
| i | Trifolium cernuum Brot., Phytogr. Lusitan. Select. (1816-1827) 1: 150 (1816) | 1:159 |
| i | Trifolium dubium Sibth., Fl. Oxon. 231 (1794) | 1:157 |
| i | Trifolium fragiferum L., Sp. Pl. 2: 772 (1753) | 1:158 |
| i | Trifolium glomeratum L., Sp. Pl. 2: 770 (1753) | 1:159 |
| i | Trifolium hybridum L., Sp. Pl. 2: 766 (1753) | 1:159 |
| i | Trifolium incarnatum L., Sp. Pl. 2: 769 (1753) | 1:160 |
| i | Trifolium lappaceum L., Sp. Pl. 2: 768 (1753) | |
| | Trifolium medium Huds. sensu Curtis & Morris (1975) = Trifolium pratense (misapplied in Tasmania) | 1:159 |
| i | Trifolium micranthum Viv., Fl. Libyc. Spec. 45, pl. 19, Fig. 1 (1824) | 1:158 |
| i | Trifolium ornithopodioides L., Sp. Pl. 2: 766 (1753) | 1:152 |
| i | Trifolium pratense L., Sp. Pl. 2: 768 (1753) | 1:160 |
| | Trifolium procumbens L. sensu Rodway (1903) = Trifolium dubium (misapplied in Tasmania) | |
| i | Trifolium repens L., Sp. Pl. 2: 767 (1753) | 1:158 |
| i | Trifolium resupinatum L., Sp. Pl. 2: 771 (1753) | 1:158 |
| i | Trifolium scabrum L., Sp. Pl. 2: 770 (1753) | 1:160 |
| i | Trifolium squamosum L., Amoen. Acad., Linnaeus ed. 4, 105 (1759) | 1:161 |
| i | Trifolium stellatum L., Sp. Pl. 2: 769 (1753) | 1:161 |
| i | Trifolium striatum L., Sp. Pl. 2: 770 (1753) | 1:160 |
| i | Trifolium subterraneum L., Sp. Pl. 2: 767 (1753) | 1:159 |
| i | Trifolium suffocatum L., Mant. Pl. 2: 276 (1771) | 1:159 |
| i | Trifolium tomentosum L., Sp. Pl. 2: 771 (1753) | 1:158 |
| | Trifolium uniflorum L. previously listed as naturalised but insufficient evidence exists to support this | |

| | | |
|----------------------|--|---------------|
| i | Trifolium vesiculosum Savi, Fl. Pis. 2: 165 (1798) | |
| | Trigonella ornithopodioides (L.) DC. = Trifolium ornithopodioides | 1:152 |
| i | Ulex europaeus L., Sp. Pl. 2: 741 (1753) | 1:150 |
| | Vicia angustifolia L. = Vicia sativa subsp. nigra | 1:167 |
| | Vicia cracca L. sensu Curtis & Morris (1975) = Vicia villosa (misapplied in Tasmania) | 1:167 |
| i | Vicia hirsuta (L.) Gray, Nat. Arr. Brit. Pl. 2: 614 (1821) | 1:166 |
| | Vicia lathyroides L. recorded in error | |
| i | Vicia sativa L. subsp. nigra (L.) Ehrh., Hannover. Mag. 15: 229 (1780) | 1:167 |
| i | Vicia sativa L. subsp. sativa , Sp. Pl. 2: 736 (1753) | 1:167 |
| i | Vicia tetrasperma (L.) Schreb., Spic. Fl. Lips. 26 (1771) | 1:167 |
| i | Vicia villosa Roth subsp. eriocarpa (Hauskn.) P.W.Ball, Feddes Repert. 79: 45 (1968) | 1:167 |
| | Viminaria denudata (Vent.) Sm. = Viminaria juncea | |
| | Viminaria juncea (Schrad. & J.C.Wendl.) Hoffmanns., Verz. Pfl.-Kult. 200 (1824) | 1:136 |
| FRANKENIACEAE | | FTO 93 |
| | Frankenia pauciflora DC. var. gunnii Summerh., J. Linn. Soc. Bot. 48: 366 (1930) | 1:64 |
| FUMARIACEAE | | FTO 45 |
| i | Fumaria bastardii Boreau, Rev. Bot. Recueil Mans. 2: 359 (1847) | 1:30 |
| i | Fumaria densiflora DC., Cat. Pl. Horti Monsp. 113 (1813) | 1:30 |
| i | Fumaria muralis Sond. ex W.D.J.Koch subsp. muralis , Syn. Fl. Germ. Helv., ed. 2: 1017 (1845) | 1:30 |
| i* | Fumaria officinalis L. subsp. officinalis , Sp. Pl. 2: 700 (1753) | |
| | <i>Pseudofumaria alba</i> (Mill.) Lidén subsp. <i>alba</i> previously listed as naturalised but insufficient evidence exists to support this | |
| GENTIANACEAE | | |
| | <i>Centaurium australe</i> (R.Br.) Druce = Schenkia australis | 3:479 |
| i | Centaurium erythraea Rafn, Danm. Holst. Fl. 2: 75 (1800) | 3:480 |
| | <i>Centaurium pulchellum</i> (Sw.) Druce sensu Curtis (1967) = Centaurium tenuiflorum (misapplied in Tasmania) | 3:480 |
| | <i>Centaurium spicatum</i> (L.) Fritsch ex Janch. sensu Buchanan (1999) = Schenkia australis (misapplied in Tasmania) | 3:479 |
| i | Centaurium tenuiflorum (Hoffmanns. & Link) Fritsch ex Janch., Mitt. Naturwiss. Vereins Univ. Wien 5: 97 (1907) | 3:480 |
| | <i>Chionogentias brevisepala</i> L.G.Adams = Gentianella brevisepala | 3:482 |
| | <i>Chionogentias cunninghamii</i> L.G.Adams = Gentianella cunninghamii | 3:482 |
| | <i>Chionogentias demissa</i> L.G.Adams = Gentianella demissa | 3:482 |
| | <i>Chionogentias diemensis</i> (Griseb.) L.G.Adams = Gentianella diemensis | 3:482 |
| | <i>Chionogentias eichleri</i> L.G.Adams = Gentianella eichleri | 3:482 |
| e ?x | Chionogentias grandis L.G.Adams, Austral. Syst. Bot. 8: 960 (1995) | 3:482 |
| | <i>Chionogentias gunniana</i> L.G.Adams = Gentianella gunniana | 3:482 |
| | <i>Chionogentias pleurogynoides</i> (Griseb.) L.G.Adams = Gentianella pleurogynoides | 3:482 |
| | <i>Chionogentias polysperes</i> L.G.Adams = Gentianella polysperes | 3:482 |
| i | Cicendia filiformis (L.) Delarbre, Fl. Auvergne (Delarbre), ed. 2, 1: 29 (1800) | 3:481 |
| | <i>Erythraea australis</i> R.Br. = Schenkia australis | |
| | <i>Exacum ovatum</i> Labill. = Sebaea ovata | |

| | | |
|-----|--|---------------|
| | Gentiana montana G.Forst. sensu Brown (1810) = Gentianella pleurogynoides (misapplied in Tasmania) | |
| | Gentiana montana G.Forst. sensu Hooker (1860) = Gentianella gunniana (misapplied in Tasmania) | |
| | Gentiana saxosa G.Forst. sensu Rodway (1903) a name of uncertain application (= Gentianella spp.) | |
| e | Gentianella brevisepala (L.G.Adams) Glenny, New Zealand J. Bot. 42: 518 (2004) | 3:482 |
| x | Gentianella cunninghamii (L.G.Adams) Glenny subsp. cunninghamii , New Zealand J. Bot. 42: 518 (2004) | 3:482 |
| e | Gentianella demissa (L.G.Adams) Glenny, Muelleria 26: 95 (2008) | 3:482 |
| e | Gentianella diemensis (Griseb.) J.H.Willis subsp. diemensis , Vict. Naturalist 73: 199 (1957) | 3:482 |
| e | Gentianella diemensis (Griseb.) J.H.Willis subsp. plantaginea (L.G.Adams) Glenny, New Zealand J. Bot. 42: 519 (2004) | 3:482 |
| e | Gentianella eichleri (L.G.Adams) Glenny, New Zealand J. Bot. 42: 519 (2004) | 3:482 |
| | Gentianella grandis (L.G.Adams) Glenny nom. illeg. non Sm. (1936) = Chionogentias grandis | 3:482 |
| | Gentianella gunniana (L.G.Adams) Glenny, New Zealand J. Bot. 42: 519 (2004) | 3:482 |
| e | Gentianella pleurogynoides (Griseb.) Glenny subsp. milliganii (L.G.Adams) Glenny, New Zealand J. Bot. 42: 519 (2004) | 3:482 |
| e | Gentianella pleurogynoides (Griseb.) Glenny subsp. pleurogynoides , New Zealand J. Bot. 42: 519 (2004) | 3:482 |
| | Gentianella polysperes (L.G.Adams) Glenny, New Zealand J. Bot. 42: 519 (2004) | 3:482 |
| | Schenkia australis (R.Br.) G.Mans., Taxon 53: 725 (2004) | 3:479 |
| | Sebaea albidiflora F.Muell., Trans. Philos. Soc. Victoria 1: 46 (1855) | 3:479 |
| | Sebaea ovata (Labill.) R.Br., Prodr. Fl. Nov. Holland. 452 (1810) | 3:478 |
| | GERANIACEAE | FTO 54 |
| i | Erodium botrys (Cav.) Bertol., Amoen. Ital. 35 (1819) | 1:98 |
| i | Erodium cicutarium (L.) L'Hér. ex Aiton, Hortus Kew. (W.Aiton) 2: 414 (1789) | 1:97 |
| i * | Erodium malacoides (L.) L'Hér. ex Aiton, Hortus Kew. (W.Aiton) 2: 415 (1789) | 1:98 |
| i | Erodium moschatum (L.) L'Hér. ex Aiton, Hortus Kew. (W.Aiton) 2: 414 (1789) | 1:97 |
| | Geranium brevicaule Hook., J. Bot. (Hooker) 1: 252 (1834) | |
| i | Geranium dissectum L., Cent. Pl. 1: 21 (1755) | 1:96 |
| | Geranium dissectum L. var. australe Benth. sensu Bentham (1863) = G. potentilloides , G. retrorsum and G. solanderi | |
| | Geranium dissectum L. var. pilosum Hook.f. = Geranium solanderi | |
| | Geranium dissectum L. var. potentilloides Benth. = Geranium potentilloides | |
| ?i | Geranium homeanum Turcz., Bull. Soc. Imp. Naturalistes Moscou 36: 591 (1863) | |
| | Geranium microphyllum Hook.f. = Geranium potentilloides var. potentilloides | |
| i | Geranium molle L., Sp. Pl. 2: 682 (1753) | 1:96 |
| | Geranium sp. Pale Pink Flowers (M.Gray 5847) Vic. Herbarium | |
| | Geranium pilosum Sol. ex Willd. nom. illeg. = Geranium solanderi | |
| | Geranium potentilloides L'Hér. ex DC. var. potentilloides , Prodr. [A. P. de Candolle] 1: 639 (1824) | 1:95 |
| | Geranium retrorsum L'Hér. ex DC., Prodr. [A. P. de Candolle] 1: 644 (1824) | |
| i t | Geranium rotundifolium L., Sp. Pl. 2: 683 (1753) | 1:96 |
| | Geranium sessiliflorum Cav. sensu Rodway (1903) = Geranium brevicaule (misapplied in Tasmania) | |
| | Geranium sessiliflorum Cav. subsp. brevicaule (Hook.) Carolin = Geranium brevicaule | 1:95 |

| | | |
|-------|---|----------------|
| | Geranium solanderi Carolin, Proc. Linn. Soc. New South Wales 89: 350 (1965) | 1:95 |
| i | Geranium yeoi Aedo & Muñoz Garm., Kew Bull. 52: 727 (1997) | |
| i | Pelargonium × asperum Ehrh. ex Willd., Sp. Pl., ed. 4 [Willdenow] 3: 678 (1800) | 1:99 |
| | Pelargonium australe Willd., Sp. Pl., ed. 4 [Willdenow] 3: 675 (1800) | 1:98 |
| i | Pelargonium capitatum (L.) L'Hér. ex Aiton, Hortus Kew. (W.Aiton)2: 425 (1789) | |
| i | Pelargonium × domesticum L.H.Bailey, Stand. Cycl. Hort. 2532 (1916) | 1:99 |
| | Pelargonium inodorum Willd., Enum. Pl. [Willdenow] 2: 702 (1809) | 1:99 |
| | Pelargonium littorale Hugel, Bot. Arch.(Hügel): t.5 (1837) | |
| | GOODENIACEAE | FTO 129 |
| | Brunonia australis Sm. ex R.Br., Prodr. Fl. Nov. Holland. 590 (1810) | 2:405 |
| x | Cooperookia barbata (R.Br.) Carolin, Proc. Linn. Soc. New South Wales 92: 213 (1968) | 2:401 |
| | Dampiera stricta (Sm.) R.Br., Prodr. Fl. Nov. Holland. 589 (1810) | 2:404 |
| | <i>Goodenia amplexans</i> F.Muell. sensu Curtis (1963) recorded in error | 2:400 |
| | <i>Goodenia barbata</i> R.Br. = Cooperookia barbata | 2:401 |
| | Goodenia elongata Labill., Nov. Holl. Pl. 1: 52, t.75 (1805) | 2:401 |
| | Goodenia geniculata R.Br., Prodr. Fl. Nov. Holland. 577 (1810) | 2:401 |
| | <i>Goodenia geniculata</i> R.Br. var. <i>lanata</i> (R.Br.) Rodway = Goodenia lanata | |
| | <i>Goodenia hederacea</i> Sm. sensu Brown (1810), Hooker (1860) = Goodenia lanata (misapplied in Tasmania) | |
| | Goodenia humilis R.Br., Prodr. Fl. Nov. Holland. 575 (1810) | 2:402 |
| | Goodenia lanata R.Br., Prodr. Fl. Nov. Holland. 577 (1810) | 2:401 |
| | Goodenia ovata Sm., Trans. Linn. Soc. London 2: 347 (1794) | 2:400 |
| x | Goodenia pinnatifida Schlttdl., Linnaea 21: 450 (1848) | |
| | <i>Goodenia repens</i> Labill. = Selliera radicans | |
| | Scaevola aemula R.Br., Prodr. Fl. Nov. Holland. 584 (1810) | 2:404 |
| | Scaevola albida (Sm.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916, Suppl. 2: 644 (1917) | 2:404 |
| | <i>Scaevola calendulacea</i> (Andrews) Druce sensu Curtis (1963) recorded in error | |
| | <i>Scaevola cuneiformis</i> Labill. recorded in error | |
| | Scaevola hookeri (de Vriese) F.Muell. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 231 (1856) | 2:403 |
| | <i>Scaevola microcarpa</i> Cav. = Scaevola albida | |
| | Selliera radicans Cav., Anales. Hist. Nat. 1: 41, t.5 (1799) | 2:402 |
| | Velleia montana Hook.f., London J. Bot. 6: 265 (1847) | 2:399 |
| | Velleia paradoxa R.Br., Prodr. Fl. Nov. Holland. 580 (1810) | 2:398 |
| | GRISELINIACEAE | FTO 136 |
| i # t | Griselinia littoralis (Raoul) Raoul, Choix Pl. Nouv.-Zél. 22, t.19 (1846) | |
| | GROSSULARIACEAE | FTO 50 |
| i t | Ribes sanguineum Pursh, Fl. Amer. Sept. (Pursh) 1: 164 (1813) | |
| | GUNNERACEAE | FTO 49 |
| e | Gunnera cordifolia (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 125 (1856) | 1:194 |
| | GYROSTEMONACEAE | FTO 83 |
| | <i>Didymotheca thesioides</i> Hook.f. = Gyrostemon thesioides | 3:581 |
| | Gyrostemon thesioides (Hook.f.) A.S.George, Fl. Australia 8: 392 (1982) | 3:581 |

HALORAGACEAE

| | | |
|----|--|-------|
| | Gonocarpus humilis Orchard, Bull. Auckland Inst. Mus. 10: 195 (1975) | |
| | Gonocarpus micranthus Thunb. subsp. micranthus , Nov. Gen. Pl. [Thunberg] 3: 55 (1783) | 1:190 |
| | Gonocarpus montanus (Hook.f.) Orchard, Bull. Auckland Inst. Mus. 10: 172 (1975) | 1:192 |
| | Gonocarpus serpyllifolius Hook.f., Icon. Pl. 3: t.290 (1840) | 1:192 |
| | Gonocarpus tetragynus Labill., Nov. Holl. Pl. 1: 39, t.53 (1805) | 1:191 |
| | Gonocarpus teucroides DC., Prodr. [A. P. de Candolle] 3: 66 (1828) | 1:191 |
| | Haloragis aspera Lindl., J. Exped. Trop. Australia [Mitchell]: 306 (1848) | 1:191 |
| | Haloragis brownii (Hook.f.) Schindl. = Meionectes brownii | 1:192 |
| | Haloragis ceratophylla Endl. sensu Bentham (1864) = Haloragis aspera (misapplied in Tasmania) | |
| | Haloragis depressa (A.Cunn.) Walp. sensu Bentham (1864) = Gonocarpus montanus (misapplied in Tasmania) | |
| | Haloragis depressa (A.Cunn.) Walp. sensu Rodway (1903) = Gonocarpus serpyllifolius (misapplied in Tasmania) | |
| | Haloragis depressa (A.Cunn.) Walp. var. montana (Hook.f.) Hook.f. = Gonocarpus montanus | |
| | Haloragis gunnii Hook.f. = Gonocarpus teucroides | |
| | Haloragis heterophylla Brongn., Voy. Monde, Atlas: t.68A (1826) | 1:190 |
| | Haloragis micrantha (Thunb.) R.Br. ex Sieber & Zucc. = Gonocarpus micranthus | 1:190 |
| | Haloragis montana Hook.f. = Gonocarpus montanus | 1:192 |
| | Haloragis myriocarpa Orchard, Bull. Auckland Inst. Mus. 10: 132 (1975) | |
| | Haloragis pinnatifida A.Gray = Haloragis aspera | |
| | Haloragis serpyllifolia (Hook.f.) Walp. = Gonocarpus serpyllifolius | 1:192 |
| | Haloragis tetragyna (Labill.) Hook.f. = Gonocarpus tetragynus | 1:191 |
| | Haloragis teucroides (DC.) Schtdl. = Gonocarpus teucroides | 1:191 |
| | Meionectes brownii Hook.f., Icon. Pl. 4: t.306 (1841) | 1:192 |
| | Myriophyllum amphibium Labill., Nov. Holl. Pl. 2: 70, t.220 (1806) | 1:193 |
| i | Myriophyllum aquaticum (Vell.) Verdc., Kew Bull. 28: 36 (1973) | 1:193 |
| e | Myriophyllum austropygmaeum Orchard, Brunonia 8: 280 (1986) | |
| | Myriophyllum brasiliense Cambess. = Myriophyllum aquaticum | 1:193 |
| ?i | Myriophyllum crispatum Orchard, Brunonia 8: 210 (1986) | |
| | Myriophyllum elatinoides Gaudich. sensu Curtis (1956), Curtis & Morris (1975) = Myriophyllum salsugineum (misapplied in Tasmania) | 1:193 |
| x | Myriophyllum glomeratum Schindl., Pflanzenr. (Engler) 23: 103 (1905) | |
| | Myriophyllum integrifolium (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 123 (1856) | 1:194 |
| | Myriophyllum muelleri Sond., Linnaea 28: 233 (1856) | |
| | Myriophyllum pedunculatum Hook.f. subsp. longibracteolatum (Schindl.) Orchard, Brunonia 8: 273 (1986) | 1:194 |
| | Myriophyllum pedunculatum Hook.f. subsp. pedunculatum , London J. Bot. 6: 474 (1847) | 1:194 |
| | Myriophyllum propinquum A.Cunn. sensu Curtis (1956), Curtis & Morris (1975) = Myriophyllum simulans (misapplied in Tasmania) | 1:193 |
| | Myriophyllum propinquum A.Cunn. sensu Curtis (1956), Curtis & Morris (1975) = Myriophyllum variifolium (misapplied in Tasmania) | |
| | Myriophyllum salsugineum Orchard, Brunonia 4: 48 (1981) | 1:193 |
| | Myriophyllum simulans Orchard, Brunonia 8: 203 (1986) | 1:193 |

| | | |
|-----------------------------|---|-------|
| | Myriophyllum variifolium Hook.f., Icon. Pl. 3: t.289 (1840) | 1:193 |
| HYPERICACEAE | | |
| | Ascyrum humifusum Labill. = Hypericum pusillum | |
| | Ascyrum involutum Labill. = Hypericum gramineum | |
| i | Hypericum androsaemum L., Sp. Pl. 2: 784 (1753) | 1:82 |
| i | Hypericum calycinum L., Mant. Pl. 1: 106 (1767) | 1:82 |
| | Hypericum gramineum G.Forst., Fl. Ins. Austr. 53 (1786) | 1:81 |
| | Hypericum humifusum L. previously listed as naturalised but insufficient evidence exists to support this | |
| | Hypericum japonicum Thunb., Fl. Jap. (Thunberg): 295 (1784) | 1:81 |
| i | Hypericum perforatum L. subsp. veronense (Schrank) H.Lindb., Öfvers. Finska Vetensk.-Soc. Förh. 48: 73 (1906) | 1:81 |
| i # t | Hypericum pulchrum L., Sp. Pl. 2: 786 (1753) | |
| t | Hypericum pusillum Choisy, Prodr. Monogr. Hyperic. 50 (1821) | |
| i | Hypericum tetrapterum Fr. var. tetrapterum , Novit. Fl. Suec. Alt. 236 (1828) | |
| LAMIACEAE (LABIATAE) | | |
| | Ajuga australis R.Br., Prodr. Fl. Nov. Holland. 503 (1810) | 3:557 |
| i | Ajuga reptans L., Sp. Pl. 2: 561 (1753) | |
| i | Cedronella canariensis (L.) Webb & Berthel., Hist. Nat. Illes Canarias (Phytogr.) 3: 87 (1845) | 3:548 |
| i | Lamium amplexicaule L., Sp. Pl. 2: 579 (1753) | 3:552 |
| i | Lamium purpureum L., Sp. Pl. 2: 579 (1753) | 3:552 |
| i | Lavandula stoechas L. subsp. stoechas , Sp. Pl. 2: 573 (1753) | |
| i | Leonotis leonurus (L.) R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 3: 410 (1811) | |
| | Lycopus australis R.Br., Prodr. Fl. Nov. Holland. 500 (1810) | 3:546 |
| i | Marrubium vulgare L., Sp. Pl. 2: 583 (1753) | 3:550 |
| i | Melissa officinalis L., Sp. Pl. 2: 592 (1753) | 3:546 |
| | Mentha australis R.Br., Prodr. Fl. Nov. Holland. 505 (1810) | 3:544 |
| | Mentha diemenica Spreng. var. diemenica , Syst. Veg. (ed. 16) [Sprengel] 2: 724 (1825) | 3:544 |
| | Mentha diemenica Spreng. var. serpyllifolia (Benth.) J.H.Willis, Muellera 1: 144 (1967) | 3:544 |
| | Mentha gracilis R.Br. nom. illeg. = Mentha diemenica | |
| i | Mentha × piperita L., Sp. Pl. 2: 576 (1753) | 3:545 |
| i | Mentha pulegium L., Sp. Pl. 2: 577 (1753) | 3:545 |
| | Mentha serpyllifolia Benth. = Mentha diemenica var. serpyllifolia | |
| i # | Mentha spicata L., Sp. Pl. 2: 576 (1753) | 3:545 |
| i | Nepeta cataria L., Sp. Pl. 2: 570 (1753) | 3:548 |
| x | Prostanthera cuneata Benth., Prodr. [A. P. de Candolle] 12: 560 (1848) | 3:554 |
| | Prostanthera lasianthos Labill. var. lasianthos , Nov. Holl. Pl. 2: 18 t.157 (1806) | 3:553 |
| | Prostanthera retusa R.Br. = Prostanthera rotundifolia | |
| | Prostanthera rotundifolia R.Br., Prodr. Fl. Nov. Holland. 509 (1810) | 3:553 |
| i | Prunella laciniata (L.) L., Sp. Pl., ed. 2, 2: 837 (1763) | 3:550 |
| i | Prunella vulgaris L., Sp. Pl. 2: 600 (1753) | 3:550 |
| | Salvia horminoides Pourr. = Salvia verbenaca | 3:547 |
| i | Salvia verbenaca L. var. verbenaca , Sp. Pl. 1: 25 (1753) | 3:547 |
| | Scutellaria humilis R.Br., Prodr. Fl. Nov. Holland. 507 (1810) | 3:549 |

| | | |
|-----|--|----------------|
| i | Stachys arvensis (L.) L., Sp. Pl., ed. 2, 2: 814 (1763) | 3:551 |
| i t | Stachys palustris L., Sp. Pl. 2: 580 (1753) | 3:551 |
| | Teucrium corymbosum R.Br., Prodr. Fl. Nov. Holland. 504 (1810) | 3:556 |
| i t | Teucrium scorodonia L., Sp. Pl. 2: 564 (1753) | |
| e | Westringia angustifolia R.Br., Prodr. Fl. Nov. Holland. 501 (1810) | 3:555 |
| e | Westringia brevifolia Benth., Labiat. Gen. Spec. 459 (1834) | 3:555 |
| | Westringia brevifolia Benth. var. raleighii (B.Boivin) W.M.Curtis = Westringia brevifolia | |
| | Westringia dampieri R.Br. sensu Bentham (1870) misapplied to W. angustifolia or W. rigida | |
| | Westringia rigida R.Br., Prodr. Fl. Nov. Holland. 501 (1810) | 3:555 |
| e | Westringia rubiifolia R.Br., Prodr. Fl. Nov. Holland. 501 (1810) | 3:556 |
| | LENTIBULARIACEAE | FTO 119 |
| | Polypompholyx tenella (R.Br.) Lehm. = Utricularia tenella | 3:537 |
| | Utricularia australis R.Br., Prodr. Fl. Nov. Holland. 430 (1810) | 3:536 |
| | Utricularia barkeri R.W.Jobson, Telopea 15: 131 (2013) | |
| | Utricularia dichotoma Labill., Nov. Holl. Pl. 1: 11, t.8 (1805) | 3:536 |
| | Utricularia dichotoma Labill. var. uniflora (R.Br.) Benth. = Utricularia uniflora | |
| | Utricularia flexuosa Vahl sensu Rodway (1903) = Utricularia australis (misapplied in Tasmania) | 3:536 |
| i * | Utricularia gibba L., Sp. Pl. 1: 18 (1753) | |
| | Utricularia lateriflora R.Br., Prodr. Fl. Nov. Holland. 431 (1810) | 3:536 |
| | Utricularia monanthos Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 299 (1857) | 3:537 |
| | Utricularia tenella R.Br., Prodr. Fl. Nov. Holland. 432 (1810) | 3:537 |
| | Utricularia uniflora R.Br., Prodr. Fl. Nov. Holland. 431 (1810) | 3:536 |
| | Utricularia violacea R.Br., Prodr. Fl. Nov. Holland. 431 (1810) | |
| | LINACEAE | FTO 77 |
| | Linum albidum Ewart & Jean White = Linum marginale | |
| i t | Linum bienne Mill., Gard. Dict., ed. 8, no.8 (1768) | |
| i t | Linum catharticum L., Sp. Pl. 1: 281 (1753) | 1:92 |
| | Linum gallicum L. = Linum trigynum | |
| | Linum marginale A.Cunn., Geogr. Mem. New South Wales [Field]: 357 (1825) | 1:92 |
| i | Linum trigynum L., Sp. Pl. 1: 279 (1753) | 1:93 |
| i | Linum usitatissimum L., Sp. Pl. 1: 277 (1753) | 1:92 |
| | LOGANIACEAE | |
| | Mitrasacme archeri Hook.f. = Schizacme archeri | 3:475 |
| | Mitrasacme distylis F.Muell. = Phyllangium distylis | 3:477 |
| | Mitrasacme divergens Hook.f. = Phyllangium divergens | 3:476 |
| | Mitrasacme montana Hook.f. ex Benth. = Schizacme montana | 3:475 |
| | Mitrasacme paradoxa R.Br. sensu Curtis (1967) = Phyllangium divergens (misapplied in Tasmania) | 3:476 |
| | Mitrasacme pilosa Labill. var. pilosa , Nov. Holl. Pl. 1: 36, t.49 (1805) | 3:476 |
| | Mitrasacme pilosa Labill. var. stuartii Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 274 (1857) | 3:476 |
| | Mitrasacme serpyllifolia R.Br., Prodr. Fl. Nov. Holland. 454 (1810) | 3:475 |
| | Phyllangium distylis (F.Muell.) Dunlop, Fl. Australia 28: 315 (1996) | 3:477 |
| | Phyllangium divergens (Hook.f.) Dunlop, Fl. Australia 28: 315 (1996) | 3:476 |
| e | Schizacme archeri (Hook.f.) Dunlop, Fl. Australia 28: 314 (1996) | 3:475 |

| | | |
|----------------------|--|----------------|
| | Schizacme montana (Hook.f. ex Benth.) Dunlop, Fl. Australia 28: 314 (1996) | 3:475 |
| LYTHRACEAE | | FTO 55 |
| | Lythrum hyssopifolia L., Sp. Pl. 1: 447 (1753) | 2:227 |
| | Lythrum salicaria L., Sp. Pl. 1: 446 (1753) | 2:226 |
| MALVACEAE | | |
| i t | Anisodonteia capensis (L.) D.M.Bates, Gentes Herb. 10: 327 (1969) | |
| e | Asterotrichion discolor (Hook.) Melville, Kew Bull. 20: 512 (1967) | 1:86 |
| | Gynatrix pulchella (Willd.) Alef., Oesterr. Bot. Z. 12: 35 (1862) | 1:87 |
| | <i>Hibiscus trionum</i> L. previously listed as naturalised but insufficient evidence exists to support this | 1:88 |
| | Lasiopetalum baueri Steetz, Pl. Preiss. [J.G.C.Lehman] 2: 339 (1848) | 1:90 |
| | <i>Lasiopetalum dasyphyllum</i> Sieber ex Steetz = Lasiopetalum macrophyllum | 1:89 |
| | Lasiopetalum discolor Hook., Companion Bot. Mag. 1: 276 (1836) | 1:89 |
| | Lasiopetalum macrophyllum Graham, Edinburg New Philos. J. 31: 390 (1841) | 1:89 |
| e | Lasiopetalum micranthum Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 51 (1855) | 1:90 |
| | <i>Lavatera arborea</i> L. = Malva arborea | 1:84 |
| | <i>Lavatera cretica</i> L. = Malva pseudolavatera has also been missapplied to M. preissiana in Curtis & Morris (1975) | 1:84 |
| | <i>Lavatera plebeia</i> Sims var. <i>tomentosa</i> Hook.f. = Malva preissiana | 1:83 |
| | Lawrencia spicata Hook., Icon. Pl. 3: t.261 (1840) | 1:88 |
| ?i # | Lawrencia squamata Nees, Pl. Preiss. [J.G.C.Lehman] 1: 242 (1845) | |
| i | Malva arborea (L.) Webb & Berthel., Hist. Nat. Iles Canaries (Phytogr.) 1: 30 (1836) | 1:84 |
| | <i>Malva australiana</i> M.F.Ray = Malva preissiana | 1:83 |
| | <i>Malva dendromorpha</i> M.F.Ray nom. illeg., nom. superfl. = Malva arborea | |
| | <i>Malva linnaei</i> M.F.Ray nom. illeg., nom. superfl. = Malva pseudolavatera | |
| i | Malva moschata L., Sp. Pl. 2: 690 (1753) | 1:85 |
| i | Malva neglecta Wallr., Syll. Pl. Nov. 1: 140 (1824) | 1:85 |
| i | Malva nicaeensis All., Fl. Pedem. 2: 40 (1785) | 1:85 |
| i | Malva parviflora L., Demonstr. Pl. 18 (1753) | 1:85 |
| | Malva preissiana Miq., Pl. Preiss. [J.G.C.Lehman] 1: 238 (1845) | 1:83 |
| i | Malva pseudolavatera Webb & Berthel., Hist. Nat. Iles Canaries (Phytogr.) 3(2(1)): 29 (1836) | 1:84 |
| | <i>Malva rotundifolia</i> L. sensu Rodway (1903) = Malva neglecta (misapplied in Tasmania) | |
| i | Malva sylvestris L., Sp. Pl. 2: 689 (1753) | 1:85 |
| i | Modiola caroliniana (L.) G.Don, Gen. Hist. 1: 466 (1831) | 1:86 |
| | <i>Plagianthus pulchellus</i> (Willd.) A.Gray ex Hook.f. = Gynatrix pulchella | 1:87 |
| | <i>Plagianthus pulchellus</i> (Willd.) Hook.f. var. <i>tomentosus</i> Rodway = Gynatrix pulchella | |
| | <i>Plagianthus sidoides</i> Hook. = Asterotrichion discolor | 1:86 |
| | <i>Plagianthus spicatus</i> (Hook.) Benth. = Lawrencia spicata | 1:88 |
| MAZACEAE | | |
| | Mazus pumilio R.Br., Prodr. Fl. Nov. Holland. 439 (1810) | 3:516 |
| MENYANTHACEAE | | FTO 128 |
| | <i>Limnanthemum exaltatum</i> (Sol. ex Sims) F.Muell. = Liparophyllum exaltatum | |
| | <i>Limnanthemum exiguum</i> F.Muell. = Liparophyllum exiguum | |
| | <i>Limnanthemum gunnii</i> (Hook.f.) Hook.f. = Liparophyllum gunnii | |

| | | |
|--------------------|--|-------|
| | Liparophyllum exaltatum (Sol. ex Sims) Tippet & Les, Novon 19: 408 (2009) | 3:483 |
| e | Liparophyllum exiguum (F.Muell.) Tippet & Les, Novon 19: 408 (2009) | 3:483 |
| t | Liparophyllum gunnii Hook.f., London J. Bot. 6: 473 bis (1847) | 3:484 |
| | Nymphoides crenata (F.Muell.) Kuntze sensu Curtis (1967) recorded in error | 3:483 |
| | Nymphoides exigua (F.Muell.) Kuntze = Liparophyllum exiguum | 3:483 |
| | Ornduffia reniformis (R.Br.) Tippet & Les, Novon 19: 410 (2009) | 3:483 |
| | Ornduffia umbricola (Aston) Tippet & Les var. umbricola , Novon 19: 410 (2009) | |
| | Villarsia exaltata (Sol. ex Sims) G.Don = Liparophyllum exaltatum | 3:483 |
| | Villarsia parnassiifolia (Labill.) R.Br. sensu Brown (1810), Hooker (1860) = Liparophyllum exaltatum (misapplied in Tasmania) | |
| | Villarsia reniformis R.Br. = Ornduffia reniformis | 3:483 |
| | Villarsia umbricola Aston = Ornduffia umbricola | |
| MYRSINACEAE | | |
| | Anagallis arvensis L. = Lysimachia arvensis | 3:467 |
| | Anagallis arvensis L. subsp. foemina sensu Curtis (1967) = Lysimachia arvensis (misapplied in Tasmania) | 3:468 |
| | Anagallis arvensis L. var. caerulea (L.) Gouan = Lysimachia arvensis | 3:468 |
| i | Lysimachia arvensis (L.) U.Manns & Anderb., Willdenowia 39: 51 (2009) | 3:467 |
| i* | Lysimachia minima (L.) U.Manns & Anderb., Willdenowia 39: 52 (2009) | |
| i | Lysimachia nummularia L., Sp. Pl. 1: 148 (1753) | 3:467 |
| MYRTACEAE | | |
| FTO 57 | | |
| | Baeckea diffusa Sieber ex DC. = Euryomyrtus ramosissima | |
| | Baeckea gunniana Schauer, Repert. Bot. Syst. (Walpers) 2: 920 (1843) | 1:198 |
| e | Baeckea leptocaulis Hook.f., Icon. Pl. 3: t.298 (1840) | 1:199 |
| | Baeckea ramosissima A.Cunn. = Euryomyrtus ramosissima | 1:198 |
| | Baeckea thymifolia Hook.f. = Euryomyrtus ramosissima | |
| | Callistemon pallidus (Bonpl.) DC. = Melaleuca pallida | 1:203 |
| | Callistemon paludosus F.Muell. sensu Curtis (1956), Curtis & Morris (1975) = Melaleuca pallida (misapplied in Tasmania) | 1:203 |
| | Callistemon salignus (Sm.) Colvill ex Sweet sensu Benth (1867) = Melaleuca pallida (misapplied in Tasmania) | |
| | Callistemon salignus (Sm.) Colvill ex Sweet var. australis Benth. = Melaleuca pallida | |
| | Callistemon salignus (Sm.) Colvill ex Sweet var. hebestachys Benth. = Melaleuca pallida | |
| | Callistemon salignus (Sm.) Colvill ex Sweet var. viridiflora Benth. = Melaleuca virens | |
| | Callistemon viridiflorus (Sims) Sweet = Melaleuca virens | 1:203 |
| | Calycothrix glabra (R.Br.) Hook.f. = Calytrix tetragona | |
| | Calytrix tetragona Labill., Nov. Holl. Pl. 2: 8, t.146 (1806) | 1:197 |
| | Eucalyptus acervula Sieber ex DC. sensu Hooker (1860), Rodway (1903) = Eucalyptus ovata var. ovata (misapplied in Tasmania) | |
| | Eucalyptus aggregata H.Deane & Maiden sensu Curtis (1956) = Eucalyptus rodwayi (misapplied in Tasmania) | |
| | Eucalyptus ambigua DC. a name of uncertain application | |
| e | Eucalyptus amygdalina Labill., Nov. Holl. Pl. 2: 14, t.154 (1806) | 1:224 |
| | Eucalyptus amygdalina Labill. var. nitida (Hook.f.) Benth. = Eucalyptus nitida | |

| | | |
|---|---|-------|
| | Eucalyptus amygdalina Labill. var. radiata (Sieber ex DC.) Benth. = Eucalyptus radiata subsp. radiata | |
| e | Eucalyptus archeri Maiden & Blakely, Crit. Rev. Eucalyptus 8: 58 (1929) | 1:215 |
| e | Eucalyptus barberi L.A.S.Johnson & Blaxell, Contr. N.S.W. Natl. Herb. 4: 288 (1972) | 1:211 |
| | Eucalyptus bicostata Maiden, Blakely & Simmonds sensu Curtis & Morris (1975) = Eucalyptus globulus subsp. pseudoglobulus (misapplied in Tasmania) | 1:219 |
| | Eucalyptus brookeriana A.M.Gray, Austral. Forest Res. 9: 111 (1979) | |
| e | Eucalyptus coccifera Hook.f., London J. Bot. 6: 477 (1847) | 1:225 |
| | Eucalyptus coccifera Hook.f. var. parviflora Benth. = Eucalyptus coccifera | |
| e | Eucalyptus cordata Labill. subsp. cordata , Nov. Holl. Pl. 2: 13, t.152 (1806) | 1:218 |
| e | Eucalyptus cordata Labill. subsp. quadrangulosa D.Nicolle, B.M.Potts & McKinnon, Pap. & Proc. Roy. Soc. Tasmania 142: 75 (2008) | |
| | Eucalyptus coriacea Schauer = Eucalyptus pauciflora subsp. pauciflora | |
| | Eucalyptus dalrympleana Maiden subsp. dalrympleana , Forest Fl. N.S.W. 7: 137, t.241 (1920) | 1:214 |
| e | Eucalyptus delegatensis R.T.Baker subsp. tasmaniensis Boland, Austral. Forest Res. 15: 177 (1985) | 1:220 |
| | Eucalyptus gigantea Hook.f. = Eucalyptus delegatensis subsp. tasmaniensis | 1:220 |
| | Eucalyptus globulus Labill. subsp. globulus , Voy. Rech. Prouse 1: 153, t.13 (1800) | 1:218 |
| | Eucalyptus globulus Labill. subsp. pseudoglobulus (Naudin) J.B.Kirkp., Bot. J. Linn. Soc. 69: 101 (1975) | 1:219 |
| e | Eucalyptus gunnii Hook.f. subsp. divaricata (McAulay & Brett) B.M.Potts, Pap. & Proc. Roy. Soc. Tasmania 135: 57 (2001) | |
| e | Eucalyptus gunnii Hook.f. subsp. gunnii , London J. Bot. 3: 499 (1844) | 1:215 |
| | Eucalyptus haemastoma Sm. sensu Rodway (1903) misapplied to a hybrid involving Eucalyptus sieberi | |
| e | Eucalyptus johnstonii Maiden, Crit. Rev. Eucalyptus 6: 280 (1922) | 1:219 |
| | Eucalyptus linearis Dehnh. = Eucalyptus pulchella | 1:224 |
| e | Eucalyptus morrisbyi Brett, Pap. & Proc. Roy. Soc. Tasmania 129: pl. 14-15 (1939) | 1:215 |
| | Eucalyptus muelleri T.B.Moore nom. illeg. = Eucalyptus johnstonii | |
| e | Eucalyptus nebulosa A.M.Gray, Kanunnah 3: 42 (2008) | |
| i | Eucalyptus nitens (H.Deane & Maiden) Maiden, Crit. Rev. Eucalyptus 2: 272 (1913) | |
| e | Eucalyptus nitida Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 137, t.29 (1856) | 1:225 |
| | Eucalyptus obliqua L'Hér., Sert. Angl. 18, t.20 (1792) | 1:220 |
| | Eucalyptus ovata Labill. var. ovata , Nov. Holl. Pl. 2: 13, t.153 (1806) | 1:211 |
| | Eucalyptus pauciflora Sieber ex Spreng. subsp. pauciflora , Syst. Veg. (ed. 16) [Sprengel] 4: 195 (1827) | 1:222 |
| | Eucalyptus perriniana F.Muell. ex Rodway, Pap. & Proc. Roy. Soc. Tasmania: 181 (1894) | 1:217 |
| e | Eucalyptus pulchella Desf., Cat. Pl. Horti Paris., ed. 3: 284, 408 (1829) | 1:224 |
| | Eucalyptus radiata Sieber ex DC. subsp. radiata , Prodr. [A. P. de Candolle] 3: 218 (1828) | 1:224 |
| | Eucalyptus radiata Sieber ex DC. subsp. robertsonii sensu Curtis & Morris (1975) = Eucalyptus radiata subsp. radiata (misapplied in Tasmania) | |
| | Eucalyptus regnans F.Muell., Ann. Rep. Victorian Acclim. Soc. 20 in obs. (1871) | 1:220 |
| e | Eucalyptus risdonii Hook.f., London J. Bot. 6: 477 bis (1847) | 1:226 |
| | Eucalyptus risdonii Hook.f. var. elata = Eucalyptus delegatensis subsp. tasmaniensis | |
| e | Eucalyptus rodwayi R.T.Baker & H.G.Sm., Pap. & Proc. Roy. Soc. Tasmania 1912: 191 (1913) | 1:211 |
| | Eucalyptus rubida H.Deane & Maiden subsp. rubida , Proc. Linn. Soc. New South Wales 24: 456, t.xl (1899) | 1:214 |

| | | |
|---|--|-------|
| | <i>Eucalyptus salicifolia</i> Cav. = <i>Eucalyptus amygdalina</i> | 1:224 |
| | <i>Eucalyptus sieberi</i> L.A.S.Johnson, Contr. New South Wales Natl. Herb. 3: 125 (1962) | 1:222 |
| | <i>Eucalyptus sieberiana</i> F.Muell. nom. illeg. = <i>Eucalyptus sieberi</i> | |
| | <i>Eucalyptus simmondsii</i> Maiden = <i>Eucalyptus nitida</i> | 1:225 |
| | <i>Eucalyptus stuartiana</i> F.Muell. ex Miq. = <i>Eucalyptus ovata</i> var. <i>ovata</i> | |
| e | <i>Eucalyptus subcrenulata</i> Maiden & Blakely, Crit. Revis. <i>Eucalyptus</i> 8: 59 (1929) | 1:219 |
| | <i>Eucalyptus tasmanica</i> Blakely = <i>Eucalyptus tenuiramis</i> | 1:226 |
| e | <i>Eucalyptus tenuiramis</i> Miq., Ned. Kruidk. Arch. 4: 128 (1856) | 1:226 |
| e | <i>Eucalyptus urnigera</i> Hook.f., London J. Bot. 6: 477 bis (1847) | 1:217 |
| | <i>Eucalyptus urnigera</i> Hook.f. var. <i>elongata</i> Rodway = <i>Eucalyptus urnigera</i> | |
| e | <i>Eucalyptus vernicosa</i> Hook.f., London J. Bot. 6: 478 bis (1847) | 1:219 |
| e | <i>Eucalyptus viminalis</i> Labill. subsp. <i>hentyensis</i> Brooker & Slee, J. Adelaide Bot. Gard. 21: 92 (2007) | |
| | <i>Eucalyptus viminalis</i> Labill. subsp. <i>viminalis</i> , Nov. Holl. Pl. 2: 12, t.151 (1806) | 1:214 |
| | <i>Euryomyrtus parviflora</i> Miq., Ned. Kruidk. Arch. 4(1): 149 (1856) | |
| | <i>Euryomyrtus ramosissima</i> (A.Cunn.) Trudgen, Nuytsia 13(3): 560 (2001) | 1:198 |
| | <i>Euryomyrtus ramosissima</i> (A.Cunn.) Trudgen subsp. <i>prostrata</i> (Hook.f.) Trudgen = <i>Euryomyrtus parviflora</i> | 1:198 |
| | <i>Fabricia laevigata</i> Gaertn. = <i>Leptospermum laevigatum</i> | |
| | <i>Kunzea ambigua</i> (Sm.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 629 (1917) | 1:202 |
| | <i>Kunzea corifolia</i> (Vent.) Schauer = <i>Kunzea ambigua</i> | |
| i | <i>Kunzea ericoides</i> (A.Rich.) Joy Thomps., Telopea 2: 379 (1983) | |
| | <i>Leptospermum flavescens</i> Sm. sensu Rodway (1903) = <i>Leptospermum glaucescens</i> (misapplied in Tasmania) | |
| | <i>Leptospermum flavescens</i> Sm. var. <i>commune</i> Benth. = <i>Leptospermum glaucescens</i> | |
| | <i>Leptospermum flavescens</i> Sm. var. <i>grandiflorum</i> (Lodd., G.Lodd & W.Lodd.) Benth. = <i>Leptospermum grandiflorum</i> | |
| | <i>Leptospermum flavescens</i> Sm. var. <i>nitidum</i> (Hook.f.) Rodway = <i>Leptospermum nitidum</i> | |
| e | <i>Leptospermum glaucescens</i> S.Schauer, Linnaea 15: 421 (1841) | 1:201 |
| e | <i>Leptospermum grandiflorum</i> Lodd., G.Lodd & W.Lodd., G.Lodd. & W.Lodd., Bot. Cab. 6: t.514 (1821) | 1:200 |
| | <i>Leptospermum humifusum</i> W.M.Curtis = <i>Leptospermum rupestre</i> nom. inval. | |
| | <i>Leptospermum laevigatum</i> (Gaertn.) F.Muell., Cat. Pl. Melb. Bot. Gard. 22 (1858) | 1:200 |
| | <i>Leptospermum lanigerum</i> (Sol. ex Aiton) Sm., Trans. Linn. Soc. London 3: 263 (1797) | 1:200 |
| | <i>Leptospermum lanigerum</i> (Sol. ex Aiton) Sm. var. <i>montanum</i> Rodway = <i>Leptospermum lanigerum</i> | |
| | <i>Leptospermum myrtifolium</i> Sieber ex DC. sensu Hooker (1960), Rodway (1903) = <i>Leptospermum glaucescens</i> (misapplied in Tasmania) | |
| e | <i>Leptospermum nitidum</i> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 139 (1856) | 1:201 |
| | <i>Leptospermum pilosum</i> Schauer a name of uncertain application | |
| e | <i>Leptospermum riparium</i> D.I.Morris, Records of the Queen Victoria Museum 50: 2 (1974) | 1:201 |
| | <i>Leptospermum rodwayanum</i> Summerh. & H.F.Comber = <i>Leptospermum grandiflorum</i> | |
| e | <i>Leptospermum rupestre</i> Hook.f., Icon. Pl. 4: t.308 (1840) | 1:202 |
| | <i>Leptospermum scoparium</i> J.R.Forst. & G.Forst., Char. Gen. Pl.: 72, t.36, figs.f-l (1775) | 1:200 |
| | <i>Leptospermum scoparium</i> J.R.Forst. & G.Forst. var. <i>eximium</i> B.L.Burtt = <i>Leptospermum scoparium</i> | |

| | | |
|---|--|------------------|
| | Leptospermum sericeum Labill. sensu Curtis (1956) = Leptospermum glaucescens (misapplied in Tasmania) | |
| | Melaleuca armillaris (Sol. ex Gaertn.) Sm. subsp. armillaris , Trans. Linn. Soc. London 3: 277 (1797) | 1:205 |
| | Melaleuca ericifolia Sm., Trans. Linn. Soc. London 3: 276 (1797) | 1:204 |
| | Melaleuca gibbosa Labill., Nov. Holl. Pl. 2: 30, t.172 (1806) | 1:204 |
| | Melaleuca pallida (Bonpl.) Craven, Novon 16: 472 (2006) | 1:203 |
| e | Melaleuca pustulata Hook.f., London J. Bot. 6: 476 bis (1847) | 1:205 |
| | Melaleuca squamea Labill., Nov. Holl. Pl. 2: 28, t.168 (1806) | 1:204 |
| | Melaleuca squarrosa Donn ex Sm., Trans. Linn. Soc. London 6: 300 (1802) | 1:204 |
| e | Melaleuca virens Craven, Novon 16: 473 (2006) | 1:203 |
| i | Syzygium smithii (Poir.) Nied., Nat. Pflanzenfam. [Engler & Prantl] III. 7: 85 (1893) | |
| | Thryptomene micrantha Hook.f., Hooker's J. Bot. Kew Gard. Misc. 5: 299, t.8 (1853) | 1:198 |
| | NOTHOFAGACEAE | FTO 66 |
| | Fagus cunninghamii Hook. = Nothofagus cunninghamii | |
| | Fagus gunnii Hook.f. = Nothofagus gunnii | |
| | Fuscospora gunnii (Hook.f.) Heenan & Smissen = Nothofagus gunnii | |
| | Lophozonia cunninghamii (Hook.) Heenan & Smissen = Nothofagus cunninghamii | |
| | Nothofagus cunninghamii (Hook.) Oerst., Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd. (ser. 5), 9: 355 (1871) | 3:647 |
| e | Nothofagus gunnii (Hook.f.) Oerst., Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd. (ser. 5), 9: 354 (1871) | 3:646 |
| | NYCTAGINACEAE | FTO 101.1 |
| i | Mirabilis jalapa L., Sp. Pl. 1: 177 (1753) | |
| | OLEACEAE | FTO 113 |
| i | Fraxinus angustifolia Vahl subsp. angustifolia , Enum. Pl. [Vahl] 1: 52 (1804) | |
| i | Ligustrum vulgare L., Sp. Pl. 1: 7 (1753) | 3:470 |
| | Notelaea ligustrina Vent., Choix Pl. 25 (1804) | 3:470 |
| | ONAGRACEAE | |
| | Boisduvalia tasmanica (Hook.f.) Munz = Epilobium curtisiae | 2:234 |
| | Epilobium adenocaulon Hausskn. = Epilobium ciliatum | 2:233 |
| | Epilobium alpinum L. nom. illeg., sensu Rodway (1903) = Epilobium tasmanicum (misapplied in Tasmania) | |
| | Epilobium billardiereanum Ser. ex DC. subsp. billardiereanum , Prodr. [A. P. de Candolle] 3: 41 (1828) | 2:232 |
| | Epilobium billardiereanum Ser. ex DC. subsp. cinereum (A.Rich.) P.H.Raven & Engelhorn, New Zealand J. Bot. 9: 349 (1971) | 2:230 |
| | Epilobium billardiereanum Ser. ex DC. subsp. hydrophilum P.H.Raven & Engelhorn Sensu Buchanan et al (1989) = Epilobium billardiereanum subsp. billardiereanum (misapplied in Tasmania) | |
| | Epilobium billardiereanum Ser. ex DC. subsp. intermedium P.H.Raven & Engelhorn, New Zealand J. Bot. 9: 348 (1971) | |
| i | Epilobium ciliatum Raf., Med. Repos. 5: 361 (1808) | 2:233 |
| | Epilobium confertifolium Hook.f. sensu Bentham (1867) = Epilobium tasmanicum (misapplied in Tasmania) | |
| | Epilobium curtisiae P.H.Raven, Aliso 5: 249, fig.1 (1963) | 2:234 |

- Epilobium erosum Hausskn. = **Epilobium gunnianum** 2:230
- e **Epilobium fugitivum** P.H.Raven & Engelhorn, New Zealand J. Bot. 9: 346 (1971)
- Epilobium glabellum G.Forst. sensu Rodway (1903) = **Epilobium billardioreanum** (misapplied in Tasmania)
- Epilobium gunnianum** Hausskn., Oesterr. Bot. Z. 29: 149 (1879) 2:230
- Epilobium hirtigerum** A.Cunn., Ann. Nat. Hist. 3: 33 (1839) 2:231
- Epilobium junceum G.Forst. ex Spreng. sensu Curtis (1963) = **Epilobium billardioreanum** subsp. **cinereum** (misapplied in Tasmania) 2:230
- i * t Epilobium nummulariifolium R.Cunn. ex A.Cunn., Ann. Nat. Hist. 3: 31 (1839)
- i t **Epilobium obscurum** Schreb., Spic. Fl. Lips. 147, 155 (1771)
- Epilobium pallidiflorum** A.Cunn., Ann. Nat. Hist. 3: 34 (1839) 2:231
- e **Epilobium perpusillum** Hausskn., Monogr. Epilobium: 301, t.21, fig. 90 (1884) 2:233
- i t **Epilobium rotundifolium** G.Forst., Fl. Ins. Austr. 27 (1786)
- Epilobium sarmentaceum** Hausskn., Oesterr. Bot. Z. 29: 149 (1879) 2:229
- Epilobium tasmanicum** Hausskn., Monogr. Epilobium: 296, t.20, fig.84 (1884) 2:233
- Epilobium tenuipes Hook.f. sensu Hooker (1860) = **Epilobium tasmanicum** (misapplied in Tasmania)
- Epilobium tetragonum L. sensu Bentham (1867) = **Epilobium billardioreanum** (misapplied in Tasmania)
- Epilobium willisii** P.H.Raven & Engelhorn, New Zealand J. Bot. 9: 347 (1971)
- i **Fuchsia magellanica** Lam., Encycl. (Lamarck) 2: 565 (1788) 2:235
- i # **Oenothera biennis** L., Sp. Pl. 1: 346 (1753)
- i **Oenothera glazioviana** Micheli, Fl. Bras. (Martius) 13: 178 (1875)
- i **Oenothera stricta** Ledeb. ex Link subsp. **stricta**, Enum. Hort. Berol. Alt. 1: 377 (1821) 2:235
- Oenothera tasmanica Hook.f. = **Epilobium curtisiae**

OROBANCHACEAE

- i **Bartsia trixago** L., Sp. Pl. 2: 602 (1753)
- Bellardia trixago (L.) All. = **Bartsia trixago**
- Euphrasia alpina R.Br. nom. illeg. a name of uncertain application
- Euphrasia alpina R.Br. var. angustifolia Benth. sensu Hooker (1860) = **Euphrasia collina** subsp. **collina**
- Euphrasia alpina R.Br. var. humilis Benth. sensu Hooker (1860) = **Euphrasia striata**
- e **Euphrasia amphisysepala** W.R.Barker, J. Adelaide Bot. Gard. 10: 204 (1987)
- e **Euphrasia amplidens** W.R.Barker, J. Adelaide Bot. Gard. 26: 23 (2013)
- e **Euphrasia sp. Bivouac Bay (W.R.Barker 7626 et al.) W.R.Barker**
- Euphrasia brownii F.Muell. nom. illeg., sensu Rodway (1903) a catch-all name that includes several **Euphrasia** species
- Euphrasia brownii F.Muell. var. striata (R.Br.) Rodway = **Euphrasia striata** 3:530
- Euphrasia collina** R.Br. subsp. **collina**, Prodr. Fl. Nov. Holland. 436 (1810) 3:527
- e **Euphrasia collina** R.Br. subsp. **deflexifolia** (Gand.) W.R.Barker, The Endemic Flora of Tasmania 6: 477 (1978)
- e **Euphrasia collina** R.Br. subsp. **diemenica** (Spreng.) W.R.Barker, The Endemic Flora of Tasmania 6: 477 (1978) 3:528
- e **Euphrasia collina** R.Br. subsp. **Dukes Marshes (A.Moscals 861) Tas Herbarium**
- e **Euphrasia collina** R.Br. subsp. **gunnii** (Du Rietz) W.R.Barker, J. Adelaide Bot. Gard. 5: 199 (1982) 3:528
- e **Euphrasia collina** R.Br. subsp. **Northwest Tasmania (M.Visoiu 216) Tas Herbarium**

| | | |
|---|---|---------------|
| e | Euphrasia collina R.Br. subsp. tasmanica (Gand.) W.R.Barker, J. Adelaide Bot. Gard. 21: 93 (2007) | |
| | Euphrasia collina R.Br. subsp. tetragona (R.Br.) W.R.Barker, J. Adelaide Bot. Gard. 5: 189 (1982) | |
| | Euphrasia cuspidata Hook.f. = Euphrasia hookeri | 3:532 |
| | Euphrasia diemenica Spreng. = Euphrasia collina subsp. diemenica | 3:528 |
| | Euphrasia sp. fabula W.C.Potts & W.R.Barker MS = Euphrasia sp. Bivouac Bay (W.R.Barker 7626 et al.) W.R.Barker | |
| e | Euphrasia fragosa W.R.Barker, J. Adelaide Bot. Gard. 17: 217 (1996) | |
| | Euphrasia gibbsiae Du Rietz subsp. aff. subglabrifolia (Du Rietz) W.R.Barker = Euphrasia gibbsiae subsp. Mt Field (W.R.Barker 1187) Tas Herbarium | |
| e | Euphrasia gibbsiae Du Rietz subsp. comberi (Du Rietz) W.R.Barker, J. Adelaide Bot. Gard. 5: 122 (1982) | 3:531 |
| e | Euphrasia gibbsiae Du Rietz subsp. discolor W.R.Barker, J. Adelaide Bot. Gard. 5: 128 (1982) | |
| e | Euphrasia gibbsiae Du Rietz subsp. gibbsiae , Svensk Bot. Tidskr. 42: 104 (1948) | 3:531 |
| e | Euphrasia gibbsiae Du Rietz subsp. kingii (W.M.Curtis) W.R.Barker, The Endemic Flora of Tasmania 6: 477 (1978) | 3:530 |
| e | Euphrasia gibbsiae Du Rietz subsp. microdonta W.R.Barker, J. Adelaide Bot. Gard. 5: 126 (1982) | |
| e | Euphrasia gibbsiae Du Rietz subsp. Mt Anne (A.M.Buchanan 5109) Tas Herbarium | |
| e | Euphrasia gibbsiae Du Rietz subsp. Mt Field (W.R.Barker 1187) Tas Herbarium | |
| e | Euphrasia gibbsiae Du Rietz subsp. psilantherea (F.Muell.) W.R.Barker, J. Adelaide Bot. Gard. 5: 119 (1982) | |
| e | Euphrasia gibbsiae Du Rietz subsp. pulvinestris W.R.Barker, J. Adelaide Bot. Gard. 5: 131 (1982) | |
| e | Euphrasia gibbsiae Du Rietz subsp. wellingtonensis W.R.Barker, J. Adelaide Bot. Gard. 5: 120 (1982) | |
| | Euphrasia gunnii Du Rietz = Euphrasia collina subsp. gunnii | 3:528 |
| e | Euphrasia hookeri Wettst., Monogr. Euphrasia: 268 (1896) | 3:532 |
| | Euphrasia kingii W.M.Curtis = Euphrasia gibbsiae subsp. kingii | 3:530 |
| | Euphrasia multicaulis Benth. sensu Hooker (1860) = Euphrasia collina subsp. tetragona | |
| e | Euphrasia phragmostoma W.R.Barker, J. Adelaide Bot. Gard. 5: 103 (1982) | |
| | Euphrasia scabra R.Br., Prodr. Fl. Nov. Holland. 437 (1810) | 3:531 |
| e | Euphrasia semipicta W.R.Barker, J. Adelaide Bot. Gard. 5: 139 (1982) | |
| e | Euphrasia striata R.Br., Prodr. Fl. Nov. Holland. 436 (1810) | 3:530 |
| | Orobanche australiana F.Muell. sensu Buchanan (1995) previously recorded in error | |
| i | Orobanche minor Sm., Engl. Bot. 6: t.422 (1797) | 3:533 |
| i | Parentucellia latifolia (L.) Caruel, Fl. Ital. (Parlatore) 6: 480 (1885) | 3:532 |
| i | Parentucellia viscosa (L.) Caruel, Fl. Ital. (Parlatore) 6: 482 (1885) | 3:533 |
| | OXALIDACEAE | FTO 70 |
| i | Oxalis articulata Savigny, Encycl. (Lamarck) 4: 686 (1798) | 1:102 |
| i | Oxalis corniculata L. subsp. corniculata , Sp. Pl. 1: 435 (1753) | 1:101 |
| | Oxalis exilis A.Cunn., Ann. Nat. Hist. 3: 316 (1839) | |
| i | Oxalis incarnata L., Sp. Pl. 1: 433 (1753) | 1:102 |
| | Oxalis lactea Hook. = Oxalis magellanica | 1:101 |
| i | Oxalis latifolia Kunth, Nov. Gen. Sp. [H.B.K.] 5: 237, t.467 (1823) | 1:101 |
| | Oxalis magellanica G.Forst., Commentat. Soc. Regiae Sci. Gott. 9: 33 (1789) | 1:101 |
| | Oxalis perennans Haw., Misc. Nat. 181 (1803) | |

| | | |
|------------------------|--|----------------|
| i | Oxalis pes-caprae L., Sp. Pl. 1: 434 (1753) | 1:101 |
| i | Oxalis purpurea L., Sp. Pl. 1: 433 (1753) | 1:102 |
| | Oxalis radicata A.Rich., Tent. Fl. Abyss. 2: 123 (1847) | |
| | Oxalis rubens Haw., Misc. Nat. 182 (1803) | |
| PAPAVERACEAE | | FTO 44 |
| | <i>Argemone mexicana</i> L. sensu Curtis (1956), Curtis & Morris (1975) = Argemone ochroleuca subsp. ochroleuca (misapplied in Tasmania) | 1:28 |
| i | Argemone ochroleuca Sweet subsp. ochroleuca , Brit. Fl. Gard. [Sweet] 3: t.242 (1828) | 1:28 |
| i | Eschscholzia californica Cham., Horae Phys. Berol. [Nees]: 73-74, t.15 (1820) | 1:29 |
| i | Glaucium flavum Crantz, Stirp. Austr. Fasc. Sp. 2: 133 (1763) | 1:28 |
| i | Papaver aculeatum Thunb., Prodr. Pl. Cap. 92 (1800) | 1:27 |
| i | Papaver argemone L., Sp. Pl. 1: 506 (1753) | 1:27 |
| i | Papaver dubium L., Sp. Pl. 2: 1196 (1753) | 1:27 |
| i | Papaver hybridum L., Sp. Pl. 1: 506 (1753) | 1:27 |
| i | Papaver rhoeas L., Sp. Pl. 1: 507 (1753) | 1:27 |
| i | Papaver somniferum L. subsp. setigerum (DC.) Arcang., Nouv. Fl. Normandie: 30 (1894) | 1:28 |
| i | Papaver somniferum L. subsp. somniferum , Sp. Pl. 1: 508 (1753) | 1:28 |
| PASSIFLORACEAE | | FTO 81 |
| ?i # | Passiflora cinnabarina Lindl., Gard. Chron. 23: 724 (1855) | |
| | <i>Passiflora mollissima</i> (Kunth) L.H.Bailey sensu Buchanan (1999) = Passiflora tarminiana (misapplied in Tasmania) | |
| i | Passiflora tarminiana Coppens & V.E.Barney, Novon 11: 9 (2001) | |
| PHRYMACEAE | | FTO 117 |
| i | Erythranthe moschata (Douglas ex Lindl.) G.L.Nesom, Phytoneuron 39: 38 (2012) | |
| | Glossostigma elatinoides (Benth.) Benth. ex Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 189 (1853) | 3:518 |
| | <i>Mimulus moschatus</i> Douglas ex Lindl. = Erythranthe moschata | 3:516 |
| | <i>Mimulus repens</i> R.Br. = Thyridia repens | 3:515 |
| | Thyridia repens (R.Br.) W.R.Barker & Beardsley, Phytoneuron 39: 20 (2012) | |
| PHYLLANTHACEAE | | |
| | <i>Oreoporphantha petalifera</i> Orchard & J.B.Davies = Poranthera petalifera | |
| | Phyllanthus australis Hook.f., London J. Bot. 6: 284 (1847) | 3:638 |
| | Phyllanthus gunnii Hook.f., London J. Bot. 6: 284 (1847) | 3:638 |
| | Poranthera microphylla Brongn., Ann. Sci. Nat. (Paris) 29: 385 (1833) | 3:634 |
| e | Poranthera petalifera (Orchard & J.B.Davies) Halford & R.J.F.Hend., Austrobaileya 7: 22 (2005) | |
| PHYTOLACCACEAE | | |
| i | Phytolacca octandra L., Sp. Pl., ed. 2: 631 (1762) | |
| PICRODENDRACEAE | | |
| | Micranthemum hexandrum Hook.f., London J. Bot. 6: 283 (1847) | 3:634 |
| e | Micranthemum serpentinum Orchard, Aspects of Tasmanian Botany: 60 (1991) | |
| | <i>Pseudanthus divaricatissimus</i> (Müll.Arg.) Benth. sensu Buchanan (1999) recorded in error | |
| | Pseudanthus ovalifolius F.Muell., Trans. & Proc. Philos. Inst. Victoria 2: 66 (1857) | 3:634 |
| | <i>Pseudanthus tasmanicus</i> Rodway = Muehlenbeckia axillaris (Polygonaceae) | 3:593 |

PITTOSPORACEAE

- Billardiera alpina (McGill.) E.M.Benn. sensu Buchanan (1989) = **Rhytidosporum inconspicuum**
(misapplied in Tasmania)
- Billardiera fusiformis Labill. sensu Buchanan (2009) = **Billardiera heterophylla** (misapplied in Tasmania) 1:59
- i **Billardiera heterophylla** (Lindl.) L.W.Cayzer & Crisp, Austral. Syst. Bot. 17: 119 (2004) 1:59
- e **Billardiera longiflora** Labill., Nov. Holl. Pl. 1: 64, t.89 (1805) 1:58
- Billardiera longiflora Labill. var. alpina Rodway = **Billardiera longiflora**
- Billardiera longiflora Labill. var. ovalis (Lindl.) E.M.Benn. = **Billardiera ovalis**
- Billardiera macrantha** Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 37 (1855)
- Billardiera mutabilis** Salisb., Parad. Lond. 1: t.48 (1806) 1:59
- Billardiera nesophila L.W.Cayzer & D.L.Jones = **Billardiera macrantha**
- e **Billardiera ovalis** Lindl., Edwards's Bot. Reg. 20: t.1719 (1834)
- Billardiera procumbens (Hook.) E.M.Benn. = **Rhytidosporum procumbens** 1:58
- Billardiera scandens Sm. sensu Curtis & Morris (1975) = **Billardiera mutabilis** (misapplied in Tasmania) 1:59
- Billardiera viridiflora L.W.Cayzer & D.L.Jones = **Billardiera macrantha**
- Bursaria spinosa** Cav. subsp. **spinosa**, Icon. [Cavanilles] 4: 30, t.350 (1797) 1:57
- Marianthus procumbens (Hook.) Benth. = **Rhytidosporum procumbens** 1:58
- Pittosporum bicolor** Hook., J. Bot. (Hooker) 1: 249 (1834) 1:56
- i **Pittosporum crassifolium** Banks & Sol. ex A.Cunn., Ann. Nat. Hist. 4: 106 (1840)
- i **Pittosporum eugenioides** A.Cunn., Ann. Nat. Hist. 4: 106 (1840)
- i **Pittosporum tenuifolium** Gaertn., Fruct. Sem. Pl. 1: 286 (1788)
- i **Pittosporum undulatum** Vent., Descr. Pl. Nouv. 76, t.76 (1802)
- i **Pittosporum undulatum** Vent. subsp. × **emmettii** W.M.Curtis, Records of the Queen Victoria Museum 50: 3(1974) = **Pittosporum bicolor** × **P. undulatum** 1:57
- Rhytidosporum alpinum McGill. sensu Buchanan (1995) = **Rhytidosporum inconspicuum** (misapplied in Tasmania)
- Rhytidosporum inconspicuum** L.W.Cayzer, Crisp & I.Telford, Austral. Syst. Bot. 12: 700 (1999)
- Rhytidosporum procumbens** (Hook.) F.Muell., Pl. Victoria 1: 75 (1862) 1:58
- Sollya heterophylla Lindl. = **Billardiera heterophylla** 1:59

PLANTAGINACEAE

- i # **Antirrhinum majus** L., Sp. Pl. 2: 617 (1753)
- Callitriche brachycarpa** Hegelm., Verh. Bot. Vereins Prov. Brandenburg 10: 115 (1868) 1:195
- i * **Callitriche brutia** Petagna subsp. **brutia**, Institutiones Botanicae 2: 10 (1787)
- Callitriche sonderi** Hegelm., Verh. Bot. Vereins Prov. Brandenburg 9: 18 (1867)
- i **Callitriche stagnalis** Scop., Fl. Carniol., ed. 2, 2: 251 (1772) 1:195
- Callitriche umbonata** Hegelm., Verh. Bot. Vereins Prov. Brandenburg 9: 19 (1867) 1:196
- Callitriche verna** L. sensu Hooker (1860), Bentham (1864) = **Callitriche stagnalis** (misapplied in Tasmania)
- Chionohebe ciliolata (Hook.f.) B.G.Briggs & Ehrend. = **Veronica ciliolata**
- i **Cymbalaria muralis** G.Gaertn., B.Mey. & Scherb., Oekon. Fl. Wetterau 2: 397 (1800) 3:514
- Derwentia derwentiana (Andrews) B.G.Briggs & Ehrend. subsp. **derwentiana** = **Veronica derwentiana** subsp. **derwentiana** 3:523

| | | |
|-----|---|-------|
| | Derwentia nivea (Lindl.) B.G.Briggs & Ehrend. = Veronica nivea | 3:523 |
| i | Digitalis purpurea L., Sp. Pl. 2: 621 (1753) | 3:519 |
| | Gratiola latifolia R.Br. = Gratiola peruviana | 3:517 |
| | Gratiola nana Benth., Prodr. [A. P. de Candolle] 10: 404 (1846) | 3:517 |
| | Gratiola peruviana L., Sp. Pl. 1: 17 (1753) | 3:517 |
| | Gratiola pubescens R.Br., Prodr. Fl. Nov. Holland. 435 (1810) | 3:517 |
| | Gratiola pumilo F.Muell., Linnaea 25: 431 (1853) | |
| | Hebe elliptica (G.Forst.) Pennell = Veronica elliptica | |
| i | Kickxia elatine (L.) Dumort. subsp. elatine , Fl. Belg. (Dumortier): 35 (1827) | 3:514 |
| i * | Kickxia spuria (L.) Dumort. subsp. integrifolia (Brot.) R.Fern., Bot. J. Linn. Soc. 64: 74 (1971) | 3:514 |
| | Limosella aquatica L. sensu Rodway (1903) = Limosella australis | 3:519 |
| | Limosella australis R.Br., Prodr. Fl. Nov. Holland. 443 (1810) | 3:519 |
| | Limosella lineata Glück = Limosella australis | 3:519 |
| | Linaria spuria (L.) Mill. = Kickxia spuria | |
| i | Linaria vulgaris Mill., Gard. Dict., ed. 8, LIN I (1768) | 3:513 |
| e | Ourisia integrifolia R.Br., Prodr. Fl. Nov. Holland. 439 (1810) | 3:520 |
| | Parahebe derwentiana (Andrews) B.G.Briggs & Ehrend. = Veronica derwentiana | 3:523 |
| | Plantago antarctica Decne., Prodr. [A. P. de Candolle] 13: 703 (1852) | 3:562 |
| | Plantago archeri Hook.f. = Plantago tasmanica var. archeri | |
| i | Plantago australis Lam., Tabl. Encycl. 1: 339 (1792) | |
| e | Plantago bellidioides Decne., Prodr. [A. P. de Candolle] 13: 701 (1852) | 3:561 |
| | Plantago brownii Rapin = Plantago triantha | |
| | Plantago carnososa R.Br. = Plantago triantha | |
| i | Plantago coronopus L. subsp. commutata (Guss.) Pilg., Repert. Spec. Nov. Regni Veg. 28: 287 (1930) | 3:560 |
| i | Plantago coronopus L. subsp. coronopus , Sp. Pl. 1: 115 (1753) | 3:560 |
| e | Plantago daltonii Decne., Prodr. [A. P. de Candolle] 13: 703 (1852) | 3:563 |
| | Plantago debilis R.Br., Prodr. Fl. Nov. Holland. 425 (1810) | 3:560 |
| | Plantago gaudichaudii Barnéoud, Monogr. Plantag. 15 (1845) | |
| e | Plantago glabrata Hook.f., London J. Bot. 6: 276 (1847) | 3:562 |
| | Plantago glacialis B.G.Briggs, Carolin & Pulley, Contr. New South Wales Natl. Herb. 4: 395 (1973) | 3:564 |
| e | Plantago gunnii Hook.f., London J. Bot. 5: 446 (1846) | 3:564 |
| | Plantago hispida R.Br., Prodr. Fl. Nov. Holland. 425 (1810) | 3:561 |
| i | Plantago lanceolata L., Sp. Pl. 1: 113 (1753) | 3:560 |
| i | Plantago major L., Sp. Pl. 1: 112 (1753) | 3:559 |
| | Plantago muelleri Pilg. sensu Curtis (1967) = Plantago glacialis (misapplied in Tasmania) | 3:564 |
| e | Plantago paradoxa Hook.f., London J. Bot. 6: 277 (1847) | 3:564 |
| e | Plantago tasmanica Hook.f. var. archeri (Hook.f.) W.M.Curtis, The Student's Flora of Tasmania 3: 650 (1967) | 3:563 |
| | Plantago tasmanica Hook.f. var. daltonii (Decne.) Hook.f. = Plantago daltonii | |
| | Plantago tasmanica Hook.f. var. glabrata (Hook.f.) Hook.f. = Plantago glabrata | |
| e | Plantago tasmanica Hook.f. var. tasmanica , London J. Bot. 6: 276 (1847) | 3:562 |
| t | Plantago triantha Spreng., Syst. Veg. (ed. 16) [Sprengel] 1: 439 (1824) | 3:563 |

| | | |
|-----|---|----------------|
| | Plantago varia R.Br., Prodr. Fl. Nov. Holland. 425 (1810) | 3:561 |
| | Pygmea ciliolata Hook.f. = Veronica ciliolata | |
| | Veronica arguta R.Br. sensu Hooker (1860) a name of uncertain application | |
| i | Veronica arvensis L., Sp. Pl. 1: 13 (1753) | 3:525 |
| | Veronica calycina R.Br., Prodr. Fl. Nov. Holland. 435 (1810) | 3:524 |
| t | Veronica ciliolata (Hook.f.) Cheeseman subsp. fiordensis (Ashwin) Meudt, Austral. Syst. Bot. 21: 413 (2008) | |
| e | Veronica continua B.G.Briggs, Telopea 11: 278 (2006) | |
| | Veronica derwentiana Andrews subsp. derwentiana , Bot. Repos 8: t. 531 (1808) | 3:523 |
| | Veronica distans R.Br. var. pubescens Benth. = Veronica novae-hollandiae | 3:523 |
| i t | Veronica elliptica G.Forst., Fl. Ins. Austr. 3 (1786) | |
| e | Veronica formosa R.Br., Prodr. Fl. Nov. Holland. 434 (1810) | 3:522 |
| | Veronica gracilis R.Br., Prodr. Fl. Nov. Holland. 435 (1810) | 3:523 |
| i | Veronica hederifolia L., Sp. Pl. 1: 13 (1753) | 3:525 |
| | Veronica labiata R.Br. nom. illeg. = Veronica derwentiana | |
| | Veronica nivea Lindl., Edwards's Bot. Reg. 28: 42 (1842) | 3:523 |
| | Veronica notabilis F.Muell. ex Benth., Fl. Austral. 4: 511 (1868) | 3:524 |
| e | Veronica novae-hollandiae Poir., Encycl. (Lamarck) 8: 526 (1808) | 3:523 |
| | Veronica peregrina L. previously listed as naturalised but insufficient evidence exists to support this | 3:525 |
| i | Veronica persica Poir., Encycl. (Lamarck) 8: 542 (1808) | 3:526 |
| | Veronica plebeia R.Br., Prodr. Fl. Nov. Holland. 435 (1810) | 3:524 |
| i t | Veronica scutellata L., Sp. Pl. 1: 12 (1753) | |
| i | Veronica serpyllifolia L., Sp. Pl. 1: 12 (1753) | 3:525 |
| | PLUMBAGINACEAE | FTO 94 |
| | Limonium australe (R.Br.) Kuntze var. australe , Revis. Gen. Pl. 2: 395 (1891) | 3:465 |
| e | Limonium australe (R.Br.) Kuntze var. baudinii (Lincz.) A.M.Gray, Kanunna 4: 117 (2010) | |
| | Limonium baudinii Lincz. = Limonium australe var. baudinii | |
| i # | Limonium sinuatum (L.) Mill., Gard. Dict., ed. 8, 6 (1768) | |
| | Statice australis (R.Br.) Spreng. = Limonium australe | |
| | Taxantheme australis R.Br. = Limonium australe | |
| | POLEMONIACEAE | FTO 102 |
| | Collomia grandiflora Douglas ex Lindl. previously listed as naturalised but insufficient evidence exists to support this | 3:485 |
| | Collomia linearis (Cav.) Nutt. sensu Curtis (1967) = Collomia grandiflora (misapplied in Tasmania) | 3:485 |
| i | Navarretia squarrosa (Eschsch.) Hook. & Arn., Bot. Beechey Voy. 368 (1839) | 3:485 |
| | POLYGALACEAE | FTO 60 |
| | Comesperma calymega Labill., Nov. Holl. Pl. 2: 23, t.162 (1806) | 1:62 |
| | Comesperma defoliatum F.Muell., Pl. Victoria 1: 189 (1862) | 1:63 |
| | Comesperma ericinum DC., Prodr. [A. P. de Candolle] 1: 334 (1824) | 1:62 |
| | Comesperma retusum Labill., Nov. Holl. Pl. 2: 22, t.160 (1806) | 1:62 |
| | Comesperma volubile Labill., Nov. Holl. Pl. 2: 24, t.163 (1806) | 1:62 |
| i | Polygala myrtifolia L., Sp. Pl. 2: 703 (1753) | 1:63 |
| i | Polygala vulgaris L., Sp. Pl. 2: 702 (1753) | |

POLYGONACEAE**FTO 95**

- i **Acetosa sagittata** (Thunb.) L.A.S.Johnson & B.G.Briggs, Contr. New South Wales Natl. Herb. 3: 166 (1962)
- i **Acetosella vulgaris** Fourr., Ann. Soc. Linn. Lyon, Ser. 2, 17: 145 (1869) 3:587
- i **Emex australis** Steinh., Ann. Sci. Nat., Bot., sér. 2, 9: 195 (1838) 3:583
- i **Fallopia convolvulus** (L.) Á.Löve, Taxon 19: 300 (1970) 3:591
- i **Fallopia japonica** (Houtt.) Ronse Decr., Bot. J. Linn. Soc. 98: 369 (1988)
- Muehlenbeckia adpressa** (Labill.) Meisn., Pl. Vasc. Gen. [Meisner] 2: 227 (1843) 3:592
- Muehlenbeckia adpressa (Labill.) Meisn. var. hastifolia Meisn. = **Muehlenbeckia gunnii**
- Muehlenbeckia adpressa (Labill.) Meisn. var. rotundifolia Benth. nom. illeg. = **Muehlenbeckia adpressa**
- Muehlenbeckia australis (G.Forst.) Meisn. sensu Buchanan (2009) = **Muehlenbeckia adpressa** (misapplied in Tasmania)
- Muehlenbeckia axillaris** (Hook.f.) Endl., Gen. Pl. [Endlicher] Suppl. 4(2): 51 (1848) 3:593
- Muehlenbeckia gunnii** (Hook.f.) Endl., Gen. Pl. [Endlicher] Suppl. 4(2): 51 (1848) 3:593
- i **Persicaria capitata** (Buch.-Ham. ex D.Don) H.Gross, Bot. Jahrb. Syst. 49: 277 (1913)
- Persicaria decipiens** (R.Br.) K.L.Wilson, Telopea 3: 178 (1988) 3:590
- Persicaria hydropiper** (L.) Delarbre, Fl. Auvergne (Delarbre) 2: 518 (1800) 3:590
- i **Persicaria lapathifolia** (L.) Gray, Nat. Arr. Brit. Pl. 2: 270 (1821) 3:591
- i **Persicaria maculosa** Gray, Nat. Arr. Brit. Pl. 2: 269 (1821) 3:591
- Persicaria praetermissa** (Hook.f.) H.Hara, Fl. E. Himalaya: 73 (1966) 3:589
- ?i **Persicaria prostrata** (R.Br.) Soják, Preslia 46: 154 (1974) 3:590
- Persicaria subsessilis** (R.Br.) K.L.Wilson, Telopea 3: 180 (1988) 3:591
- Polygonum adpressum Labill. = **Muehlenbeckia adpressa**
- i **Polygonum arenastrum** Jord. ex Boreau, Fl. Centre France, ed. 3 [Boreau], 2: 559 (1857) 3:589
- i **Polygonum aviculare** L., Sp. Pl. 1: 362 (1753) 3:588
- Polygonum convolvulus L. = **Fallopia convolvulus** 3:591
- Polygonum decipiens R.Br. = **Persicaria decipiens** 3:590
- Polygonum hydropiper L. = **Persicaria hydropiper** 3:590
- Polygonum lapathifolium L. = **Persicaria lapathifolia** 3:591
- Polygonum minus Huds. sensu Bentham (1870) = **Persicaria decipiens** (misapplied in Tasmania)
- Polygonum persicaria L. = **Persicaria maculosa** 3:591
- Polygonum plebeium R.Br. sensu Curtis (1967) = **Polygonum arenastrum** (misapplied in Tasmania) 3:589
- Polygonum prostratum R.Br. = **Persicaria prostrata** 3:590
- Polygonum strigosum R.Br. sensu Curtis (1967) = **Persicaria praetermissa** (misapplied in Tasmania) 3:589
- Polygonum subsessile R.Br. = **Persicaria subsessilis** 3:591
- Reynoutria japonica Houtt. = **Fallopia japonica**
- Rumex acetosella L. = **Acetosella vulgaris** 3:587
- Rumex angiocarpus Murb. sensu Curtis (1967) = **Acetosella vulgaris** (misapplied in Tasmania) 3:586
- Rumex bidens** R.Br., Prodr. Fl. Nov. Holland. 421 (1810) 3:586
- Rumex brownii** Campd., Monogr. Rumex 64: 81 (1819) 3:585
- i **Rumex conglomeratus** Murray, Prodr. Stirp. Gott. 52 (1770) 3:585
- i **Rumex crispus** L., Sp. Pl. 1: 335 (1753) 3:584
- Rumex dumosiformis Rech. = **Rumex dumosus** 3:586

| | | |
|----------------------|---|----------------|
| | Rumex dumosus A.Cunn. ex Meisn., Prodr. [A. P. de Candolle] 14: 62 (1856) | 3:586 |
| | Rumex dumosus A.Cunn. ex Meisn. var. dumosiformis (Rech.f.) Rech.f. = Rumex dumosus | 3:586 |
| i | Rumex obtusifolius L., Sp. Pl. 1: 335 (1753) | 3:585 |
| | Rumex obtusifolius L. subsp. transiens (Simonk.) Rech.f. sensu Buchanan (1999) = Rumex obtusifolius (misapplied in Tasmania) | |
| i | Rumex pulcher L. subsp. pulcher , Sp. Pl. 1: 336 (1753) | 3:585 |
| | Rumex sagittatus Thunb. = Acetosa sagittata | |
| | Rumex sanguineus L. sensu Rodway (1903) = Rumex conglomeratus (misapplied in Tasmania) | |
| PORTULACACEAE | | FTO 100 |
| | Calandrinia calyptrata Hook.f., Icon. Pl. 3: t.296 (1840) | 1:78 |
| | Calandrinia caulescens Kunth = Calandrinia ciliata | 1:79 |
| i | Calandrinia ciliata (Ruiz & Pav.) DC., Prodr. [A. P. de Candolle] 3: 359 (1828) | 1:79 |
| | Calandrinia eremaea Ewart, Fl. Victoria: 486 (1931) | |
| | Calandrinia granulifera Benth., Fl. Austral. 1: 176 (1863) | 1:78 |
| | Calandrinia menziesii (Hook.) Torr. & A.Gray = Calandrinia ciliata | 1:79 |
| | Calandrinia neesiana H.Eichler = Calandrinia granulifera | 1:78 |
| | Calandrinia pygmaea F.Muell. nom. illeg. = Calandrinia granulifera | |
| | Claytonia australasica Hook.f. = Montia australasica | 1:79 |
| | Claytonia perfoliata Donn ex Willd. subsp. perfoliata previously listed as naturalised but insufficient evidence exists to support this | |
| | Montia australasica (Hook.f.) Pax & K.Hoffm., Nat. Pflanzenfam., ed. 2 [Engler & Prantl] 16c: 259 (1934) | 1:79 |
| | Montia fontana L. subsp. chondrosperma (Fenzl) Walters, Watsonia 3: 4 (1953) | 1:79 |
| | Neopaxia australasica (Hook.f.) O.Nilsson = Montia australasica | 1:79 |
| | Portulaca oleracea L., Sp. Pl. 1: 445 (1753) | 1:78 |
| PROTEACEAE | | FTO 48 |
| e | Agastachys odorata R.Br., Trans. Linn. Soc. London 10: 158 (1810) | 3:601 |
| | Banksia australis R.Br. = Banksia marginata | |
| | Banksia depressa R.Br. = Banksia marginata | |
| | Banksia insularis R.Br. = Banksia marginata | |
| x | Banksia integrifolia L.f. subsp. integrifolia , Suppl. Pl. 127 (1782) | 3:616 |
| | Banksia marginata Cav., Anales Hist. Nat. 1(3): 227, t.13 (1800) | 3:615 |
| | Banksia media R.Br. sensu Hooker (1860) = Banksia serrata (misapplied in Tasmania) | |
| | Banksia serrata L.f., Suppl. Pl. 126 (1782) | 3:616 |
| e | Bellenden montana R.Br., Trans. Linn. Soc. London 10: 166 (1810) | 3:600 |
| e | Cenarrhenes nitida Labill., Nov. Holl. Pl. 1: 36, t.50 (1805) | 3:601 |
| e | Conospermum hookeri (Meisn.) E.M.Benn., Fl. Australia 16: 485 (1995) | 3:600 |
| | Conospermum taxifolium C.F.Gaertn. sensu Curtis (1967) = Conospermum hookeri (misapplied in Tasmania) | 3:600 |
| | Embothrium tinctorium Labill. = Lomatia tinctoria | |
| | Embothrium truncatum Labill. = Telopea truncata | |
| i | Grevillea hybrids | |
| | Grevillea australis R.Br., Trans. Linn. Soc. London 10: 171 (1810) | 3:606 |

| | | |
|-----|---|-------|
| | <i>Grevillea australis</i> R.Br. var. <i>brevifolia</i> Hook.f. = Grevillea australis | 3:606 |
| | <i>Grevillea australis</i> R.Br. var. <i>erecta</i> Hook.f. = Grevillea australis | 3:606 |
| | <i>Grevillea australis</i> R.Br. var. <i>linearifolia</i> Hook.f. = Grevillea australis | 3:606 |
| | <i>Grevillea australis</i> R.Br. var. <i>montana</i> Hook.f. = Grevillea australis | 3:606 |
| | <i>Grevillea australis</i> R.Br. var. <i>planifolia</i> Hook.f. = Grevillea australis | 3:606 |
| | <i>Grevillea australis</i> R.Br. var. <i>subulata</i> Hook.f. = Grevillea australis | 3:606 |
| | <i>Grevillea australis</i> R.Br. var. <i>tenuifolia</i> (R.Br.) Meisn. = Grevillea australis | 3:606 |
| | <i>Grevillea seymouriae</i> Sweet ex Meisn. sensu Meisner (1856) mistakenly attributed to Tasmania | |
| | <i>Grevillea stuartii</i> Meisn. sensu Hooker (1860) = Grevillea australis | |
| | <i>Grevillea tenuifolia</i> R.Br. = Grevillea australis | |
| | <i>Hakea acicularis</i> (Sm. ex Vent.) Knight nom. illeg. sensu Hooker (1860) = Hakea decurrens subsp. platytaenia (misapplied in Tasmania) | |
| | <i>Hakea acicularis</i> (Sm. ex Vent.) Knight var. <i>lissosperma</i> (R.Br.) Benth. = Hakea lissosperma | |
| | Hakea decurrens R.Br. subsp. physocarpa W.R.Barker, J. Adelaide Bot. Gard. 17: 193 (1996) | 3:610 |
| | Hakea decurrens R.Br. subsp. platytaenia W.R.Barker, J. Adelaide Bot. Gard. 17: 196 (1996) | |
| e | Hakea epiglottis Labill. subsp. epiglottis , Nov. Holl. Pl. 1: 30, t.40 (1805) | 3:609 |
| e | Hakea epiglottis Labill. subsp. milliganii (Meisn.) R.M.Barker, Fl. Australia 17B: 394 (1999) | |
| i x | Hakea laurina R.Br., Suppl. Prodr. Fl. Nov. Holl. 29 (1830) | |
| | Hakea lissosperma R.Br., Trans. Linn. Soc. London 10: 180 (1810) | 3:611 |
| e | Hakea megadenia R.M.Barker, Aspects of Tasmanian Botany: 83 (1991) | |
| | Hakea microcarpa R.Br., Trans. Linn. Soc. London 10: 182 (1810) | 3:611 |
| | Hakea nodosa R.Br., Trans. Linn. Soc. London 10: 179 (1810) | 3:610 |
| | <i>Hakea pugioniformis</i> Cav. = Hakea teretifolia | |
| | <i>Hakea rostrata</i> F.Muell. ex Meisn. sensu Curtis (1967) = Hakea epiglottis & H. megadenia (misapplied in Tasmania) | 3:609 |
| | <i>Hakea rugosa</i> R.Br. sensu Curtis (1967) = Hakea epiglottis & H. megadenia (misapplied in Tasmania) | 3:609 |
| i | Hakea salicifolia (Vent.) B.L.Burt subsp. salicifolia , Bull. Misc. Inform. Kew 1: 33 (1941) | |
| | <i>Hakea sericea</i> Schrad. & J.C.Wendl. sensu Curtis (1967) = Hakea decurrens (misapplied in Tasmania) | 3:610 |
| | Hakea teretifolia (Salisb.) Britten subsp. hirsuta (Endl.) R.M.Barker, J. Adelaide Bot. Gard. 13: 105 (1990) | 3:609 |
| | Hakea ulicina R.Br., Suppl. Prodr. Fl. Nov. Holl. 29 (1830) | 3:611 |
| | <i>Hylogyne australis</i> R.Br. = Telopea truncata | |
| | Isopogon ceratophyllus R.Br., Trans. Linn. Soc. London 10: 72 (1810) | 3:599 |
| i # | Lomatia fraseri R.Br., Suppl. Prodr. Fl. Nov. Holl. 34 (1830) | |
| e | Lomatia polymorpha R.Br., Trans. Linn. Soc. London 10: 200 (1810) | 3:613 |
| e | Lomatia tasmanica W.M.Curtis, The Student's Flora of Tasmania 3: 651 (1967) | 3:614 |
| e | Lomatia tinctoria (Labill.) R.Br., Trans. Linn. Soc. London 10: 199 (1810) | 3:613 |
| e | Orites acicularis (R.Br.) Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 3: 427 (1818) | 3:605 |
| e | Orites diversifolius R.Br., Trans. Linn. Soc. London 10: 190 (1810) | 3:604 |
| e | Orites milliganii Meisn., Prodr. [A. P. de Candolle] 14: 424 (1856) | 3:604 |
| e | Orites revolutus R.Br., Trans. Linn. Soc. London 10: 190 (1810) | 3:605 |
| e | Persoonia gunnii Hook.f., London J. Bot. 6: 283 (1847) | |

| | | |
|----|--|---------------|
| | <i>Persoonia gunnii</i> Hook.f. var. <i>angustifolia</i> Benth. = <i>Persoonia muelleri</i> subsp. <i>angustifolia</i> | 3:603 |
| | <i>Persoonia gunnii</i> Hook.f. var. <i>oblanceolata</i> Orchard = <i>Persoonia gunnii</i> | 3:603 |
| | <i>Persoonia juniperina</i> Labill., Nov. Holl. Pl. 1: 33, t.45 (1805) | 3:603 |
| | <i>Persoonia juniperina</i> Labill. var. <i>brevifolia</i> Meisn. = <i>Persoonia juniperina</i> | 3:603 |
| | <i>Persoonia juniperina</i> Labill. var. <i>mollis</i> Orchard = <i>Persoonia juniperina</i> | 3:603 |
| | <i>Persoonia juniperina</i> Labill. var. <i>ulicina</i> Meisn. = <i>Persoonia juniperina</i> | 3:603 |
| e | <i>Persoonia moscalii</i> Orchard, Brunonia 6: 238, fig.12 (1984) | |
| e | <i>Persoonia muelleri</i> (P.Parm.) Orchard subsp. <i>angustifolia</i> (Benth.) L.A.S.Johnson & P.H.Weston, Fl. Australia 16: 472 (1995) | 3:603 |
| e | <i>Persoonia muelleri</i> (P.Parm.) Orchard subsp. <i>densifolia</i> (Orchard) L.A.S.Johnson & P.H.Weston, Fl. Australia 16: 472 (1995) | |
| e | <i>Persoonia muelleri</i> (P.Parm.) Orchard subsp. <i>muelleri</i> , Brunonia 6: 226 (1984) | |
| e | <i>Telopea truncata</i> (Labill.) R.Br., Trans. Linn. Soc. London 10: 198 (1810) | 3:612 |
| | RANUNCULACEAE | FTO 47 |
| i* | <i>Adonis microcarpa</i> DC., Syst. Nat. [Candolle] 1: 223 (1817) | |
| e | <i>Anemone crassifolia</i> Hook., Icon. Pl. 3(6): t.257 (1840) | 1:11 |
| i | <i>Aquilegia vulgaris</i> L., Sp. Pl. 1: 533 (1753) | |
| | <i>Batrachium trichophyllum</i> (Chaix) F.W.Schultz = <i>Ranunculus trichophyllus</i> | 1:14 |
| | <i>Caltha introloba</i> F.Muell. sensu Bentham (1863) = <i>Psychrophila phylloptera</i> (misapplied in Tasmania) | |
| | <i>Caltha novae-zelandiae</i> Hook. sensu Rodway (1903) = <i>Psychrophila phylloptera</i> (misapplied in Tasmania) | |
| | <i>Caltha phylloptera</i> A.W.Hill = <i>Psychrophila phylloptera</i> | 1:20 |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl., Bot. Reg. 3: t.238 (1817) | 1:9 |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl. subsp. <i>confertissima</i> Kuntze = <i>Clematis aristata</i> | |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl. subsp. <i>gentianoides</i> (DC.) Kuntze = <i>Clematis gentianoides</i> | |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl. subsp. <i>gunniana</i> Kuntze = <i>Clematis clitorioides</i> | |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl. subsp. <i>procumbens</i> Kuntze = <i>Clematis gentianoides</i> | |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl. subsp. <i>tasmanica</i> Kuntze = <i>Clematis gentianoides</i> | |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl. var. <i>blanda</i> (Hook.) Benth. = <i>Clematis clitorioides</i> | |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl. var. <i>coriacea</i> (DC.) Benth. = <i>Clematis aristata</i> | |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl. var. <i>gentianoides</i> (DC.) F.Muell. = <i>Clematis gentianoides</i> | |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl. var. <i>gunniana</i> (Kuntze) Domin = <i>Clematis clitorioides</i> | |
| | <i>Clematis aristata</i> R.Br. ex Ker Gawl. var. <i>minor</i> Hook.f. = <i>Clematis clitorioides</i> | |
| | <i>Clematis blanda</i> Hook. = <i>Clematis clitorioides</i> | |
| | <i>Clematis clitorioides</i> DC., Syst. Nat. [Candolle] 1: 158 (1817) | |
| | <i>Clematis clitorioides</i> DC. var. <i>decipiens</i> Domin = <i>Clematis clitorioides</i> | |
| | <i>Clematis coriacea</i> DC. = <i>Clematis aristata</i> | |
| | <i>Clematis decipiens</i> H.Eichler ex Jeanes, Fl. Australia 2: 461 (2007) | |
| e | <i>Clematis gentianoides</i> DC., Syst. Nat. [Candolle] 1: 159 (1817) | 1:10 |
| | <i>Clematis gentianoides</i> DC. var. <i>normalis</i> Domin nom. inval. = <i>Clematis gentianoides</i> | |
| | <i>Clematis gentianoides</i> var. <i>procumbens</i> = <i>Clematis gentianoides</i> | |
| | <i>Clematis gentianoides</i> DC. var. <i>tasmanica</i> (Kuntze) Domin = <i>Clematis gentianoides</i> | |
| | <i>Clematis hexapetala</i> L.f. subsp. <i>brachystemon</i> Kuntze = <i>Clematis clitorioides</i> | |

| | | |
|---|--|------|
| | Clematis linearifolia Steud. sensu Hooker (1860) = Clematis clitorioides (misapplied in Tasmania) | |
| | Clematis microphylla DC., Syst. Nat. [Candolle] 1: 147 (1817) | 1:10 |
| i | Clematis vitalba L. var. vitalba , Sp. Pl. 1: 544 (1753) | 1:10 |
| | Myosurus australis F.Muell., Trans. Philos. Soc. Victoria 1: 6 (1854) | 1:11 |
| | Myosurus minimus L. sensu Curtis & Morris (1975) = Myosurus australis (misapplied in Tasmania) | 1:11 |
| e | Psychrophila phylloptera (A.W.Hill) H.Eichler, Fl. Australia 2: 459 (2007) | 1:20 |
| t | Ranunculus acaulis Banks & Sol. ex DC., Syst. Nat. [Candolle] 1: 270 (1817) | 1:17 |
| i | Ranunculus acris L. subsp. acris , Sp. Pl. 1: 554 (1753) | 1:18 |
| | Ranunculus amphitrichus Colenso, Trans. & Proc. New Zealand Inst. 17: 237 (1885) | 1:17 |
| | Ranunculus arvensis L. previously listed as naturalised but insufficient evidence exists to support this | 1:18 |
| e | Ranunculus collicola Menadue, Brunonia 8(2): 373 (1986) | |
| | Ranunculus collinus R.Br. ex DC., Syst. Nat. [Candolle] 1: 271 (1817) | 1:17 |
| | Ranunculus concinnus (Hook.f.) Melville = Ranunculus decurvus | 1:15 |
| | Ranunculus cuneatus Hook. nom. illeg. = Ranunculus triplodontus | |
| e | Ranunculus decurvus (Hook.f.) Melville, Kew Bull. 10: 202 (1955) | 1:15 |
| | Ranunculus diminutus B.G.Briggs, Telopea 5(4): 583 (1994) | |
| | Ranunculus flammula L. subsp. flammula previously listed as naturalised but insufficient evidence exists to support this | |
| | Ranunculus glabrifolius Hook., J. Bot. (Hooker) 1: 243 (1834) | 1:17 |
| | Ranunculus gunnianus Hook., J. Bot. (Hooker) 1: 244, t.133 (1834) | 1:14 |
| | Ranunculus hirtus Banks & Sol. ex DC. nom. illeg. = Ranunculus pimpinellifolius | |
| | Ranunculus inconspicuus Hook.f. = Ranunculus collinus | |
| | Ranunculus inundatus R.Br. ex DC. sensu Curtis & Morris (1975) = Ranunculus amphitrichus (misapplied in Tasmania) | 1:17 |
| e | Ranunculus jugosus Menadue, Brunonia 8(2): 377 (1986) | |
| | Ranunculus lappaceus Sm., Cycl. (Rees) 29(2): no.61 (1815) | 1:14 |
| | Ranunculus lappaceus Sm. var. nanus (Hook.) Benth. = Ranunculus nanus | |
| | Ranunculus lappaceus Sm. var. pimpinellifolius (Hook.) Benth. = Ranunculus pimpinellifolius | |
| | Ranunculus lappaceus Sm. var. scapiger (Hook.) Benth. = Ranunculus scapiger | |
| | Ranunculus lappaceus Sm. var. subsericeus Benth. = Ranunculus pascuinus | |
| | Ranunculus millanii F.Muell. sensu Rodway (1903) = Ranunculus setaceus (misapplied in Tasmania) | |
| i | Ranunculus muricatus L., Sp. Pl. 1: 555 (1753) | 1:18 |
| e | Ranunculus nanus Hook., J. Bot. (Hooker) 1: 242 (1834) | 1:16 |
| i | Ranunculus parviflorus L., Syst. Nat., ed. 10, 2: 1087 (1759) | 1:19 |
| e | Ranunculus pascuinus (Hook.f.) Melville, Kew Bull. 10: 198 (1955) | 1:15 |
| | Ranunculus philonotis Retz. sensu Bentham (1863) = Ranunculus sardous (misapplied in Tasmania) | |
| | Ranunculus pimpinellifolius Hook., J. Bot. (Hooker) 1: 243 (1834) | 1:14 |
| | Ranunculus plebeius R.Br. ex DC. sensu Curtis (1956) mistakenly attributed to Tasmania | |
| e | Ranunculus prasinus Menadue, Brunonia 8: 375 (1986) | |
| | Ranunculus pumilio R.Br. ex DC. var. pumilio , Syst. Nat. [Candolle] 1: 271 (1817) | 1:19 |
| i | Ranunculus repens L., Sp. Pl. 1: 554 (1753) | 1:18 |
| | Ranunculus rivularis Banks & Sol. ex DC. sensu Curtis & Morris (1975) = Ranunculus amphitrichus | 1:17 |
| | Ranunculus rivularis Banks & Sol. ex DC. var. inconspicuus Benth. = Ranunculus collinus | |

| | | |
|----|--|---------------|
| | Ranunculus rivularis Banks & Sol. ex DC. var. major Benth. = Ranunculus glabrifolius | |
| | Ranunculus rivularis Banks & Sol. ex DC. var. subfluitans Benth. nom. illeg. = Ranunculus amphitrichus | |
| i | Ranunculus sardous Crantz, Stirp. Austr. Fasc. 2: 84 (1763) | 1:19 |
| | Ranunculus scapiger Hook., J. Bot. (Hooker) 1: 224 (1834) | 1:15 |
| | Ranunculus sceleratus L. subsp. sceleratus previously listed as naturalised but insufficient evidence exists to support this | 1:20 |
| | Ranunculus sessiliflorus R.Br. ex DC. var. sessiliflorus , Syst. Nat. [Candolle] 1: 302 (1817) | 1:19 |
| e | Ranunculus setaceus Rodway, Pap. & Proc. Roy. Soc. Tasmania 1900-1901: 107 (1902) | 1:14 |
| ?i | Ranunculus trichophyllus Chaix, Hist. Pl. Dauphiné (Villars) 1: 335 (1786) | 1:14 |
| | Ranunculus trichophyllus Chaix subsp. drouetii (F.W.Schultz ex Godr.) A.R.Clapham = Ranunculus trichophyllus | |
| i | Ranunculus trilobus Desf., Fl. Atlant. 1: 437, t.113 (1798) | 1:19 |
| e | Ranunculus triplodontus Melville, Kew Bull. 10: 204 (1955) | 1:16 |
| | Valvaria gentianoides (DC.) Ser. = Clematis gentianoides | |
| | RESEDACEAE | FTO 84 |
| i | Reseda alba L., Sp. Pl. 1: 449 (1753) | 1:53 |
| i | Reseda lutea L., Sp. Pl. 1: 449 (1753) | 1:52 |
| i | Reseda luteola L., Sp. Pl. 1: 448 (1753) | 1:52 |
| | Reseda odorata L. sensu Baker & de Salas (2012) = Reseda lutea (misapplied in Tasmania) | |
| | RHAMNACEAE | |
| e | Cryptandra alpina Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 75, t.12B (1855) | 1:121 |
| | Cryptandra amara Sm., Cycl. (Rees) 10: no.2 (1808) | 1:120 |
| e | Cryptandra exilis D.I.Morris, Aspects of Tasmanian Botany: 57 (1991) | |
| | Cryptandra mollis Hook.f. = Spyridium parvifolium var. molle | |
| | Cryptandra obovata Hook.f. = Spyridium obovatum | |
| | Cryptandra parvifolia Hook.f. = Spyridium parvifolium var. parvifolium | |
| | Cryptandra pimelioides Hook.f. = Stenanthemum pimeleoides | |
| | Cryptandra sieberi Fenzl = Cryptandra amara | |
| | Cryptandra tomentosa Lindl. sensu Buchanan et al. (1989) = Cryptandra exilis | |
| | Cryptandra vexillifera Hook. = Spyridium vexilliferum | |
| | Discaria australis Hook. nom. illeg. = Discaria pubescens (misapplied in Tasmania) | |
| | Discaria pubescens (Brongn.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 620 (1917) | 1:121 |
| | Pomaderris affinis N.A.Wakef. = Pomaderris intermedia | 1:114 |
| | Pomaderris apetala Labill. subsp. apetala , Nov. Holl. Pl. 1: 62, t.87 (1805) | 1:115 |
| | Pomaderris apetala Labill. subsp. maritima N.G.Walsh & Coates, Muelleria 10: 34 (1997) | |
| | Pomaderris aspera Sieber ex DC., Prodr. [A. P. de Candolle] 2: 33 (1825) | |
| | Pomaderris discolor (Vent.) Poir. sensu Hooker (1860) = Pomaderris intermedia (misapplied in Tasmania) | |
| | Pomaderris elachophylla F.Muell., Fragm. (Mueller) 2: 131 (1861) | 1:116 |
| e | Pomaderris elliptica Labill. var. diemenica N.G.Walsh & Coates, Muelleria 10: 51 (1997) | |
| | Pomaderris elliptica Labill. var. elliptica , Nov. Holl. Pl. 1: 61, t.86 (1805) | 1:114 |
| | Pomaderris ferruginea Sieber ex Fenzl sensu Bentham (1863) recorded in error | |

| | | |
|-----------------|---|-------|
| | Pomaderris intermedia Sieber ex DC., Prodr. [A. P. de Candolle] 2: 33 (1825) | 1:114 |
| | <i>Pomaderris kumeraho</i> A.Cunn. sensu Curtis (1956) mistakenly attributed to Tasmania | |
| | Pomaderris oraria F.Muell. ex Reissek subsp. oraria , Linnaea 29: 268 (1858) | 1:116 |
| | Pomaderris paniculosa F.Muell. ex Reissek subsp. paralia N.G.Walsh, Muelleria 7: 274 (1990) | |
| | Pomaderris phyllicifolia Lodd. ex Link subsp. ericoides (Maiden & Betche) N.G.Walsh & Coates, Muelleria 10: 52 (1997) | |
| | Pomaderris phyllicifolia Lodd. ex Link subsp. phyllicifolia , Enum. Hort. Berol. Alt. 1: 232 (1821) | 1:116 |
| | Pomaderris pilifera N.A.Wakef. subsp. pilifera , Vict. Naturalist 68: 140 (1951) | 1:115 |
| e | Pomaderris pilifera N.A.Wakef. subsp. talpicutica A.M.Gray & Wapstra, Muelleria 25: 129 (2007) | |
| | Pomaderris racemosa Hook., J. Bot. (Hooker) 1: 256 (1834-35) | 1:116 |
| i | Rhamnus alaternus L., Sp. Pl. 1: 193 (1753) | |
| | Spyridium eriocephalum Fenzl var. eriocephalum , Enum. Pl. [Endlicher]: 24 (1837) | 1:119 |
| e | Spyridium gunnii (Hook.f.) Benth., Fl. Austral. 1: 429 (1863) | 1:118 |
| e | Spyridium lawrencei (Hook.f.) Benth., Fl. Austral. 1: 430 (1863) | 1:119 |
| | <i>Spyridium microphyllum</i> (F.Muell. ex Reissek) Druce = Spyridium lawrencei | 1:119 |
| e | Spyridium obcordatum (Hook.f.) W.M.Curtis, Vict. Naturalist 87: 251 (1970) | 1:117 |
| | <i>Spyridium obovatum</i> (Hook.f.) Benth. var. <i>gunnii</i> (Hook.f.) Rodway = Spyridium gunnii | |
| e | Spyridium obovatum (Hook.) Benth. var. obovatum , Fl. Austral. 1: 429 (1863) | 1:118 |
| e | Spyridium obovatum (Hook.) Benth. var. velutinum (F.Muell. ex Reissek) Benth., Fl. Austral. 1: 429 (1863) | 1:118 |
| e | Spyridium parvifolium (Hook.) F.Muell. var. molle (Hook.f.) Benth., Fl. Austral. 1: 428 (1863) | 1:118 |
| | Spyridium parvifolium (Hook.) F.Muell. var. parvifolium , Fragm. (Mueller) 3: 79 (1862) | 1:118 |
| | <i>Spyridium serpyllaceum</i> (Reissek & F.Muell.) F.Muell. nom. illeg. = Spyridium obcordatum | |
| e | Spyridium ulicinum (Hook.) Benth., Fl. Austral. 1: 434 (1863) | 1:119 |
| | Spyridium vexilliferum (Hook.) Reissek var. vexilliferum , Linnaea 29: 285 (1858) | 1:119 |
| e | Stenanthemum pimeleoides (Hook.f.) Benth., Fl. Austral. 1: 436 (1863) | 1:120 |
| ROSACEAE | | |
| | <i>Acaena agnipila</i> Gand. var. <i>aequispina</i> Orchard = Acaena ovina | 1:176 |
| | <i>Acaena agnipila</i> Gand. var. <i>tenuispica</i> (Bitter) Orchard = Acaena ovina | 1:176 |
| | <i>Acaena anserinifolia</i> (J.R.Forst. & G.Forst.) Druce sensu Curtis (1956) = Acaena novae-zelandiae | |
| | Acaena × anserovina Orchard, Trans. Roy. Soc. South Australia 93: 104 (1969) | 1:176 |
| | Acaena echinata Nees, Pl. Preiss. [J.G.C.Lehman] 1: 95 (1844) | 1:175 |
| | <i>Acaena echinata</i> Nees var. <i>retrorsumpilosa</i> (Bitter) Orchard = Acaena echinata | 1:175 |
| | <i>Acaena echinata</i> Nees var. <i>subglabralyx</i> (Bitter) Orchard = Acaena echinata | 1:175 |
| | <i>Acaena echinata</i> Nees var. <i>tylacantha</i> Orchard = Acaena echinata | 1:175 |
| e | Acaena montana Hook.f., London J. Bot. 6: 476 bis (1847) | 1:175 |
| | Acaena novae-zelandiae Kirk, Trans. Proc. New Zealand Inst. 3: 177 (1871) | 1:175 |
| | Acaena ovina A.Cunn., Geog. Mem. New South Wales [Field]: 358 (1825) | 1:175 |
| | <i>Acaena ovina</i> A.Cunn. var. <i>velutina</i> Orchard = Acaena ovina | 1:175 |
| | Acaena pallida (Kirk) Allan, Fl. New Zealand 1: 360 (1961) | |
| | <i>Acaena sanguisorbae</i> (L.) Vahl sensu Bentham (1864) = Acaena novae-zelandiae (misapplied in Tasmania) | |
| | <i>Acaena sanguisorbae</i> (L.) Vahl var. <i>montana</i> (Hook.f.) Hook.f. = Acaena montana | |

| | | |
|-----|--|-------|
| | Alchemilla arvensis (L.) Scop. = Aphanes arvensis | |
| i | Aphanes arvensis L., Sp. Pl. 1: 123 (1753) | 1:174 |
| | Aphanes australiana (Rothm.) Rothm., Bull. Misc. Inform. Kew 1938: 270 (1938) | |
| | Aphanes inexpectata W.Lippert sensu Buchanan (1999) = Aphanes microcarpa | |
| i | Aphanes microcarpa (Boiss. & Reut.) Rothm., Repert. Spec. Nov. Regni Veg. 42: 172 (1937) | |
| ?i | Argentina anserina (L.) Rydb., Mem. Dept. Bot. Columbia Coll. 2: 159 (1898) | 1:173 |
| i | Cotoneaster franchetii Bois, Rev. Hort. [Paris] 74: 379 (1902) | |
| i | Cotoneaster glaucophyllus Franch. var. serotinus (Hutch.) L.T.Lu & Brach, Novon 12: 495 (2002) | |
| i | Cotoneaster pannosus Franch., Pl. Delavay. 223 (1890) | |
| i | Cotoneaster simonsii Baker, Refug. Bot. [Saunders] 1: t.55 (1869) | |
| | Cotoneaster symondsii T.Moore = Cotoneaster simonsii | |
| i | Crataegus monogyna Jacq., Fl. Austriac. (Jacquin) 3: 50 (1775) | 1:178 |
| | Geum renifolium F.Muell. = Geum talbotianum | |
| e | Geum talbotianum W.M.Curtis, Records of the Queen Victoria Museum 50: 4 (1974) | 1:172 |
| x | Geum urbanum L. var. strictum Hook.f., Bot. Antarct. Voy. III (Fl. Tasman.) 1: 114 (1856) | 1:172 |
| | Malus × domestica Borkh. = Malus pumila | |
| i | Malus pumila Mill., Gard. Dict., ed. 8, Malus no 3 (1768) | |
| i | Potentilla anglica Laichard., Veg. Eur. 1: 475 (1790) | 1:173 |
| | Potentilla anserina L. = Argentina anserina | 1:173 |
| i | Potentilla recta L., Sp. Pl. 1: 497 (1753) | 1:173 |
| i | Potentilla reptans L., Sp. Pl. 1: 499 (1753) | 1:173 |
| | Poterium polygamum Waldst. & Kit. = Sanguisorba minor | 1:176 |
| | Poterium sanguisorba L. sensu Rodway (1903) = Sanguisorba minor | 1:176 |
| i | Prunus domestica L. subsp. insititia (L.) Bonnier & Layens, Fl. France [Rouy & Foucard]: 95 (1894) | 1:178 |
| | Prunus insititia L. = Prunus domestica subsp. insititia | |
| i | Prunus laurocerasus L., Sp. Pl. 1: 474 (1753) | |
| i | Prunus spinosa L., Sp. Pl. 1: 475 (1753) | 1:178 |
| i | Rosa canina L., Sp. Pl. 1: 491 (1753) | 1:177 |
| i | Rosa rubiginosa L., Mant. Pl. 2: 564 (1771) | 1:177 |
| i | Rubus anglocandicans A.Newton, Watsonia 11: 243 (1977) | 1:171 |
| i | Rubus echinatus Lindl., Syn. Brit. Fl. 94 (1829) | |
| i | Rubus erythrops Eedes & A.Newton, Watsonia 12: 135 (1978) | |
| | Rubus fruticosus L. a catch-all name for all the blackberry species of Rubus | |
| e | Rubus gunnianus Hook., Icon. Pl. 3: t.291 (1840) | 1:171 |
| i | Rubus laciniatus Willd., Hort. Berol. [Willdenow] 2(8): t.82 (1806) | 1:171 |
| i | Rubus leucostachys Schleich. ex Sm., Engl. Fl. 2: 403 (1824) | 1:171 |
| i | Rubus loganobaccus L.H.Bailey, Gentes Herb. 1: 155 (1923) | |
| | Rubus macropodus Ser. ex DC. sensu Hooker (1860) = Rubus parvifolius | |
| | Rubus parvifolius L., Sp. Pl. 2: 1197 (1753) | 1:171 |
| i | Rubus philadelphicus Blanch., Torreyia 7: 56 (1907) | |
| i | Rubus polyanthemus Lindeb., Bot. Not. 1883: 105 (1883) | |
| i | Rubus rubritinctus W.C.R.Watson, Watsonia 3: 287 (1956) | |
| i # | Rubus rugosus Sm., Cycl. (Rees) 30: no 34 (1815) | |

| | | |
|------------------|---|-------|
| i | Rubus sp. Tasmania (J.R.Hosking 1551) SA Herbarium | |
| i | Rubus vestitus Weihe, Comp. Fl. German. 1: 684 (1825) | |
| i | Sanguisorba minor Scop., Fl. Carniol., ed. 2, 1: 110 (1771) | 1:176 |
| | Sanguisorba minor Scop. subsp. muricata (Spach) Briq. = Sanguisorba minor | 1:176 |
| i | Sorbus aucuparia L., Sp. Pl. 1: 477 (1753) | |
| RUBIACEAE | | |
| | <i>Asperula charophyton</i> Airy Shaw & Turrill sensu Curtis (1963) recorded in error | 2:276 |
| | Asperula conferta Hook.f., London J. Bot. 6: 464 bis (1847) | 2:275 |
| | <i>Asperula conferta</i> Hook.f. var. <i>abbreviata</i> Airy Shaw & Turrill = Asperula conferta | 2:275 |
| | <i>Asperula conferta</i> Hook.f. var. <i>scoparioides</i> Airy Shaw & Turrill = Asperula conferta | 2:275 |
| | Asperula gunnii Hook.f., London J. Bot. 6: 464 bis (1847) | 2:275 |
| | <i>Asperula gunnii</i> Hook.f. var. <i>curta</i> (Hook.f.) Airy Shaw & Turrill = Asperula gunnii | 2:276 |
| | <i>Asperula gunnii</i> Hook.f. var. <i>pusilla</i> (Hook.f.) Benth. = Asperula pusilla | |
| | Asperula minima Hook.f., London J. Bot. 6: 464 bis (1847) | 2:276 |
| | Asperula oblanceolata I.Thomps., Muelleria 27: 68 (2009) | |
| | <i>Asperula oligantha</i> F.Muell. nom. illeg., sensu Rodway (1903) a catch-all name that includes most Tasmanian Asperula species | |
| | Asperula pusilla Hook.f., London J. Bot. 6: 464 bis (1847) | 2:276 |
| | Asperula scoparia Hook.f. subsp. scoparia , London J. Bot. 6: 463 bis (1847) | 2:275 |
| | Asperula subsimplex Hook.f., London J. Bot. 6: 463 bis (1847) | 2:276 |
| | <i>Canthium quadrifidum</i> Labill. = Coprosma quadrifida | |
| | <i>Coprosma billardierei</i> Hook.f. nom. illeg. = Coprosma quadrifida | |
| | Coprosma hirtella Labill., Nov. Holl. Pl. 1: 70, t.95 (1805) | 2:268 |
| | Coprosma moorei F.Muell. ex Rodway, Pap. & Proc. Roy. Soc. Tasmania 1893: 179, t.1 (1894) | 2:269 |
| | Coprosma nitida Hook.f., London J. Bot. 6: 465 bis (1847) | 2:269 |
| | Coprosma nivalis W.R.B.Oliv., Bull. Bernice P. Bishop Mus. 132: 37 (1935) | |
| | Coprosma perpusilla Colenso subsp. perpusilla , Trans. & Proc. New Zealand Inst. 22: 466 (1890) | |
| | Coprosma pumila Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.). 2: 543 (1847) | 2:269 |
| | Coprosma quadrifida (Labill.) B.L.Rob., Proc. Amer. Acad. Arts 45: 409 (1910) | 2:268 |
| i | Coprosma repens A.Rich., Voy. Astrolabe 1: 264 (1832) | |
| i | Coprosma robusta Raoul, Ann. Sci. Nat., Bot., sér. 3, 2: 121 (1844) | |
| e | Galium albescens Hook.f., London J. Bot. 6: 462 bis (1847) | 2:273 |
| i | Galium aparine L., Sp. Pl. 1: 108 (1753) | 2:273 |
| | Galium australe DC., Prodr. [A. P. de Candolle] 4: 608 (1830) | 2:273 |
| | Galium binifolium N.A.Wakef. subsp. conforme I.Thomps., Muelleria 27: 88 (2009) | |
| | Galium ciliare Hook.f. subsp. ciliare , London J. Bot. 6: 461 bis (1847) | 2:272 |
| | Galium ciliare Hook.f. subsp. terminale I.Thomps., Muelleria 27: 82 (2009) | |
| | Galium compactum Ehrend. & McGill., Telopea 2: 370 (1983) | |
| | Galium densum Hook.f., London J. Bot. 6: 461 bis (1847) | |
| i | Galium divaricatum Pourr. ex Lam., Encycl. (Lamarck) 2: 580 (1788) | |
| | Galium gaudichaudii DC. subsp. parviflorum I.Thomps., Muelleria 27: 98 (2009) | 2:272 |
| | Galium leiocarpum I.Thomps., Muelleria 27: 79 (2009) | |
| i | Galium murale (L.) All., Fl. Pedem. 1: 8 (1785) | 2:273 |

| | | |
|-------|---|---------------|
| i | Galium palustre L., Sp. Pl. 1: 105 (1753) | |
| | Galium squalidum Hook.f. = Galium australe | |
| | Galium tricornutum Dandy previously listed as naturalised but insufficient evidence exists to support this | |
| | Galium umbrosum G.Forst. nom. illeg., sensu Rodway (1903) a catch-all name that includes all glabrous-fruited Tasmanian Galium | |
| | Galium vagans Hook.f. = Galium binifolium subsp. conforme (uncertain) | |
| i # t | Galium verum L., Sp. Pl. 1: 107 (1753) | |
| | Nertera depressa Banks & Sol. ex Gaertn. = Nertera granadensis | 2:270 |
| | Nertera granadensis (Mutis ex L.f.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 637 (1917) | 2:270 |
| | Opercularia ovata Hook.f., London J. Bot. 6: 465 bis (1847) | 2:270 |
| | Opercularia varia Hook.f., London J. Bot. 6: 466 bis (1847) | 2:271 |
| i | Sherardia arvensis L., Sp. Pl. 1: 102 (1753) | 2:277 |
| | RUTACEAE | FTO 87 |
| e | Acradenia frankliniae Kippist, Proc. Linn. Soc. London 2: 201 (1853) | 1:106 |
| | Boronia anemonifolia A.Cunn. subsp. anemonifolia recorded in error | |
| | Boronia anemonifolia A.Cunn. subsp. variabilis (Hook.) P.G.Neish, Muelleria 14: 11 (2000) | 1:105 |
| | Boronia anemonifolia A.Cunn. var. dentigera (F.Muell.) Benth. = Boronia anemonifolia subsp. anemonifolia | |
| e | Boronia citriodora Gunn ex Hook.f. subsp. citriodora , Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 68 (1855) | 1:105 |
| e | Boronia citriodora Gunn ex Hook.f. subsp. orientalis Duretto, Muelleria 17: 83 (2003) | |
| e | Boronia citriodora Gunn ex Hook.f. subsp. paulwilsonii Duretto, Muelleria 17: 81 (2003) | |
| e | Boronia elisabethiae Duretto, Muelleria 17: 88 (2003) | |
| e | Boronia gunnii Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 68, t.10 (1855) | |
| e | Boronia hemichiton Duretto, Muelleria 17: 87 (2003) | |
| e | Boronia hippopala Duretto, Muelleria 17: 84 (2003) | |
| | Boronia nana Hook. var. hyssopifolia Melville, Kew Bull. 9: 463 (1954) | 1:105 |
| | Boronia nana Hook. var. nana , Icon. Pl. 3: t.270 (1840) | 1:105 |
| | Boronia parviflora Sm., Tracts Nat. Hist. 295: t.6 (1798) | 1:105 |
| | Boronia pilonema Labill. = Boronia parviflora | |
| | Boronia pilosa Labill. subsp. pilosa , Nov. Holl. Pl. 1: 97, t.124 (1805) | 1:105 |
| e | Boronia pilosa Labill. subsp. tasmanensis Duretto, Muelleria 17: 97 (2003) | |
| | Boronia pilosa Labill. var. floribunda (Hook.) Hook.f. = Boronia pilosa subsp. pilosa | 1:105 |
| | Boronia pilosa Labill. var. laricifolia (Hook.) Hook.f. = Boronia pilosa subsp. pilosa | 1:105 |
| | Boronia pinnata Sm. var. citriodora (Gunn ex Hook.f.) Rodway = Boronia citriodora | |
| | Boronia pinnata Sm. var. gunnii (Hook.f.) Benth. = Boronia gunnii | |
| | Boronia polygalifolia Sm. sensu Bentham (1863) = Boronia nana var. nana (misapplied in Tasmania) | |
| | Boronia polygalifolia Sm. var. trifoliata Benth. = Boronia nana var. nana | |
| | Boronia rhomboidea Hook., Icon. Pl. 8: t.722 (1845) | 1:105 |
| e | Boronia rozefeldsii Duretto, Muelleria 17: 101 (2003) | |
| | Boronia tetrandra Labill. var. grandiflora Hook. = Boronia gunnii | |
| | Boronia variabilis Hook. = Boronia anemonifolia subsp. variabilis | |
| i | Coleonema pulchellum I.Williams, J. S. African Bot. 47: 89 (1981) | |

| | | |
|---|---|-------|
| | Correa alba Andrews var. alba , Bot. Repos. 1: t.18 (1798) | 1:109 |
| e | Correa alba Andrews var. rotundifolia DC., Prodr. [A. P. de Candolle] 1: 719 (1824) | 1:109 |
| | Correa backhouseana Hook. var. backhouseana , J. Bot. (Hooker) 1: 253 (1834) | 1:110 |
| e | Correa lawrenceana Hook. var. ferruginea Hook.f., Bot. Antarct. Voy. III (Fl. Tasman.) 1: 62 (1855) | |
| e | Correa lawrenceana Hook. var. lawrenceana , J. Bot. (Hooker) 1: 254 (1834) | 1:111 |
| | <i>Correa lawrenciana</i> Hook. var. <i>glabra</i> Benth. = Correa lawrenceana var. lawrenceana | |
| e | Correa reflexa (Labill.) Vent. var. nummulariifolia (Hook.f.) Paul G.Wilson, Trans. Roy. Soc. South Australia 85: 30 (1961) | 1:110 |
| | Correa reflexa (Labill.) Vent. var. reflexa , Jard. Malmaison 1: t.13 (1803) | 1:110 |
| | <i>Correa rufa</i> (Labill.) Vent. sensu Brown (1810) = Correa alba | |
| | <i>Correa speciosa</i> Donn ex Andrews = Correa reflexa | |
| | <i>Correa speciosa</i> Donn ex Andrews var. <i>backhouseana</i> (Hook.) Benth. = Correa backhouseana | |
| | <i>Correa speciosa</i> Donn ex Andrews var. <i>normalis</i> Benth. nom. inval., nom. superfl. = Correa reflexa | |
| | <i>Eriostemon daviesii</i> (Hook.f.) F.Muell. = Phebalium daviesii | |
| | <i>Eriostemon hillebrandii</i> F.Muell. nom. illeg., sensu Rodway (1903) = Leionema bilobum | |
| | <i>Eriostemon montanus</i> (Hook.f.) F.Muell. = Leionema montanum | |
| | <i>Eriostemon obovalis</i> A.Cunn. sensu Bentham (1863) = Philothea verrucosa (misapplied in Tasmania) | |
| | <i>Eriostemon oldfieldii</i> F.Muell. = Leionema oldfieldii | |
| | <i>Eriostemon squameus</i> Labill. = Nematolepis squamea | |
| | <i>Eriostemon verrucosus</i> A.Rich. = Philothea verrucosa | 1:107 |
| | <i>Eriostemon virgatus</i> A.Cunn. ex Hook. = Philothea virgata | 1:107 |
| e | Leionema bilobum (Lindl.) Paul G.Wilson subsp. truncatum (Hook.f.) Duretto & K.L.Durham, Muellera 23: 11 (2006) | 1:108 |
| e | Leionema montanum (Hook.) Paul G.Wilson, Nuytsia 12: 275 (1998) | 1:108 |
| e | Leionema oldfieldii (F.Muell.) Paul G.Wilson, Nuytsia 12: 276 (1998) | 1:108 |
| e | Nematolepis squamea (Labill.) Paul G.Wilson subsp. retusa (Hook.) Paul G.Wilson, Nuytsia 12: 279 (1998) | 1:109 |
| | Nematolepis squamea (Labill.) Paul G.Wilson subsp. squamea , Nuytsia 12: 279 (1998) | 1:109 |
| | <i>Phebalium billardierei</i> A.Juss. = Nematolepis squamea | |
| | <i>Phebalium bilobum</i> Lindl. = Leionema bilobum | 1:108 |
| e | Phebalium daviesii Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 358 (1859) | 1:108 |
| | <i>Phebalium glandulosum</i> Hook. var. <i>daviesii</i> (Hook.f.) Benth. = Phebalium daviesii | |
| | <i>Phebalium montanum</i> Hook. = Leionema montanum | 1:108 |
| | <i>Phebalium oldfieldii</i> (F.Muell.) Benth. = Leionema oldfieldii | 1:108 |
| | <i>Phebalium squameum</i> (Labill.) Engl. = Nematolepis squamea | 1:109 |
| | <i>Phebalium truncatum</i> Hook.f. = Leionema bilobum subsp. truncatum | |
| e | Philothea freyciana Rozefelds, Muellera 15: 23 (2001) | |
| | Philothea verrucosa (A.Rich.) Paul G.Wilson, Nuytsia 12: 260 (1998) | 1:107 |
| | Philothea virgata (Hook.f.) Paul G.Wilson, Nuytsia 12: 260 (1998) | 1:107 |
| | Zieria arborescens Sims subsp. arborescens , Bot. Mag. 34: t.1395 (1811) | 1:103 |
| | <i>Zieria cytisoides</i> Sm. sensu Curtis & Morris (1975) = Zieria littoralis | |
| | Zieria littoralis J.A.Armstr., Austral. Syst. Bot. 15: 389 (2002) | 1:103 |

Zieria smithii Jacks. sensu Bentham (1863) = **Zieria arborescens** subsp. **arborescens** (misapplied in Tasmania)

Zieria smithii Jacks. var. *macrophylla* (Bonpl.) Benth. = **Zieria arborescens** subsp. **arborescens**

Zieria veronicea (F.Muell.) Benth. subsp. **veronicea**, Fl. Austral. 1: 305 (1863) 1:103

SALICACEAE

FTO 80

- i **Populus alba** L., Sp. Pl. 2: 1034 (1753)
- i * *Salix alba* L. var. *vitellina* (L.) Stokes, Bot. Mat. Med. 4: 506 (1812)
- Salix alba* × *fragilis* sensu Curtis (1967) = **Salix × fragilis** nothovar. **fragilis** 3:648
- Salix atrocinerea* Brot. sensu Curtis (1967) = **Salix × reichardtii** 3:649
- Salix babylonica* L. sensu Curtis (1967) = *Salix pendulina* nothovar. *pendulina* 3:648
- i * *Salix × calodendron* Wimm., Sal. Eur. (Wimm.): 187 (1866)
- i **Salix cinerea** L. subsp. **cinerea**, Sp. Pl. 2: 1021 (1753) 3:649
- i **Salix cinerea** L. subsp. **oleifolia** (Sm.) Macreight, Man. Brit. Bot. (Macreight): 212 (1837) 3:649
- i **Salix × fragilis** L. nothovar. **fragilis**, Sp. Pl. 2: 1017 (1753) 3:648
- i * *Salix matsudana* Koidz. cv. *tortuosa*, Bot. Mag. (Tokyo) 29: 312 (1915)
- Salix × pendulina* Wender. nothovar. *pendulina* previously listed as naturalised but insufficient evidence exists to support this
- i # **Salix purpurea** L., Sp. Pl. 2: 1017 (1753)
- i **Salix × reichardtii** A.Kern., Verh. Zool.-Bot. Ges. Wien 10: 249 (1860) 3:649
- i * *Salix × rubens* Schrank, Baier. Fl. 1: 226 (1789) 3:648
- i * *Salix × sepulcralis* Simonk. nothovar. *chrysocoma* (Dode) Meikle, Watsonia 15: 274 (1985)

SANTALACEAE

FTO 91

- Exocarpos cupressiformis** Labill., Voy. Rech. Perouse 1: 156, t.14 (1800) 3:628
- e **Exocarpos humifusus** R.Br., Prodr. Fl. Nov. Holland. 356 (1810) 3:629
- Exocarpos nanus** Hook.f., London J. Bot. 6: 281 (1847) 3:629
- Exocarpos strictus** R.Br., Prodr. Fl. Nov. Holland. 357 (1810) 3:628
- Exocarpos syrticola** (F.Muell. ex Miq.) Stauffer, Mitt. Bot. Mus. Univ. Zurich 213: 173 (1959) 3:628
- Leptomeria billardierei* R.Br. = **Leptomeria drupacea** 3:626
- Leptomeria drupacea** (Labill.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 632 (1917) 3:626
- e **Leptomeria glomerata** F.Muell. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 370 (1859) 3:627
- x **Thesium australe** R.Br., Prodr. Fl. Nov. Holland. 353 (1810) 3:626
- Thesium drupaceum* Labill. = **Leptomeria drupacea**

SAPINDACEAE

FTO 86

- i **Acer pseudoplatanus** L., Sp. Pl. 2: 1054 (1753)
- Dodonaea ericifolia* G.Don = **Dodonaea filiformis** 1:122
- e **Dodonaea filiformis** Link, Enum. Hort. Berol. Alt. 1: 381 (1821) 1:122
- Dodonaea viscosa** Jacq. subsp. **spatulata** (Sm.) J.G.West, Brunonia 7: 43 (1984) 1:122

SCROPHULARIACEAE

- i **Buddleja davidii** Franch., Nouv. Arch. Mus. Hist. Nat. Paris, Ser. 2, 10: 65 (1888)
- Celsia cretica* L. = **Verbascum creticum** 3:513
- Myoporum adscendens* R.Br. = **Myoporum insulare**
- Myoporum insulare** R.Br., Prodr. Fl. Nov. Holland. 516 (1810) 3:540
- Myoporum parvifolium** R.Br., Prodr. Fl. Nov. Holland. 516 (1810) 3:540

| | | |
|-----|--|----------------|
| | Myoporum serratum R.Br. nom. illeg., sensu Bentham (1870) = Myoporum insulare (misapplied in Tasmania) | |
| | Myoporum serratum R.Br. var. obovatum Benth. = Myoporum insulare | |
| | Myoporum tasmanicum A.DC. = Myoporum insulare | |
| | Scrophularia aquatica L. sensu Curtis (1967) = Scrophularia auriculata | 3:515 |
| i | Scrophularia auriculata L., Sp. Pl. 2: 620 (1753) | 3:515 |
| i | Verbascum blattaria L., Sp. Pl. 1: 178 (1753) | 3:512 |
| i | Verbascum creticum (L.) Cav., Elench. Pl. Horti Matr. 39 (1803) | 3:513 |
| i | Verbascum thapsus L., Sp. Pl. 1: 177 (1753) | 3:512 |
| i | Verbascum virgatum Stokes, Bot. Arr. Brit. Pl., ed. 2: 227 (1787) | 3:512 |
| | SOLANACEAE | FTO 112 |
| | Anthocercis tasmanica (Miers) Hook.f. = Cyphanthera tasmanica | 3:509 |
| e | Cyphanthera tasmanica Miers, Ann. Mag. Nat. Hist., Ser. 2, 11: 377 (1853) | 3:509 |
| i | Datura ferox L., Amoen. Acad., Linnaeus ed. 3: 403 (1756) | 3:508 |
| i | Datura stramonium L., Sp. Pl. 1: 179 (1753) | 3:508 |
| | Hyoscyamus albus L. previously listed as naturalised but insufficient evidence exists to support this | 3:509 |
| | Hyoscyamus niger L. sensu Curtis (1967) = Hyoscyamus albus | 3:509 |
| i | Lycium barbarum L., Sp. Pl. 1: 192 (1753) | |
| i | Lycium ferocissimum Miers, Ann. Mag. Nat. Hist., Ser.2, 14: 187 (1854) | 3:507 |
| i t | Nicotiana rustica L., Sp. Pl. 1: 180 (1753) | |
| | Nicotiana sylvestris Speg. previously listed as naturalised but insufficient evidence exists to support this | |
| i | Physalis peruviana L., Sp. Pl., ed. 2, 2: 1670 (1763) | |
| i | Salpichroa organifolia (Lam.) Thell., Mem. Soc. Sci. Nat. Math. Cherbourg 38: 452 (1912) | 3:506 |
| | Solanum americanum Mill. snasu Buchanan (2004) = Solanum nodiflorum (misapplied in Tasmania) | |
| | Solanum aviculare G.Forst. sensu Bentham (1868) = Solanum laciniatum (misapplied in Tasmania) | |
| i t | Solanum dulcamara L., Sp. Pl. 1: 185 (1753) | 3:504 |
| i | Solanum furcatum Dunal ex Poir., Encycl. (Lamarck) Suppl. 3: 750 (1814) | 3:504 |
| | Solanum laciniatum Aiton, Hortus Kew. (W.Aiton) 1: 247 (1789) | 3:505 |
| i | Solanum marginatum L.f., Suppl. Pl. 147 (1782) | 3:506 |
| i | Solanum mauritianum Scop., Delic. Fl. Faun. Insubr. 3: 16, t. 8 (1788) | |
| i | Solanum nigrum L., Sp. Pl. 1: 186 (1753) | 3:504 |
| | Solanum nitidibaccatum Bitter = Solanum physalifolium var. nitidibaccatum | 3:505 |
| i * | Solanum nodiflorum Jacq., Collectanea [Jacquin] 2: 288 (1789) | |
| | Solanum opacum A.Braun & Bouché, Index Seminum Hort. Berol. 8: 18 no.38 (1853) | |
| i | Solanum physalifolium Rusby var. nitidibaccatum (Bitter) Edmonds, Bot. J. Linn. Soc. 92: 27 (1986) | 3:505 |
| i | Solanum pseudocapsicum L., Sp. Pl. 1: 184 (1753) | 3:506 |
| | Solanum sarrachoides Sendtn. = Solanum physalifolium var. nitidibaccatum | |
| i | Solanum triflorum Nutt., Gen. N. Amer. Pl. [Nuttall] 1: 128 (1818) | |
| | Solanum vescum F.Muell., Trans. & Proc. Victorian Inst. Advancem. Sci. 1: 69 (1855) | 3:506 |
| | Solanum vescum F.Muell. var. davidii Geras. (a cultivar name.) | |
| | STYLIDIACEAE | FTO 127 |
| | Candollea armeria Labill. = Stylidium armeria | |
| | Candollea serrulata Labill. = Stylidium graminifolium | |

| | | |
|-----|--|----------------|
| e | Forstera bellidifolia Hook., Icon. Pl. 9: t.851 (1851) | 2:395 |
| x | Levenhookia dubia Sond., Pl. Preiss. [J.G.C.Lehman] 1: 392 (1845) | 2:395 |
| t | Phyllachne colensoi (Hook.f.) Berggr., Minneskr. Kongl. Fysiogr. Sällsk. Lund 8: 11, t.3 figs.1-27 (1877) | 2:396 |
| | Stylidium armeria (Labill.) Labill. subsp. armeria , Nov. Holl. Pl. 2: 66 (1806) | |
| | Stylidium beagleholei J.H.Willis, Muellera 1: 153 (1967) | 2:394 |
| | <i>Stylidium brachyphyllum</i> Sond. sensu Curtis (1963) = Stylidium beagleholei | 2:394 |
| | Stylidium despectum R.Br., Prodr. Fl. Nov. Holland. 571 (1810) | 2:394 |
| | <i>Stylidium dilatatum</i> W.D.Jacks. & R.J.E.Wiltshire = Stylidium armeria subsp. armeria | |
| | Stylidium graminifolium Sw., Sp. Pl., ed. 4 [Willdenow], 4: 146 (1805) | 2:393 |
| | <i>Stylidium inundatum</i> R.Br. sensu Buchanan (2005) = Stylidium beagleholei | 2:394 |
| | <i>Stylidium melastachys</i> R.Br. = Stylidium armeria | |
| | Stylidium perpusillum Hook.f., London J. Bot. 6: 266 (1847) | 2:394 |
| | <i>Stylidium umbellatum</i> (Labill.) Labill. sensu Brown (1810) a name of uncertain application | |
| | TETRACARPAEACEAE | FTO 52 |
| e | Tetracarpaea tasmannica Hook., Icon. Pl. 3: t.264 (1840) | 1:182 |
| | THEOPHRASTACEAE | FTO 103 |
| | <i>Samolus littoralis</i> R.Br. nom. illeg. = Samolus repens | |
| | Samolus repens (J.R.Forst. & G.Forst.) Pers. var. repens , Syn. Pl. (Persoon) 1: 171 (1805) | 3:468 |
| | <i>Sheffieldia incana</i> Labill. = Samolus repens | |
| | THYMELAEACEAE | FTO 89 |
| i t | Daphne laureola L., Sp. Pl. 1: 356 (1753) | |
| | <i>Drapetes tasmanicus</i> Hook.f. = Kelleria dieffenbachii | 3:625 |
| | Kelleria dieffenbachii (Hook.) Endl., Gen. Pl. [Endlicher] Suppl. 4(2): 61 (1848) | 3:625 |
| | Pimelea axiflora F.Muell. ex Meisn. subsp. axiflora , Linnaea 26: 345 (1854) | 3:622 |
| | <i>Pimelea cernua</i> R.Br. = Pimelea linifolia | |
| e | Pimelea cinerea R.Br., Prodr. Fl. Nov. Holland. 361 (1810) | 3:619 |
| | Pimelea curviflora R.Br., Prodr. Fl. Nov. Holland. 362 (1810) | 3:624 |
| | <i>Pimelea curviflora</i> R.Br. var. <i>gracilis</i> (R.Br.) Threlfall = Pimelea curviflora | 3:624 |
| | <i>Pimelea curviflora</i> R.Br. var. <i>sericea</i> Benth. = Pimelea curviflora | 3:624 |
| | Pimelea drupacea Labill., Nov. Holl. Pl. 1: 10 (1805) | 3:622 |
| e | Pimelea filiformis Hook.f., London J. Bot. 6: 280 (1847) | 3:622 |
| | Pimelea flava R.Br. subsp. flava , Prodr. Fl. Nov. Holland. 361 (1810) | 3:623 |
| | <i>Pimelea</i> sp. Freycinet = Pimelea leiophylla (A.M.Buchanan 15902) Tas Herbarium | |
| | Pimelea glauca R.Br., Prodr. Fl. Nov. Holland. 360 (1810) | 3:619 |
| | <i>Pimelea gracilis</i> R.Br. = Pimelea curviflora | |
| | <i>Pimelea gunnii</i> Hook.f. = Pimelea cinerea | |
| | Pimelea humilis R.Br., Prodr. Fl. Nov. Holland. 361 (1810) | 3:620 |
| | <i>Pimelea incana</i> R.Br. = Pimelea nivea | |
| e | Pimelea leiophylla A.M.Gray & M.Baker, Muellera 35: 16 (2016) | |
| | Pimelea ligustrina Labill. subsp. ligustrina , Nov. Holl. Pl. 1: 9, f.3 (1805) | |
| | <i>Pimelea lindleyana</i> Meisn. = Pimelea linifolia | 3:620 |
| | Pimelea linifolia Sm., Spec. Bot. New Holland 31, t.11 (1793) | 3:620 |
| | <i>Pimelea linifolia</i> Sm. subsp. <i>linifolia</i> = Pimelea linifolia | |

| | | |
|-----|--|----------------|
| | <i>Pimelea linifolia</i> Sm. subsp. <i>linoides</i> (A.Cunn.) Threlfall = <i>Pimelea linifolia</i> | |
| | <i>Pimelea micrantha</i> F.Muell. ex Meisn., <i>Linnaea</i> 26: 351 (1854) | 3:624 |
| e | <i>Pimelea milliganii</i> Meisn., <i>Prodr.</i> [A. P. de Candolle] 14: 509 (1857) | 3:619 |
| e | <i>Pimelea nivea</i> Labill., <i>Nov. Holl. Pl.</i> 1: 10 (1805) | 3:621 |
| | <i>Pimelea pauciflora</i> R.Br., <i>Prodr. Fl. Nov. Holland.</i> 360 (1810) | 3:622 |
| | <i>Pimelea phyllicoides</i> Meisn. sensu Rye (1990) probably recorded in error | |
| e | <i>Pimelea pygmaea</i> F.Muell. & C.Stuart ex Meisn., <i>Linnaea</i> 26: 346 (1853) | 3:623 |
| e | <i>Pimelea sericea</i> R.Br., <i>Prodr. Fl. Nov. Holland.</i> 361 (1810) | 3:621 |
| | <i>Pimelea serpyllifolia</i> R.Br. subsp. <i>serpyllifolia</i> , <i>Prodr. Fl. Nov. Holland.</i> 360 (1810) | 3:623 |
| | <i>Pimelea spatulata</i> Labill. = <i>Pimelea linifolia</i> | |
| | <i>Pimelea stricta</i> Meisn. sensu Curtis (1967) = <i>Pimelea linifolia</i> | 3:624 |
| | <i>Pimelea</i> sp. Tunbridge (A.Moscal 9026) Tas Herbarium | |
| | TROPAEOLACEAE | FTO 82 |
| i | <i>Tropaeolum majus</i> L., <i>Sp. Pl.</i> 1: 345 (1753) | |
| | ULMACEAE | FTO 63 |
| i | <i>Ulmus</i> × <i>hollandica</i> Mill., <i>Gard. Dict.</i> , ed. 8, 5 (1768) | 3:642 |
| | URTICACEAE | FTO 64 |
| | <i>Australina muelleri</i> Wedd. = <i>Australina pusilla</i> subsp. <i>muelleri</i> | 3:641 |
| | <i>Australina pusilla</i> (Poir.) Gaudich. subsp. <i>muelleri</i> (Wedd.) Friis & Wilmot-Dear, <i>Nordic. J. Bot.</i> 7: 126 (1987) | 3:641 |
| | <i>Australina pusilla</i> (Poir.) Gaudich. subsp. <i>pusilla</i> , <i>Voy. Bonite, Bot.</i> 3: t.114A (1852) | 3:641 |
| | <i>Helxine soleirolii</i> Req. sensu Curtis (1967) = <i>Soleirolia soleirolii</i> | 3:640 |
| | <i>Parietaria cardiostegia</i> Greuter, <i>Fl. Australia</i> 3: 190 (1989) | |
| | <i>Parietaria debilis</i> G.Forst., <i>Fl. Ins. Austr.</i> 73 (1786) | 3:640 |
| i * | <i>Parietaria judaica</i> L., <i>Sp. Pl.</i> , ed. 2, 2: 1492 (1763) | |
| i | <i>Soleirolia soleirolii</i> (Req.) Dandy, <i>Feddes Repert.</i> 70: 4 (1965) | 3:640 |
| | <i>Urtica incisa</i> Poir., <i>Encycl. (Lamarck) Suppl.</i> 4: 224 (1816) | 3:639 |
| i | <i>Urtica urens</i> L., <i>Sp. Pl.</i> 2: 984 (1753) | 3:639 |
| | VALERIANACEAE | FTO 135 |
| n i | <i>Centranthus ruber</i> (L.) DC. subsp. <i>ruber</i> , <i>Fl. Franc. (DC. & Lamarck)</i> , ed. 3, 4(2): 239 (1805) | 2:278 |
| i | <i>Valerianella eriocarpa</i> Desv., <i>J. Bot. (Desvaux)</i> 2: 314, t.11 fig.2 (1809) | 2:278 |
| i | <i>Valerianella locusta</i> (L.) Laterr., <i>Fl. Bordel.</i> , ed. 2, 93 (1821) | |
| i t | <i>Valerianella rimosa</i> Bastard, <i>J. Bot. Agric.</i> 3: 20 (1814) | |
| | VERBENACEAE | |
| i | <i>Verbena bonariensis</i> L., <i>Sp. Pl.</i> 1: 20 (1753) | |
| | <i>Verbena incompta</i> P.W.Michael sensu Buchanan (1999) = <i>Verbena bonariensis</i> (misapplied in Tasmania) | |
| i | <i>Verbena officinalis</i> L., <i>Sp. Pl.</i> 1: 20 (1753) | 3:541 |
| | VIOLACEAE | FTO 79 |
| | <i>Hymenanthera angustifolia</i> DC. = <i>Melicytus angustifolius</i> | |
| | <i>Hymenanthera banksii</i> F.Muell. nom. illeg., nom. superfl. = <i>Melicytus dentatus</i> | |
| | <i>Hymenanthera dentata</i> DC. sensu Curtis & Morris (1975) = <i>Melicytus dentatus</i> and <i>M. angustifolius</i> | 1:55 |

- e **Melicytus angustifolius** (DC.) Garn.-Jones subsp. **angustifolius**, New Zealand J. Bot. 25: 127 (1987)
Melicytus angustifolius (DC.) Garn.-Jones subsp. **divaricatus** Stajsic & R.Douglas, Austral. Syst. Bot. 27: 320 (2014)
- Melicytus dentatus** (DC.) Molloy & Mabb., Curtis's Bot. Mag. 17: 234 (2000) 1:55
- i **Viola arvensis** Murray, Prodr. Stirp. Gott. 73 (1770) 1:55
Viola betonicifolia Sm. subsp. **betonicifolia**, Cycl. (Rees) 37: no.7 (1817) 1:54
Viola calejana G.Don, Gen. Hist. 1: 329 (1831) 1:55
Viola cleistogamoides (L.G.Adams) Seppelt, Fl. S. Australia, ed. 4, 2: 870 (1986)
- t **Viola cunninghamii** Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 16 (1852) 1:54
Viola curtisiae (L.G.Adams) K.R.Thiele, Muelleria 36: 109 (2018)
Viola fuscoviolacea (L.G.Adams) T.A.James, Muelleria 9: 35 (1996)
Viola hederacea Labill. subsp. **cleistogamoides** L.G.Adams = **Viola cleistogamoides**
Viola hederacea Labill. subsp. **curtisiae** L.G.Adams = **Viola curtisiae**
Viola hederacea Labill. subsp. **fuscoviolacea** L.G.Adams = **Viola fuscoviolacea**
Viola hederacea Labill. subsp. **hederacea**, Nov. Holl. Pl. 1: 66, t.91 (1805) 1:54
- i **Viola odorata** L., Sp. Pl. 2: 934 (1753)
- e **Viola serpentinicola** de Salas, Muelleria 36: 113 (2018)
- n *Viola sieberiana* Spreng. sensu de Salas & Baker (2017) and earlier = **Viola hederacea** subsp. **hederacea** (misapplied in Tasmania) 1:54

ZYGOPHYLLACEAE

FTO 58

- Roepera billardierei* (DC.) G.Don = **Zygophyllum billardierei**
Roepera latifolia Hook.f. sensu Hooker (1860) recorded in error
Zygophyllum apiculatum F.Muell. sensu Curtis & Morris (1975) recorded in error 1:93
Zygophyllum billardierei DC., Prodr. [A. P. de Candolle] 1: 705 (1824) 1:93

MONOCOTS

AGAPANTHACEAE

- i **Agapanthus praecox** Willd. subsp. **orientalis** (F.M.Leight.) F.M.Leight., J. S. African Bot. 4: 21 (1965)

ALISMATACEAE

- i **Alisma lanceolatum** With., Arr. Brit. Pl., ed. 3, 2: 362 (1796)
- i **Alisma plantago-aquatica** L., Sp. Pl. 1: 342 (1753) 4b:4
Damasonium australe Salisb. = **Damasonium minus** 4b:6
Damasonium minus (R.Br.) Buchenau, Abh. Naturwiss. Vereins Bremen 2: 20 (1869) 4b:6

ALLIACEAE

- i **Allium ampeloprasum** L., Sp. Pl. 1: 294 (1753)
- i **Allium neapolitanum** Cirillo, Pl. Rar. Neap. 1: 13, t.4 (1788) 4b:392
- i **Allium triquetrum** L., Sp. Pl. 1: 300 (1753) 4b:392
- i **Allium vineale** L., Sp. Pl. 1: 299 (1753) 4b:392
- i **Nothoscordum borbonicum** Kunth, Enum. Pl. [Kunth] 4: 463 (1843) 4b:394
Nothoscordum gracile (Aiton) Stearn sensu Curtis & Morris (1994) = **Nothoscordum borbonicum** 4b:394

ALSTROEMERIACEAE

- i # **Alstroemeria aurea** Graham, Edinburgh New Philos. J. 15: 181 (1833)

AMARYLLIDACEAE

- i **Leucojum aestivum** L., Syst. Nat., ed. 10. 2: 975 (1759)
- i **Narcissus pseudonarcissus** L., Sp. Pl. 1: 289 (1753) 4b:394
- i **Narcissus tazetta** L., Sp. Pl. 1: 290 (1753)

APONOGETONACEAE

- i **Aponogeton distachyos** L.f., Suppl. Pl. 215 (1782) 4b:13

ARACEAE

- Landoltia punctata** (G.Mey.) Les & D.J.Crawford, Novon 9: 532 (1999)
- Lemna disperma** Hegelm., Bot. Zeitung (Berlin) 29: 655 (1871) 4b:32
- Lemna minor L. sensu Rodway (1903) = **Lemna disperma** 4b:32
- Lemna trisulca** L., Sp. Pl. 2: 970 (1753) 4b:32
- Spirodela polyrhiza (L.) Schleid. probably misapplied to **Landoltia punctata**
- Wolffia arrhiza (L.) Horkel ex Wimm. sensu Willis (1973) = **Wolffia australiana** 4b:32
- Wolffia australiana** (Benth.) Hartog & Plas, Blumea 20: 151 (1972) 4b:32
- i **Zantedeschia aethiopica** (L.) Spreng., Sys. Veg. (ed. 16) [Sprengel] 3: 765 (1826) 4b:30

ASPARAGACEAE

- i **Agave americana** L., Sp. Pl. 1: 323 (1753)
- i **Asparagus asparagoides** (L.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 3: 414 (1914)
- i **Asparagus officinalis** L., Sp. Pl. 1: 313 (1753) 4b:370
- i **Asparagus scandens** Thunb., Prod. Pl. Cap. 66 (1794)
- i# **Furcraea foetida** (L.) Haw., Syn. Pl. Succ. 73 (1812)
- Lomandra glauca (R.Br.) Ewart sensu Willis (1973) = **Lomandra nana** 4b:422
- Lomandra longifolia** Labill., Nov. Holl. Pl. 1: 92, t.119 (1805) 4b:422
- Lomandra nana** (A.T.Lee) A.T.Lee, Fl. Australia 46: 223 (1986) 4b:422
- i **Muscari armeniacum** Leichtlin ex Baker, Gard. Chron. 798 (1878)
- Myrsiphyllum asparagoides (L.) Willd. = **Asparagus asparagoides**
- Myrsiphyllum scandens (Thunb.) Oberm. = **Asparagus scandens**
- Xerotes glauca R.Br. sensu Rodway (1903) = **Lomandra nana** 4b:422
- Xerotes longifolia (Labill.) R.Br. = **Lomandra longifolia** 4b:422

ASPHODELACEAE

- i **Aloe maculata** All., Auct. Syn. Meth. Stirp. Hort. Regii Taur. 13 (1773)
- Aloe saponaria (Aiton) Haw. = **Aloe maculata**
- Anthericum semibarbatum R.Br. = **Bulbine semibarbata**
- i **Asphodelus fistulosus** L., Sp. Pl. 1: 309 (1753) 4b:380
- Bulbine bulbosa** (R.Br.) Haw., Revis. Pl. Succ. 33 (1821) 4b:383
- Bulbine crassa** D.I.Morris & Duretto, Muelleria 22: 93 (2006)
- Bulbine glauca** (Raf.) E.M.Watson, Fl. Australia 45: 469 (1987) 4b:382
- Bulbine semibarbata** (R.Br.) Haw., Revis. Pl. Succ. 33 (1821) 4b:382
- i **Kniphofia uvaria** (L.) Oken, Allg. Naturgesch. 3: 566 (1841)

ASTELIACEAE

- e **Astelia alpina** R.Br. var. **alpina**, Prodr. Fl. Nov. Holland. 291 (1810) 4b:370
- e **Astelia sp. Quartzite (A.M.Buchanan 16828) Tas Herbarium**
- Astelia stylosa F.Muell. ex Hook.f. = **Milligania stylosa**

- e **Milligania densiflora** Hook.f., Hooker's J. Bot. Kew Gard. Misc. 5: 298 (1853) 4b:374
 e **Milligania johnstonii** F.Muell. ex Benth., Fl. Austral. 7: 26 (1878) 4b:371
 e **Milligania lindoniana** Rodway ex W.M.Curtis, Records of the Queen Victoria Museum 45: 1 (1972) 4b:372
 e **Milligania longifolia** Hook.f., Hooker's J. Bot. Kew Gard. Misc. 5: 297 (1853) 4b:373
 e **Milligania stylosa** (F.Muell. ex Hook.f.) F.Muell. ex Benth., Fl. Austral. 7: 27 (1878) 4b:373

BLANDFORDIACEAE

Blandfordia grandiflora Herb. var. backhousii (Lindl.) Hook.f. = **Blandfordia punicea**

Blandfordia marginata Herb. = **Blandfordia punicea** 4b:374

- e **Blandfordia punicea** (Labill.) Sweet, Hort. Brit. [Sweet], ed 2: 517 (1830) 4b:374

BURMANNIACEAE

FTO 16

Thismia rodwayi F.Muell., Vict. Naturalist 7: 115 (1890) 4b:423

CAMPYNEMATACEAE

- e **Campynema lineare** Labill., Nov. Holl. Pl. 1: 93, t.121 (1805) 4b:395

Campynema pygmaeum F.Muell. ex Benth. = **Campynema lineare**

CENTROLEPIDACEAE

Alepyrum monogynum Hook.f. = **Centrolepis monogyna**

Alepyrum muelleri Hook.f. = **Centrolepis glabra**

Alepyrum muscoides Hook.f. = **Centrolepis muscoides**

Alepyrum polygynum R.Br. = **Centrolepis polygyna**

Aphelia gracilis Sond., Linnaea 28: 227 (1856) 4b:49

Aphelia gunnii Hook.f. = **Aphelia gracilis**

Aphelia pumilio F.Muell. ex Sond., Linnaea 28: 226 (1856) 4b:49

Centrolepis aristata (R.Br.) Poir., Syst. Veg., ed. 15 bis [Roemer & Schultes] 1: 44 (1817) 4b:54

Centrolepis fascicularis Labill., Nov. Holl. Pl. 1: 7, t.1 (1804) 4b:55

Centrolepis glabra (F.Muell. ex Sond.) Hieron., Abh. Naturf. Ges. Halle 12: 209 (1873) 4b:52

- e **Centrolepis monogyna** (Hook.f.) Benth., Fl. Austral. 7: 205 (1878) 4b:53

Centrolepis monogyna (Hook.f.) Benth. subsp. paludicola (W.M.Curtis) D.A.Cooke = **Centrolepis monogyna** 4b:53

- e **Centrolepis muscoides** (Hook.f.) Hieron., Abh. Naturf. Ges. Halle 12: 209 (1873) 4b:53

Centrolepis paludicola W.M.Curtis = **Centrolepis monogyna** 4b:53

- e **Centrolepis pedderensis** W.M.Curtis, Brunonia 7: 299 (1984) 4b:53

Centrolepis polygyna (R.Br.) Hieron., Abh. Naturf. Ges. Halle 12: 210 (1873) 4b:51

Centrolepis pulvinata (R.Br.) Poir. = **Centrolepis strigosa** subsp. **pulvinata** 4b:55

- e **Centrolepis strigosa** (R.Br.) Poir. subsp. **pulvinata** (R.Br.) D.A.Cooke, J. Adelaide Bot. Gard. 15: 28 (1992) 4b:55

Centrolepis strigosa (R.Br.) Poir. subsp. **strigosa**, Syst. Veg., ed. 15 bis [Roemer & Schultes] 1: 43 (1817) 4b:55

Centrolepis tenuior (R.Br.) Poir. sensu Hooker (1860) = **Centrolepis strigosa** subsp. **strigosa**

Devauxia billardierei R.Br. = **Centrolepis fascicularis**

Devauxia pulvinata R.Br. = **Centrolepis strigosa** subsp. **pulvinata**

Devauxia tenuior R.Br. = **Centrolepis strigosa** subsp. **strigosa**

- e **Gaimardia amblyphylla** W.M.Curtis, Brunonia 7: 299 (1984) 4b:57

- e **Gaimardia fitzgeraldii** F.Muell. & Rodway, Pap. & Proc. Roy. Soc. Tasmania 1894-5: 55 (1896) 4b:57

| | | |
|----------------------|--|--------|
| t | Gaimardia setacea Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 267 (1853) | 4b:57 |
| COLCHICACEAE | | |
| | Anguillaria dioica R.Br. pro spec. Tasm. = Wurmbea uniflora, W. dioica & W. latifolia | |
| | Anguillaria uniflora R.Br. = Wurmbea uniflora | |
| | Burchardia umbellata R.Br., Prodr. Fl. Nov. Holland. 273 (1810) | 4b:396 |
| | Wurmbea biglandulosa (R.Br.) T.D.Macfarl. subsp. biglandulosa , Brunonia 3: 191 (1980) | |
| | Wurmbea dioica (R.Br.) F.Muell. subsp. dioica , Fragm. (Mueller) 10: 119 (1877) | 4b:396 |
| | Wurmbea latifolia T.D.Macfarl. subsp. vanessae R.J.Bates, J. Adelaide Bot. Gard. 16: 48 (1995) | 4b:398 |
| | Wurmbea uniflora (R.Br.) T.D.Macfarl., Brunonia 3: 194 (1980) | 4b:398 |
| COMMELINACEAE | | |
| | Tradescantia albiflora Kunth = Tradescantia fluminensis | 4b:36 |
| i | Tradescantia fluminensis Vell., Fl. Flumin. 140 (1829) | 4b:36 |
| CYMODOCEACEAE | | |
| | Amphibolis antarctica (Labill.) Sond. & Asch. ex Asch., Linnaea 35: 164 (1867) | 4b:27 |
| | Cymodocea antarctica (Labill.) Endl. ex Kunth = Amphibolis antarctica | 4b:27 |
| CYPERACEAE | | |
| | Baumea acuta (Labill.) Palla, Allg. Bot. Z. Syst. 8: 69 (1902) | 4b:131 |
| | Baumea arthropphylla (Nees) Boeckeler, Linnaea 38: 242 (1874) | 4b:129 |
| | Baumea articulata (R.Br.) S.T.Blake, Contr. Queensland Herb. 8: 28 (1969) | 4b:128 |
| | Baumea gunnii (Hook.f.) S.T.Blake, Contr. Queensland Herb. 8: 27 (1969) | 4b:130 |
| | Baumea juncea (R.Br.) Palla, Allg. Bot. Z. Syst. 15: 113 (1909) | 4b:129 |
| ?i | Baumea planifolia (Benth.) K.L.Wilson, Telopea 5: 589 (1994) | |
| | Baumea rubiginosa (Spreng.) Boeckeler, Linnaea 38: 241 (1874) | 4b:129 |
| | Baumea tetragona (Labill.) S.T.Blake, Contr. Queensland Herb. 8: 30 (1969) | 4b:131 |
| | Bolboschoenus caldwellii (V.J.Cook) Soják, Cas. Nár. Mus., Odd. Prír. 141: 62 (1972) | 4b:92 |
| | Bolboschoenus medianus (V.J.Cook) Soják, Cas. Nár. Mus., Odd. Prír. 141: 63 (1972) | 4b:93 |
| | Carex acicularis Boott = Carex archeri | 4b:148 |
| | Carex albula Allan sensu Curtis & Morris (1994) recorded in error | 4b:164 |
| e | Carex sp. Algonkian Rivulet (S.J.Jarman HO110282) Tas Herbarium | 4b:163 |
| | Carex appressa R.Br., Prodr. Fl. Nov. Holland. 242 (1810) | 4b:152 |
| | Carex appressa R.Br. f. minor Kük. = Carex appressa | 4b:152 |
| | Carex appressa R.Br. var. virgata (Boott) Kük. = Carex appressa | 4b:153 |
| | Carex archeri Boott, Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 98 (1858) | 4b:148 |
| e | Carex barbata Boott, Ill. Gen. Carex 1: 68, t.187 (1858) | 4b:160 |
| | Carex bichenoviana Boott, Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 101 (1858) | 4b:167 |
| | Carex breviculmis R.Br., Prodr. Fl. Nov. Holland. 242 (1810) | 4b:158 |
| i* | Carex buxbaumii Wahlenb., Kongl. Vetensk. Acad. Nya Handl. 24: 163 (1803) | |
| | Carex caespitosa L. sensu Brown (1810) = Carex gaudichaudiana | |
| | Carex canescens L. sensu Hooker (1860) = Carex curta (misapplied in Tasmania) | 4b:155 |
| | Carex capillacea Boott, Ill. Gen. Carex 1: 44, t.110 (1858) | 4b:148 |
| e | Carex cataractae R.Br., Prodr. Fl. Nov. Holland. 242 (1810) | 4b:162 |
| | Carex cephalotes F.Muell., Trans. Philos. Soc. Victoria 1: 110 (1855) | 4b:149 |
| | Carex chlorantha R.Br., Prodr. Fl. Nov. Holland. 242 (1810) | 4b:149 |

| | | |
|-------|---|--------|
| | Carex aff. cirrhosa Berggr. sensu Curtis & Morris (1994) = Carex sp. Algonkian Rivulet (S.J.Jarman HO110282) Tas Herbarium | 4b:163 |
| | Carex curta Gooden., Trans. Linn. Soc. London 2: 145 (1794) | 4b:155 |
| i t | Carex demissa Hornem., Fl. Dan., 7(21): 4, t.1342 (1806) | 4b:161 |
| | Carex aff. diandra Schrank sensu Curtis & Morris (1994) = Carex sp. Western Tasmania (A.Moscal 7489) Tas Herbarium | 4b:150 |
| i | Carex divisa Huds., Fl. Angl. (Hudson): 348 (1762) | 4b:149 |
| i | Carex divulsa Stokes, Bot. Arr. Brit. Pl., ed. 2. 2: 1035 (1787) | 4b:150 |
| | Carex fascicularis Sol. ex Boott, Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 283 (1853) | 4b:166 |
| i | Carex flacca Schreb., Spicil. Fl. Lips. 178 (1771) | 4b:157 |
| i t | Carex flagellifera Colenso, Trans. & Proc. New Zealand Inst. 16: 342 (1884) | 4b:163 |
| | Carex flava L. sensu Rodway (1903) = Carex cataractae, C. demissa, C. flaviformis & C. lepidocarpa | |
| t | Carex flaviformis Nelmes, Kew Bull. 10: 84 (1955) | 4b:162 |
| | Carex gaudichaudiana Kunth, Enum. Pl. [Kunth] 2: 417 (1837) | 4b:156 |
| | Carex gunniana Boott, Proc. Linn. Soc. London 1: 258 (1845) | 4b:160 |
| | Carex hypandra F.Muell. ex Benth., Fl. Austral. 7: 439 (1878) | 4b:157 |
| | Carex inversa R.Br., Prodr. Fl. Nov. Holland. 242 (1810) | 4b:153 |
| | Carex iynx Nelmes, Proc. Linn. Soc. London 155: 279 (1944) | 4b:159 |
| i t | Carex lepidocarpa Tausch, Flora 17: 179 (1834) | 4b:161 |
| i | Carex leporina L., Sp. Pl. 2: 973 (1753) | 4b:155 |
| | Carex littorea Labill. = Carex pumila | |
| | Carex longebrachiata Boeckeler, Linnaea 41: 282 (1877) | 4b:158 |
| | Carex longifolia R.Br. = Carex longebrachiata | |
| | Carex longifolia R.Br. sensu Rodway (1903) = Carex iynx (misapplied in Tasmania) | |
| | Carex ovalis Gooden. = Carex leporina | |
| | Carex paniculata L. sensu Rodway (1903) = Carex appressa (misapplied in Tasmania) | 4b:152 |
| i * t | Carex pilulifera L., Sp. Pl. 2: 976 (1753) | |
| | Carex polyantha F.Muell., Trans. Philos. Soc. Victoria 1: 110 (1855) | 4b:156 |
| | Carex pseudocyperus L. sensu Rodway (1903) = Carex fascicularis (misapplied in Tasmania) | 4b:166 |
| | Carex pumila Thunb. ex Murray, Syst. Veg., ed. 14 (J. A. Murray). 846 (1784) | 4b:166 |
| | Carex raleighii Nelmes, Bull. Misc. Inform. Kew 1939: 310 (1939) | 4b:154 |
| i * | Carex scoparia Schkuhr ex Willd., Sp. Pl., ed. 4 [Willdenow], 4: 230 (1805) | 4b:154 |
| | Carex tasmanica Kük., Bull. Herb. Boissier 4: 59 (1904) | 4b:165 |
| | Carex tereticaulis F.Muell., Fragm. (Mueller) 8: 256 (1874) | 4b:151 |
| | Carex testacea Sol. ex Boott previously listed as naturalised but insufficient evidence exists to support this | 4b:164 |
| | Carex virgata Sol. ex Boott = Carex appressa | 4b:153 |
| | Carex vulgaris Fr. sensu Rodway (1903) = Carex gaudichaudiana (misapplied in Tasmania) | 4b:156 |
| e | Carex sp. Western Tasmania (A.Moscal 7489) Tas Herbarium | 4b:150 |
| | Carpha alpina R.Br., Prodr. Fl. Nov. Holland. 230 (1810) | 4b:87 |
| e | Carpha curvata W.M.Curtis, Brunonia 7: 303 (1985) | 4b:88 |
| e | Carpha rodwayi W.M.Curtis, Brunonia 7: 302 (1985) | 4b:88 |

| | | |
|------|--|--------|
| | Caustis pentandra R.Br., Prodr. Fl. Nov. Holland. 240 (1810) | 4b:137 |
| | Chaetospora axillaris R.Br. = Schoenus maschalinus | |
| | Chaetospora capillacea Hook.f. nom. illeg. = Tetraria capillaris | |
| | Chaetospora imberbis R.Br. = Schoenus apogon | |
| | Chaetospora nitens R.Br. = Schoenus nitens | |
| | Chaetospora tenuissima Hook.f. nom. illeg. = Schoenus lepidosperma | |
| | Chorizandra australis K.L.Wilson, Telopea 5: 594 (1994) | 4b:138 |
| | Chorizandra cymbaria R.Br. sensu Rodway (1903) = Chorizandra australis | 4b:138 |
| | Chorizandra enodis Nees, Pl. Preiss. [J.G.C.Lehman] 2: 73 (1846) | 4b:138 |
| | Cladium acutum (Labill.) Poir. = Baumea acuta | |
| | Cladium filum (Labill.) R.Br. = Gahnia filum | 4b:136 |
| | Cladium glomeratum R.Br. = Baumea rubiginosa | 4b:129 |
| | Cladium gunnii Hook.f. = Baumea gunnii | 4b:130 |
| | Cladium junceum R.Br. = Baumea juncea | 4b:129 |
| | Cladium laxiflorum Hook.f. = Baumea gunnii | |
| | Cladium mariscus (L.) Pohl recorded in error | |
| ?i # | Cladium procerum S.T.Blake, Trans. Roy. Soc. South Australia 67: 57 (1943) | 4b:428 |
| | Cladium schoenoides R.Br. = Baumea acuta | 4b:131 |
| | Cladium tetraquetrum Hook.f. = Baumea tetragona | 4b:131 |
| i | Cyperus congestus Vahl, Enum. Pl. [Vahl] 2: 358 (1805) | 4b:91 |
| i | Cyperus eragrostis Lam., Tabl. Encycl. 1: 146 (1791) | 4b:90 |
| | Cyperus gunnii Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 80, t.140, fig.A (1858) | 4b:91 |
| | Cyperus lucidus R.Br., Prodr. Fl. Nov. Holland. 218 (1810) | 4b:91 |
| | Cyperus sanguineo-fuscus Nees = Cyperus lucidus | |
| ?i # | Cyperus sanguinolentus Vahl, Enum. Pl. [Vahl] 2: 351 (1805) | 4b:90 |
| | Cyperus tenellus L.f. = Isolepis levynsiana | 4b:90 |
| | Eleocharis acicularis (L.) Roem. & Schult. sensu Rodway (1903) = Eleocharis pusilla | 4b:108 |
| | Eleocharis acuta R.Br., Prodr. Fl. Nov. Holland. 224 (1810) | 4b:107 |
| | Eleocharis gracilis R.Br., Prodr. Fl. Nov. Holland. 224 (1810) | 4b:107 |
| | Eleocharis pusilla R.Br., Prodr. Fl. Nov. Holland. 225 (1810) | 4b:108 |
| | Eleocharis sphacelata R.Br., Prodr. Fl. Nov. Holland. 224 (1810) | 4b:107 |
| | Elynanthus capillaceus Benth. = Tetraria capillaris | 4b:132 |
| | Ficinia nodosa (Rottb.) Goetgh., Muasya & D.A.Simpson, Novon 10: 133 (2000) | 4b:105 |
| | Gahnia filum (Labill.) F.Muell., Key to the System of Victorian Plants 1: 456 (1888) | 4b:136 |
| | Gahnia fitzgeraldii F.Muell. ex Rodway = Gahnia microstachya | 4b:134 |
| | Gahnia graminifolia Rodway nom. illeg. = Gahnia rodwayi | 4b:134 |
| | Gahnia grandis (Labill.) S.T.Blake, Contr. Queensland Herb. 8: 33 (1969) | 4b:135 |
| | Gahnia melanocarpa R.Br. sensu Buchanan et al. (1989) = Gahnia grandis (misapplied in Tasmania) | |
| | Gahnia microstachya Benth., Fl. Austral. 7: 414 (1878) | 4b:134 |
| | Gahnia psittacorum Labill. = Gahnia grandis | 4b:135 |
| | Gahnia radula (R.Br.) Benth., Fl. Austral. 7: 417 (1878) | 4b:135 |
| e | Gahnia rodwayi F.Muell. ex Rodway, Pap. & Proc. Roy. Soc. Tasmania 1892: 93 (1893) | 4b:134 |
| | Gahnia sieberiana Kunth, Enum. Pl. [Kunth] 2: 332 (1837) | 4b:135 |

| | | |
|-----|---|--------|
| | Gahnia trifida Labill., Nov. Holl. Pl. 1: 89, t.116 (1805) | 4b:136 |
| | Gymnoschoenus sphaerocephalus (R.Br.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 83, t.142 (1858) | 4b:119 |
| n e | Isolepis alpina Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 86, t.143, fig.B (1858) | 4b:103 |
| | Isolepis aucklandica Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 88 (1844), pl.50 (1845) | 4b:102 |
| | <i>Isolepis cartilaginea</i> R.Br. = Isolepis marginata | |
| | Isolepis cernua (Vahl) Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 2: 106 (1817) | 4b:99 |
| | Isolepis crassiuscula Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 86, t.143, fig.A (1858) | 4b:98 |
| | Isolepis fluitans (L.) R.Br., Prodr. Fl. Nov. Holland. 221 (1810) | 4b:98 |
| | Isolepis habra (Edgar) Soják, Casopsis Narodniho Muzea v Praze 148: 194 (1980) | 4b:101 |
| | Isolepis hookeriana Boeckeler, Flora 41: 418 (1858) | 4b:104 |
| i | Isolepis hystrix (Thunb.) Nees, Linnaea 7: 496 (1832) | |
| | Isolepis inundata R.Br., Prodr. Fl. Nov. Holland. 222 (1810) | 4b:100 |
| | <i>Isolepis lenticularis</i> R.Br. sensu Hooker (1860) = Isolepis fluitans | |
| ?i | Isolepis levynsiana Muasya & D.A.Simpson, Novon 17: 59 (2007) | 4b:90 |
| e | Isolepis limbata W.M.Curtis, Brunonia 7: 300 (1985) | 4b:103 |
| | Isolepis marginata (Thunb.) A.Dietr., Sp. Pl., ed. 6. 1(2): 110 (1832) | 4b:104 |
| | Isolepis montivaga (S.T.Blake) K.L.Wilson, Telopea 2: 168 (1981) | 4b:103 |
| | <i>Isolepis nodosa</i> (Rottb.) R.Br. = Ficinia nodosa | 4b:105 |
| | Isolepis platycarpa (S.T.Blake) Soják, Cas. Nár. Muz. Praze, Rada Prír. 148: 194 (1980) | 4b:99 |
| | Isolepis producta (C.B.Clarke) K.L.Wilson, Telopea 2: 168 (1981) | 4b:98 |
| i | Isolepis prolifera (Rottb.) R.Br., Prodr. Fl. Nov. Holland. 223 (1810) | |
| | <i>Isolepis riparia</i> R.Br. = Isolepis cernua | |
| | <i>Isolepis saviana</i> Schult. nom. illeg. sensu Hooker (1860) a name of uncertain application | |
| i t | Isolepis setacea (L.) R.Br., Prodr. Fl. Nov. Holland. 222 (1810) | 4b:105 |
| | Isolepis stellata (C.B.Clarke) K.L.Wilson, Telopea 2: 169 (1981) | 4b:104 |
| | Isolepis subtilissima Boeckeler, Flora 41: 416 (1858) | 4b:101 |
| e | Isolepis tasmanica (S.T.Blake) K.L.Wilson, Telopea 2: 170 (1981) | 4b:102 |
| | Isolepis wakefieldiana (S.T.Blake) K.L.Wilson, Telopea 2: 170 (1981) | 4b:100 |
| | <i>Lampocarya hexandra</i> R.Br. = Gahnia trifida | |
| | <i>Lepidosperma angustifolia</i> Hook.f. sensu Hooker (1860) = Lepidosperma laterale | |
| | Lepidosperma concavum R.Br., Prodr. Fl. Nov. Holland. 234 (1810) | 4b:126 |
| | Lepidosperma curtisiae K.L.Wilson & D.I.Morris, Telopea 5: 423 (1993) | 4b:123 |
| | <i>Lepidosperma elatior</i> Labill. orth. var. sensu Rodway (1903) = Lepidosperma elatius | 4b:124 |
| | Lepidosperma elatius Labill., Nov. Holl. Pl. 1: 15, t.11 (1805) | 4b:124 |
| | <i>Lepidosperma elatius</i> Labill. var. ensiforme Rodway = Lepidosperma ensiforme | 4b:125 |
| | <i>Lepidosperma elatius</i> Labill. var. oldfieldii (Hook.f.) Rodway = Lepidosperma oldfieldii | 4b:125 |
| | Lepidosperma ensiforme (Rodway) D.I.Morris, The Student's Flora of Tasmania 4b: 425 (1994) | 4b:125 |
| | <i>Lepidosperma falcatum</i> Rodway = Lepidosperma tortuosum | |
| | Lepidosperma filiforme Labill., Nov. Holl. Pl. 1: 17, t.15 (1805) | 4b:121 |
| | Lepidosperma forsythii A.A.Ham., Proc. Linn. Soc. New South Wales 35: 411 (1910) | 4b:122 |
| | Lepidosperma gladiatum Labill., Nov. Holl. Pl. 1: 15, t.12 (1805) | 4b:125 |
| e | Lepidosperma globosum Labill., Nov. Holl. Pl. 1: 16, t.14 (1805) | 4b:123 |

| | | |
|---|---|--------|
| | Lepidosperma gunnii Boeckeler, Linnaea 38: 325 (1874) | 4b:123 |
| e | Lepidosperma inops F.Muell. ex Rodway, Pap. & Proc. Roy. Soc. Tasmania 1892: 93 (1893) | 4b:122 |
| | Lepidosperma laterale R.Br., Prodr. Fl. Nov. Holland. 234 (1810) | 4b:124 |
| | Lepidosperma lineare R.Br. sensu Rodway (1903) = Lepidosperma gunnii | 4b:123 |
| | Lepidosperma lineare R.Br. var. inops (F.Muell. ex Rodway) Rodway = Lepidosperma inops | 4b:122 |
| | Lepidosperma longitudinale Labill., Nov. Holl. Pl. 1: 16, t.13 (1805) | 4b:124 |
| e | Lepidosperma monticola G.T.Plunkett & J.J.Bruhl, Phytokeys 28: 26 (2013) | |
| | Lepidosperma neesii Kunth, Enum. Pl. [Kunth] 2: 319 (1837) | 4b:127 |
| e | Lepidosperma oldfieldii Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 91, t.146A (1858) | 4b:125 |
| | Lepidosperma squamatum Labill. sensu Rodway (1903) = Lepidosperma concavum | 4b:126 |
| | Lepidosperma tetragonum Labill. = Baumea tetragona | |
| | Lepidosperma tortuosum F.Muell., Fragm. (Mueller) 9: 23 (1875) | 4b:121 |
| | Lepidosperma viscidum R.Br., Prodr. Fl. Nov. Holland. 234 (1810) | 4b:126 |
| | Mesomelaena sphaerocephala (R.Br.) Benth. = Gymnoschoenus sphaerocephalus | 4b:119 |
| e | Oreobolus acutifolius S.T.Blake, Contr. Queensland Herb. 8: 39 (1969) | 4b:109 |
| | Oreobolus distichus F.Muell., Trans. Philos. Soc. Victoria 1: 109 (1855) | 4b:109 |
| e | Oreobolus oligocephalus W.M.Curtis, Brunonia 7: 301 (1985) | 4b:110 |
| e | Oreobolus oxycarpus S.T.Blake subsp. brownii Seberg, Bot. J. Linn. Soc. 96: 178 (1988) | 4b:110 |
| | Oreobolus pumilio R.Br. subsp. pumilio , Prodr. Fl. Nov. Holland. 236 (1810) | 4b:109 |
| e | Oreobolus tholicarpus D.I.Morris, Muellera 15: 28 (2001) | |
| | Schoenoides oligocephala (W.M.Curtis) Seberg = Oreobolus oligocephalus | 4b:110 |
| | Schoenoplectus pungens (Vahl) Palla, Bot. Jahrb. Syst. 10: 299 (1888) | 4b:94 |
| | Schoenoplectus tabernaemontani (C.C.Gmel.) Palla, Bot. Jahrb. Syst. 10: 299 (1888) | 4b:95 |
| | Schoenoplectus validus (Vahl) Á.Löve & D.Löve = Schoenoplectus tabernaemontani | 4b:95 |
| e | Schoenus absconditus Kük., Repert. Spec. Nov. Regni Veg. 44: 99 (1938) | 4b:117 |
| | Schoenus acutus Labill. = Baumea acuta | |
| | Schoenus apogon Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 2: 77 (1817) | 4b:114 |
| | Schoenus axillaris (R.Br.) Poir. = Schoenus maschalinus | 4b:116 |
| e | Schoenus biglumis Kük., Repert. Spec. Nov. Regni Veg. 44: 164 (1938) | 4b:118 |
| | Schoenus brevifolius R.Br., Prodr. Fl. Nov. Holland. 231 (1810) | 4b:113 |
| | Schoenus brownii Hook.f. = Schoenus apogon | 4b:114 |
| | Schoenus calyptratus Kük., Repert. Spec. Nov. Regni Veg. 48: 248 (1940) | 4b:117 |
| | Schoenus carsei Cheeseman, Man. New Zealand Fl. 781 (1906) | 4b:113 |
| | Schoenus filum Labill. = Gahnia filum | |
| | Schoenus fluitans Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 81 (1858) | 4b:114 |
| | Schoenus latelaminatus Kük., Repert. Spec. Nov. Regni Veg. 44: 88 (1938) | 4b:116 |
| | Schoenus lepidosperma (F.Muell.) K.L.Wilson subsp. lepidosperma , Telopea 5: 619 (1994) | 4b:114 |
| | Schoenus maschalinus Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 2: 77 (1817) | 4b:116 |
| | Schoenus nitens (R.Br.) Poir., Encycl. (Lamarck) Suppl. 2: 252 (1812) | 4b:118 |
| e | Schoenus pygmaeus S.T.Blake, Contr. Queensland Herb. 8: 41 (1969) | 4b:116 |
| | Schoenus tenuissimus (Hook.f.) Benth. sensu Rodway (1903), Curtis & Morris (1994) = Schoenus lepidosperma subsp. lepidosperma | 4b:114 |
| | Schoenus tesquorum J.M.Black, Fl. S. Austral. [J.M. Black] 1: 90 (1922) | 4b:115 |

| | | |
|--------------------------|--|--------|
| | Schoenus turbinatus (R.Br.) Poir., Encycl. (Lamarck) Suppl. 2: 251 (1811) | 4b:118 |
| | Scirpus americanus Pers. sensu Willis (1970) = Schoenoplectus pungens | 4b:94 |
| | Scirpus antarcticus L. sensu Willis (1970) = Isolepis marginata | 4b:104 |
| | Scirpus aucklandicus (Hook.f.) Boeckeler = Isolepis aucklandica | 4b:102 |
| | Scirpus caldwellii V.J.Cook = Bolboschoenus caldwellii | 4b:92 |
| | Scirpus cartilagineus (R.Br.) Poir. = Isolepis marginata | 4b:104 |
| | Scirpus cartilagineus (R.Br.) Poir. var. propinquus (Nees) Benth. = Isolepis inundata | |
| | Scirpus cernuus Vahl = Isolepis cernua | 4b:99 |
| | Scirpus crassiusculus (Hook.f.) Benth. = Isolepis crassiuscula | 4b:98 |
| | Scirpus fluitans L. = Isolepis fluitans | 4b:98 |
| | Scirpus fluviatilis (Torr.) A.Gray sensu Willis (1970) = Bolboschoenus medianus | 4b:93 |
| | Scirpus hookerianus (Boeckeler) S.T.Blake = Isolepis hookeriana | 4b:104 |
| | Scirpus inundatus (R.Br.) Poir. = Isolepis inundata | 4b:100 |
| | Scirpus lacustris L. sensu Rodway (1903) = Schoenoplectus tabernaemontani | 4b:95 |
| | Scirpus lenticularis (R.Br.) Poir. sensu Rodway (1903) = Isolepis fluitans | 4b:98 |
| | Scirpus maritimus L. sensu Rodway (1903) = Bolboschoenus caldwellii | 4b:92 |
| | Scirpus montivagus S.T.Blake = Isolepis montivaga | 4b:103 |
| | Scirpus nodosus Rottb. = Ficinia nodosa | 4b:105 |
| | Scirpus platycarpus S.T.Blake = Isolepis platycarpa | 4b:99 |
| | Scirpus productus C.B.Clark = Isolepis producta | 4b:98 |
| | Scirpus pungens Vahl = Schoenoplectus pungens | 4b:94 |
| | Scirpus riparius (R.Br.) Poir. = Isolepis cernua | 4b:99 |
| | Scirpus setaceus L. = Isolepis setacea | 4b:105 |
| | Scirpus stellatus C.B.Clark = Isolepis stellata | 4b:104 |
| | Scirpus subtilissimus (Boeckeler) S.T.Blake = Isolepis subtilissima | 4b:101 |
| | Scirpus tasmanicus S.T.Blake = Isolepis tasmanica | 4b:102 |
| | Scirpus triqueter L. sensu Brown (1810) = Schoenoplectus pungens (misapplied in Tasmania) | |
| | Scirpus validus Vahl = Schoenoplectus tabernaemontani | 4b:95 |
| | Scirpus wakefieldianus S.T.Blake = Isolepis wakefieldiana | 4b:100 |
| | Tetraria capillaris (F.Muell.) J.M.Black, Trans. Roy. Soc. South Australia 58: 169 (1934) | 4b:132 |
| | Tricostularia pauciflora (F.Muell.) Benth., Fl. Austral. 7: 383 (1878) | 4b:127 |
| | Uncinia compacta R.Br., Prodr. Fl. Nov. Holland. 241 (1810) | 4b:142 |
| t | Uncinia elegans (Kük.) Hamlin, Rec. Domin. Mus. 19: 11 (1959) | 4b:140 |
| | Uncinia flaccida S.T.Blake, Proc. Roy. Soc. Queensland 51: 49 (1940) | 4b:143 |
| t | Uncinia nervosa Boott, Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 102 (1858) | 4b:142 |
| | Uncinia riparia R.Br., Prodr. Fl. Nov. Holland. 241 (1810) | 4b:140 |
| | Uncinia tenella R.Br., Prodr. Fl. Nov. Holland. 241 (1810) | 4b:143 |
| HAEMODORACEAE | | |
| i # | Anigozanthos flavidus DC., Liliac. (Redouté) 3: t. 176 (1807) | |
| e | Haemodorum distichophyllum Hook., Icon. Pl. 9: t.866 (1851) | 4b:364 |
| HEMEROCALLIDACEAE | | |
| | Caesia alpina Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 373 (1859) | 4b:386 |
| | Caesia calliantha R.J.F.Hend., Fl. Australia 45: 471 (1987) | 4b:387 |

| | | |
|-------------------------|---|--------|
| | Caesia corymbosa R.Br. = Chamaescilla corymbosa | |
| | Caesia parviflora R.Br. var. minor R.J.F.Hend., Fl. Australia 45: 471 (1987) | 4b:386 |
| | Caesia parviflora R.Br. var. parviflora , Prodr. Fl. Nov. Holland. 277 (1810) | 4b:386 |
| | Caesia parviflora R.Br. var. vittata (R.Br.) R.J.F.Hend., Fl. Australia 45: 472 (1987) | 4b:386 |
| | Caesia vittata R.Br. sensu Rodway (1903) = Caesia calliantha | 4b:387 |
| | Caesia vittata R.Br. = Caesia parviflora var. vittata | |
| | Chlorophytum alpinum (Hook.f.) Benth. ex Baker = Caesia alpina | 4b:386 |
| | Dianella amoena G.W.Carr & P.F.Horsfall, Muelleria 8: 369 (1995) | 4b:377 |
| | Dianella archeri Hook.f. = Dianella tasmanica | |
| | Dianella brevicaulis (Ostenf.) G.W.Carr & P.F.Horsfall, Muelleria 8: 375 (1995) | 4b:378 |
| | Dianella caerulea Sims sensu Henderson (1987) = Dianella amoena (Tasmanian plants) | |
| | Dianella laevis R.Br. = Dianella amoena | 4b:377 |
| | Dianella longifolia R.Br. sensu Curtis & Morris (1994) = Dianella amoena | 4b:377 |
| | Dianella longifolia R.Br. var. aspera Rodway a name of uncertain application | |
| | Dianella revoluta R.Br. var. brevicaulis Ostenf. = Dianella brevicaulis | 4b:378 |
| | Dianella revoluta R.Br. var. revoluta , Prodr. Fl. Nov. Holland. 280 (1810) | 4b:378 |
| | Dianella tasmanica Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 57, t.133A (1858) | 4b:378 |
| | Herpolirion novae-zelandiae Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 258 (1853) | 4b:383 |
| | Herpolirion tasmaniae Hook.f. = Herpolirion novae-zelandiae | |
| i | Phormium tenax J.R.Forst. & G.Forst., Char. Gen. Pl.: 48, t.24 (1775) | 4b:418 |
| | Stypandra caespitosa R.Br. = Thelionema caespitosum | 4b:379 |
| | Stypandra umbellata R.Br. = Thelionema umbellatum | |
| | Thelionema caespitosum (R.Br.) R.J.F.Hend., Austrobaileya 2: 109 (1985) | 4b:379 |
| | Thelionema umbellatum (R.Br.) R.J.F.Hend., Austrobaileya 2: 109 (1985) | 4b:380 |
| | Tricoryne elatior R.Br., Prodr. Fl. Nov. Holland. 278 (1810) | 4b:391 |
| HYACINTHACEAE | | |
| | Ornithogalum angustifolium Boreau sensu Buchanan (1999) = Ornithogalum umbellatum | |
| i | Ornithogalum umbellatum L., Sp. Pl. 1: 307 (1753) | 4b:369 |
| i # | Scilla peruviana L., Sp. Pl. 1: 309 (1753) | 4b:391 |
| HYDROCHARITACEAE | | |
| i | Egeria densa Planch., Ann. Sci. Nat., Bot., sér. 3, 11: 80 (1849) | 4b:8 |
| i | Elodea canadensis Michx., Fl. Bor.-Amer. (Michaux) 1: 20 (1803) | 4b:8 |
| | Elodea densa (Planch.) Casp. = Egeria densa | 4b:8 |
| | Halophila australis Doty & B.C.Stone, Brittonia 18: 306, fig.2 (1966) | 4b:13 |
| | Halophila ovalis (R.Br.) Hook.f. sensu Rodway (1903) = Halophila australis | 4b:13 |
| | Lagarosiphon major (Ridl.) Moss previously listed as naturalised but insufficient evidence exists to support this | 4b:9 |
| | Vallisneria americana Michx. sensu Buchanan (2007) = Vallisneria australis | 4b:11 |
| | Vallisneria australis S.W.L.Jacobs & Les, Syst. Bot. 33: 62 (2008) | 4b:11 |
| | Vallisneria gigantea Graebn. sensu Curtis & Morris (1994) = Vallisneria australis | 4b:11 |
| | Vallisneria spiralis L. sensu Rodway (1903) = Vallisneria australis | 4b:11 |
| HYPOXIDACEAE | | |
| | Hypoxis glabella R.Br. var. glabella = Pauridia glabella var. glabella | |

| | | |
|------------------|---|--------|
| | Hypoxis hygrometrica Labill. var. hygrometrica , Nov. Holl. Pl. 1: 82 t.108 (1805) | 4b:376 |
| | Hypoxis hygrometrica Labill. var. villosisepala R.J.F.Hend., Fl. Australia 45: 488 (1987) | 4b:377 |
| | Hypoxis pusilla Hook.f. = Pauridia glabella var. glabella | 4b:375 |
| | Hypoxis vaginata Schltld. var. brevistigmata R.J.F.Hend. = Pauridia vaginata var. brevistigmata | 4b:376 |
| | Hypoxis vaginata Schltld. var. vaginata = Pauridia vaginata var. vaginata | 4b:376 |
| | Pauridia glabella (R.Br.) Snijman & Kocyan var. glabella , Phytotaxa 116: 27 (2013) | 4b:375 |
| | Pauridia vaginata (Schltld.) Snijman & Kocyan var. brevistigmata (R.J.F.Hend.) Snijman & Kocyan, Phytotaxa 116: 31 (2013) | 4b:376 |
| | Pauridia vaginata (Schltld.) Snijman & Kocyan var. vaginata (Schltld.) Snijman & Kocyan, Phytotaxa 116: 31 (2013) | 4b:376 |
| IRIDACEAE | | |
| i | Chasmanthe floribunda (Salisb.) N.E.Br., Trans. Roy. Soc. South Africa 20: 274 (1932) | |
| i | Crocosmia × crocosmiiflora (Lemoine ex E.Morren) N.E.Br., Trans. Roy. Soc. South Africa 20: 264 (1932) | 4b:417 |
| e | Diplarrena latifolia Benth., Fl. Austral. 6: 400 (1873) | 4b:405 |
| | Diplarrena moraea Labill., Voy. Rech. Pérouse 1: 157 (1800) | 4b:405 |
| | Diplarrena moraea Labill. var. alpina Hook.f. = Diplarrena moraea | |
| i | Freesia a complex of garden hybrids involving <i>F. alba</i> and <i>F. leichtlinii</i> | 4b:416 |
| | Genosiris fragilis Labill. = Patersonia fragilis | |
| i | Gladiolus × colvillii Sweet, Brit. Fl. Gard. [Sweet] 2: t.155 (1826) | |
| i | Gladiolus communis L. subsp. byzantinus (Mill.) A.P.Ham., Bot. J. Linn. Soc. 76: 358 (1978) | 4b:413 |
| | Gladiolus cuspidatus Jacq. = Gladiolus undulatus | 4b:413 |
| i | Gladiolus tristis L., Sp. Pl., ed. 2: 53 (1762) | 4b:411 |
| i | Gladiolus undulatus L., Mant. Pl. 1: 27 (1767) | 4b:413 |
| i t | Hesperantha coccinea (Backh. & Harv.) Goldblatt & J.C.Manning, Novon 6: 263 (1996) | 4b:417 |
| | Hewardia tasmanica Hook. = Isophysis tasmanica | 4b:401 |
| | Homeria breyniana (L.) G.J.Lewis sensu Willis (1973) = Moraea flaccida | 4b:408 |
| | Homeria flaccida Sweet = Moraea flaccida | 4b:408 |
| i | Iris foetidissima L., Sp. Pl. 1: 39 (1753) | 4b:406 |
| i | Iris germanica L., Sp. Pl. 1: 38 (1753) | 4b:406 |
| e | Isophysis tasmanica (Hook.) T.Moore, Proc. Linn. Soc. London 2: 212 (1853) | 4b:401 |
| i | Ixia flexuosa L., Sp. Pl., ed. 2: 51 (1762) | 4b:414 |
| i | Ixia maculata L., Sp. Pl., ed. 2: 1664 (1763) | 4b:414 |
| i | Ixia polystachya L., Sp. Pl., ed. 2: 51 (1762) | 4b:414 |
| | Libertia laurencei Hook.f. = Libertia pulchella | |
| | Libertia pulchella (R.Br.) Spreng. var. pulchella , Syst. Veg. (ed. 16) [Sprengel] 1: 69 (1824) | 4b:402 |
| e | Libertia pulchella (R.Br.) Spreng. var. pygmaea D.I.Morris, The Student's Flora of Tasmania 4b: 425 (1994) | 4b:402 |
| i | Moraea flaccida (Sweet) Steud., Nomencl. Bot. [Steudel], ed. 2. 2: 160 (1841) | 4b:408 |
| | Patersonia fragilis (Labill.) Asch. & Graebn., Syn. Mitteleur. Fl. [Ascherson & Graebner]. 3: 532 (1906) | 4b:403 |
| | Patersonia glauca R.Br. = Patersonia fragilis | 4b:403 |
| | Patersonia longiscapa Sweet = Patersonia occidentalis | 4b:403 |
| | Patersonia occidentalis R.Br. var. occidentalis , Prodr. Fl. Nov. Holland. 304 (1810) | 4b:403 |

| | | |
|-----|--|--------|
| | Romulea longifolia (Salisb.) Baker = Romulea rosea var. australis | 4b:409 |
| | Romulea minutiflora Klatt sensu Buchanan (2009) probably Romulea rosea | |
| i | Romulea rosea (L.) Eckl. var. australis (Ewart) M.P.de Vos, J. S. African Bot. Suppl. 9: 254 (1972) | 4b:409 |
| | Schizostylis coccinea Backh. & Harv. = Hesperantha coccinea | 4b:417 |
| i | Sisyrinchium iridifolium Kunth, Nov. Gen. Sp. [H.B.K.] 1: 324 (1816) | 4b:402 |
| i | Sparaxis bulbifera (L.) Ker Gawl., Ann. Bot. [König & Sims] 1: 226 (1804) | 4b:415 |
| | Trichonema roseum Ker Gawl. sensu Rodway (1903) = Romulea rosea | 4b:409 |
| i # | Tritonia gladiolaris (Lam.) Goldblatt & J.C.Manning, Bothalia 36: 57 (2006) | 4b:416 |
| | Tritonia lineata (Salisb.) Ker Gawl. = Tritonia gladiolaris | 4b:416 |
| | Watsonia bulbifera J.W.Mathews & L.Bolus = Watsonia meriana var. bulbillifera | 4b:408 |
| i | Watsonia meriana (L.) Mill. var. bulbillifera (J.W.Mathews & L.Bolus) D.A.Cooke, J. Adelaide Bot. Gard. 18: 6 (1998) | 4b:408 |
| i | Watsonia meriana (L.) Mill. var. meriana , Gard. Dict., ed. 8. no.1 (1768) | 4b:409 |
| i | Watsonia versfeldii J.W.Mathews & L.Bolus, Ann. Bolus Herb. 3: 140 (1922) | 4b:409 |
| | Watsonia versfeldii J.W.Mathews & L.Bolus var. alba J.W.Mathews & L.Bolus = Watsonia versfeldii | 4b:409 |

JUNCACEAE

| | | |
|---|---|-------|
| i | Juncus acuminatus Michx., Fl. Bor.-Amer. (Michaux) 1: 192 (1803) | 4b:75 |
| i | Juncus acutiflorus Ehrh. ex Hoffm., Deutschl. Fl. (Hoffm.) 1: 125 (1791) | 4b:77 |
| i | Juncus acutus L., Sp. Pl. 1: 325 (1753) | |
| | Juncus amabilis Edgar, New Zealand J. Bot. 2: 186, figs.7, 21 (1964) | 4b:67 |
| | Juncus antarcticus Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 79, pl.46 (1844) | 4b:71 |
| i | Juncus articulatus L., Sp. Pl. 1: 327 (1753) | 4b:77 |
| e | Juncus astreptus L.A.S.Johnson, Aspects of Tasmanian Botany: 41 (1991) | 4b:66 |
| | Juncus australis Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 66, t.134a (1858) | 4b:67 |
| | Juncus bassianus L.A.S.Johnson, Aspects of Tasmanian Botany: 40 (1991) | 4b:66 |
| | Juncus bufonius L., Sp. Pl. 1: 328 (1753) | 4b:63 |
| i | Juncus bulbosus L., Sp. Pl. 1: 327 (1753) | 4b:73 |
| | Juncus caespiticus E.Mey., Pl. Preiss. [J.G.C.Lehman] 2: 47 (1846) | 4b:72 |
| | Juncus capillaceus Hook.f. sensu Rodway (1903) = Juncus sandwithii | 4b:74 |
| i | Juncus capitatus Weigel, Observ. Bot. (Weigel): 28 (1772) | 4b:70 |
| | Juncus communis E.Mey. sensu Rodway (1903) = Juncus australis | 4b:67 |
| i | Juncus conglomeratus L., Sp. Pl. 1: 326 (1753) | 4b:69 |
| e | Juncus curtisiae L.A.S.Johnson, Aspects of Tasmanian Botany: 44 (1991) | 4b:74 |
| i | Juncus effusus L., Sp. Pl. 1: 326 (1753) | 4b:69 |
| | Juncus falcatus E.Mey. subsp. falcatus , Syn. Luzul. 34 (1823) | 4b:71 |
| | Juncus filicaulis Buchenau, Proc. Linn. Soc. New South Wales 28: 912 (1904) | 4b:68 |
| | Juncus fockei Buchenau, Bot. Jahrb. Syst. 12: 358 (1890) | 4b:75 |
| i | Juncus gerardii Loisel., J. Bot. (Desvaux) 2: 284 (1809) | |
| | Juncus gregiflorus L.A.S.Johnson, Contr. New South Wales Natl. Herb. 3: 243 (1963) | 4b:66 |
| | Juncus gunnii Hook.f. = Juncus pauciflorus | |
| | Juncus holoschoenus R.Br., Prodr. Fl. Nov. Holland. 259 (1810) | 4b:75 |
| i | Juncus indescriptus Steud., Syn. Pl. Glum. 2: 304 (1855) | 4b:72 |
| | Juncus kraussii Hochst. subsp. australiensis (Buchenau) Snogerup, Willdenowia 23: 61 (1993) | 4b:70 |

| | | |
|-----|---|-------|
| | <i>Juncus maritimus</i> Lam. sensu Rodway (1903) = <i>Juncus kraussii</i> subsp. <i>australiensis</i> | 4b:70 |
| i # | <i>Juncus microcephalus</i> Kunth, Bonpland & Kunth, Nov. Gen. Sp. [H.B.K.] 1: 237 (1816) | 4b:75 |
| | <i>Juncus pallidus</i> R.Br., Prodr. Fl. Nov. Holland. 258 (1810) | 4b:64 |
| | <i>Juncus pauciflorus</i> R.Br., Prodr. Fl. Nov. Holland. 259 (1810) | 4b:65 |
| | <i>Juncus planifolius</i> R.Br., Prodr. Fl. Nov. Holland. 259 (1810) | 4b:71 |
| | <i>Juncus prismatocarpus</i> R.Br., Prodr. Fl. Nov. Holland. 259 (1810) | 4b:73 |
| | <i>Juncus procerus</i> E.Mey., Linnaea 3: 367 (1828) | 4b:65 |
| | <i>Juncus pusillus</i> Buchenau sensu Willis (1973) = <i>Juncus sandwithii</i> | 4b:74 |
| e | <i>Juncus ratkowskyanus</i> L.A.S.Johnson, Aspects of Tasmanian Botany: 43 (1991) | 4b:74 |
| | <i>Juncus revolutus</i> R.Br., Prodr. Fl. Nov. Holland. 259 (1810) | 4b:63 |
| | <i>Juncus sandwithii</i> Lourteig, Comite National Francais des Recherches Antarctiques 23: 44 (1968) | 4b:74 |
| | <i>Juncus sarophorus</i> L.A.S.Johnson, Contr. New South Wales Natl. Herb. 3: 242 (1963) | 4b:68 |
| i t | <i>Juncus squarrosus</i> L., Sp. Pl. 1: 327 (1753) | 4b:62 |
| | <i>Juncus subsecundus</i> N.A.Wakef., Vict. Naturalist 73: 211 (1957) | 4b:69 |
| i | <i>Juncus tenuis</i> Willd., Sp. Pl., ed. 4 [Willdenow] 2: 214 (1799) | 4b:62 |
| | <i>Juncus vaginatus</i> R.Br., Prodr. Fl. Nov. Holland. 258 (1810) | 4b:65 |
| | <i>Luzula acutifolia</i> H.Nordensk. subsp. <i>acutifolia</i> , Bot. Not. 122: 85 (1969) | 4b:84 |
| | <i>Luzula atrata</i> Edgar, New Zealand J. Bot. 13: 794 (1975) | 4b:83 |
| e | <i>Luzula australasica</i> Steud. subsp. <i>australasica</i> , Syn. Pl. Glumac. 2: 294 (1855) | 4b:82 |
| i t | <i>Luzula campestris</i> (L.) DC., Fl. Franc. (DC. & Lamarck), ed. 3. 3: 161 (1805) | 4b:81 |
| i t | <i>Luzula congesta</i> (Thuill.) Lej., Fl. Spa 1: 168 (1811) | 4b:81 |
| | <i>Luzula densiflora</i> (H.Nordensk.) Edgar, New Zealand J. Bot. 13: 786 (1975) | 4b:80 |
| | <i>Luzula flaccida</i> (Buchenau) Edgar, New Zealand J. Bot. 13: 786 (1975) | 4b:80 |
| | <i>Luzula meridionalis</i> H.Nordensk., Bot. Not. 122: 71 (1969) | 4b:78 |
| | <i>Luzula modesta</i> Buchenau, Pflanzenr. (Engler) 25: 80, fig.51 (1906) | 4b:82 |
| i t | <i>Luzula multiflora</i> (Ehrh.) Lej., Fl. Spa 1: 169 (1811) | 4b:81 |
| | <i>Luzula novae-cambriae</i> Gand., Bull. Soc. Bot. France 46: 392 (1900) | 4b:82 |
| | <i>Luzula oldfieldii</i> Hook.f. = <i>Luzula australasica</i> subsp. <i>australasica</i> | 4b:82 |
| e | <i>Luzula poimena</i> W.M.Curtis, Brunonia 7: 297 (1985) | 4b:83 |

JUNCAGINACEAE

| | | |
|--|---|--------|
| | <i>Cycnogeton alcockiae</i> (Aston) Mering & Kadereit, Diversity, Phylogeny, and Evolution in the Monocotyledons: 73 (2010) | 4b:427 |
| | <i>Cycnogeton procerum</i> (R.Br.) Buchenau, Abh. Naturwiss. Vereins Bremen 224 (1867) | 4b:14 |
| | <i>Cycnogeton rheophilum</i> (Aston) Mering & Kadereit, Diversity, Phylogeny, and Evolution in the Monocotyledons: 73 (2010) | 4b:427 |
| | <i>Triglochin alcockiae</i> Aston = <i>Cycnogeton alcockiae</i> | 4b:427 |
| | <i>Triglochin centrocarpum</i> Hook. = <i>Triglochin nana</i> (Tasmanian plants) | 4b:16 |
| | <i>Triglochin decipiens</i> R.Br. = <i>Triglochin striata</i> | |
| | <i>Triglochin minutissima</i> F.Muell., Fragm. (Mueller) 6: 82 (1867) | 4b:16 |
| | <i>Triglochin mucronata</i> R.Br., Prodr. Fl. Nov. Holland. 343 (1810) | 4b:16 |
| | <i>Triglochin nana</i> F.Muell., Fragm. (Mueller) 6: 82 (1867) | 4b:16 |
| | <i>Triglochin procera</i> R.Br. = <i>Cycnogeton procerum</i> | 4b:14 |
| | <i>Triglochin rheophila</i> Aston = <i>Cycnogeton rheophilum</i> | 4b:427 |

- Triglochin striata** Ruiz & Pav., Fl. Peruv. 3: 72 (1802) 4b:16
 Triglochin triandra Michx. sensu Hooker (1860) = **Triglochin striata**

LAXMANNIACEAE

- Arthropodium laxum Hook.f. nom. illeg. = **Arthropodium strictum**
Arthropodium milleflorum (DC.) J.F.Macbr., Contr. Gray Herb. 3(56): 2 (1918) 4b:384
Arthropodium minus R.Br., Prodr. Fl. Nov. Holland. 276 (1810) 4b:385
 Arthropodium paniculatum (Haw.) R.Br. = **Arthropodium milleflorum** 4b:384
 e **Arthropodium pendulum** DC., Cat. Pl. Horti Monsp. 80 (1813)
Arthropodium strictum R.Br., Prodr. Fl. Nov. Holland. 276 (1810) 4b:388
Chamaescilla corymbosa (R.Br.) F.Muell. ex Benth. var. **corymbosa**, Fl. Austral. 7: 48 (1878) 4b:387
 i **Cordyline australis** (G.Forst.) Endl., Prodr. Fl. Norfolk. 29 (1833)
 Dichopogon strictus (R.Br.) Baker = **Arthropodium strictum** 4b:388
 Laxmannia minor R.Br. sensu Hooker (1860) = **Laxmannia orientalis**
 Laxmannia morrisii Keighery = **Laxmannia squarrosa**
Laxmannia orientalis Keighery, Fl. Australia 45: 492 (1987) 4b:388
 Laxmannia sessiliflora Decne. sensu Rodway (1903) = **Laxmannia orientalis** 4b:388
 Laxmannia squarrosa Lindl. recorded in error
Sowerbaea juncea Andrews, Bot. Repos. 2: t.81 (1800) 4b:389
Thysanotus patersonii R.Br., Prodr. Fl. Nov. Holland. 284 (1810) 4b:389

LUZURIAGACEAE

- Drymophila cyanocarpa** R.Br., Prodr. Fl. Nov. Holland. 292 (1810) 4b:369

MELANTHIACEAE

- Aletris punicea Labill. = **Blandfordia punicea**

ORCHIDACEAE

- Acianthus bifolius R.Br. = **Chiloglottis reflexa**
Acianthus caudatus R.Br., Prodr. Fl. Nov. Holland. 321 (1810) 4a:117
 Acianthus exsertus R.Br. sensu Curtis (1979) = **Acianthus pusillus** 4a:117
Acianthus pusillus D.L.Jones, Austral. Orchid Res. 2: 7 (1991) 4a:117
 Acianthus reniformis (R.Br.) Schltr. = **Cyrtostylis reniformis** 4a:119
 Acianthus viridis Hook.f. = **Townsonia viridis** 4a:118
 Anzybas fordhamii (Rupp) D.L.Jones & M.A.Clem. = **Corybas fordhamii** 4a:124
 Anzybas unguiculatus (R.Br.) D.L.Jones & M.A.Clem. = **Corybas unguiculatus** 4a:124
 Arachnorchis anthracina (D.L.Jones) D.L.Jones & M.A.Clem. = **Caladenia anthracina**
 Arachnorchis australis (G.W.Carr) D.L.Jones & M.A.Clem. = **Caladenia australis**
 Arachnorchis brachyscapa (G.W.Carr) D.L.Jones & M.A.Clem. = **Caladenia brachyscapa**
 Arachnorchis cardiochila (Tate) D.L.Jones & M.A.Clem. = **Caladenia cardiochila**
 Arachnorchis caudata (Nicholls) D.L.Jones & M.A.Clem. = **Caladenia caudata** 4a:101
 Arachnorchis clavigera (A.Cunn. ex Lindl.) D.L.Jones & M.A.Clem. = **Caladenia clavigera** 4a:105
 Arachnorchis dienema (D.L.Jones) D.L.Jones & M.A.Clem. = **Caladenia dienema**
 Arachnorchis dilatata (R.Br.) D.L.Jones & M.A.Clem. = **Caladenia dilatata** 4a:100
 Arachnorchis echidnachila (Nicholls) D.L.Jones & M.A.Clem. = **Caladenia echidnachila** 4a:101
 Arachnorchis helvina (D.L.Jones) D.L.Jones & M.A.Clem. = **Caladenia helvina**
 Arachnorchis lindleyana (Rchb.f.) D.L.Jones & M.A.Clem. = **Caladenia lindleyana**

| | | |
|---|---|--------|
| | Arachnorchis pallida (Lindl.) D.L.Jones & M.A.Clem. = Caladenia pallida | 4a:104 |
| | Arachnorchis patersonii (R.Br.) D.L.Jones & M.A.Clem. = Caladenia patersonii | 4a:100 |
| | Arachnorchis saggicola (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia saggicola | |
| | Arthrochilus huntianus (F.Muell.) Blaxell subsp. huntianus = Thynninorchis huntiana | 4a:89 |
| | Arthrochilus huntianus (F.Muell.) Blaxell subsp. nothofagicola D.L.Jones = Thynninorchis nothofagicola | |
| | Bunochilus melagrammus (D.L.Jones) D.L.Jones & M.A.Clem. = Pterostylis melagramma | 4a:27 |
| | Bunochilus stenochilus (D.L.Jones) D.L.Jones & M.A.Clem. = Pterostylis stenochila | |
| | Bunochilus tunstallii (D.L.Jones & M.A.Clem.) D.L.Jones & M.A.Clem. = Pterostylis tunstallii | |
| | Bunochilus williamsonii (D.L.Jones) D.L.Jones & M.A.Clem. = Pterostylis williamsonii | |
| | Burnettia cuneata Lindl., Gen. Sp. Orchid. Pl. 518 (1840) | 4a:92 |
| | Caladenia alata R.Br., Prodr. Fl. Nov. Holland. 324 (1810) | 4a:107 |
| | Caladenia alba R.Br. recorded in error | |
| | Caladenia alpina R.S.Rogers, Trans. Roy. Soc. South Australia 51: 12 (1927) | 4a:111 |
| e | Caladenia angustata Lindl., Gen. Sp. Orchid. Pl. 420 (1840) | 4a:108 |
| e | Caladenia anthracina D.L.Jones, Austral. Orchid Res. 3: 21 (1998) | |
| | Caladenia atkinsonii Rodway = Caladenia carnea | 4a:107 |
| e | Caladenia atrata D.L.Jones, Muelleria 8: 178 (1994) | 4a:111 |
| e | Caladenia atrochila D.L.Jones, Austral. Orchid Res. 3: 23 (1998) | |
| | Caladenia aurantiaca (R.S.Rogers) Rupp, Proc. Linn. Soc. New South Wales 71: 280 (1947) | |
| | Caladenia australis G.W.Carr, Indigenous Flora and Fauna Association Miscellaneous Paper 1: 2 (1991) | |
| | Caladenia barbata Lindl. sensu Hooker (1860) = Pheladenia deformis | |
| | Caladenia brachyscapa G.W.Carr, Muelleria 6: 439 (1988) | |
| | Caladenia caerulea R.Br. sensu Bentham (1873) = <i>Cyanicula caerulea</i> | |
| e | Caladenia campbellii D.L.Jones, Austral. Orchid Res. 3: 25 (1998) | |
| x | Caladenia cardiochila Tate, Trans. Roy. Soc. South Australia 9: 60, t.2 (1887) | |
| | Caladenia carnea R.Br., Prodr. Fl. Nov. Holland. 324 (1810) | 4a:106 |
| | Caladenia carnea R.Br. var. pygmaea R.S.Rogers sensu Curtis (1953) = Caladenia pusilla | |
| | Caladenia carnea R.Br. var. quadriseriata Benth. = Caladenia gracilis | |
| | Caladenia catenata (Sm.) Druce var. catenata sensu Curtis (1979) = Caladenia carnea & C. sylvicola | 4a:106 |
| | Caladenia catenata (Sm.) Druce var. exigua sensu Curtis (1979) = Caladenia alata | 4a:107 |
| | Caladenia catenata (Sm.) Druce var. gigantea sensu Curtis (1979) = Caladenia carnea | 4a:106 |
| | Caladenia catenata (Sm.) Druce var. minor sensu Curtis (1979) = Caladenia mentiens | 4a:106 |
| e | Caladenia caudata Nicholls, Vict. Naturalist 64: 231 (1948) | 4a:101 |
| | Caladenia clavigera A.Cunn. ex Lindl., Gen. Sp. Orchid. Pl. 422 (1840) | 4a:105 |
| | Caladenia congesta R.Br., Prodr. Fl. Nov. Holland. 324 (1810) | 4a:112 |
| | Caladenia cordiformis R.S.Rogers sensu Nicholls & Olsen 1941 = Caladenia clavigera | |
| | Caladenia corynepetala D.L.Jones = Caladenia dilatata | 4a:100 |
| e | Caladenia cracens D.L.Jones, Muelleria 9: 46 (1996) | 4a:111 |
| | Caladenia cucullata Fitzg. sensu Curtis (1979) = Caladenia atrata & C. gracilis | 4a:111 |
| | Caladenia deformis R.Br. = Pheladenia deformis | 4a:113 |
| e | Caladenia dienema D.L.Jones, Austral. Orchid Res. 3: 28 (1998) | |
| | Caladenia dilatata R.Br., Prodr. Fl. Nov. Holland. 325 (1810) | 4a:100 |

| | | |
|---|--|--------|
| | Caladenia dimorpha Fitzg. sensu Curtis (1979) = Caladenia gracilis (?) | 4a:109 |
| e | Caladenia echidnachila Nicholls, Pap. & Proc. Roy. Soc. Tasmania 1932: 13, t.6 (1933) | 4a:101 |
| | Caladenia filamentosa R.Br., Prodr. Fl. Nov. Holland. 324 (1810) | 4a:99 |
| | Caladenia fuscata (Rchb.f.) M.A.Clem. & D.L.Jones, Austral. Orchid Res. 1: 25 (1989) | |
| | Caladenia gracilis R.Br., Prodr. Fl. Nov. Holland. 324 (1810) | 4a:110 |
| e | Caladenia helvina D.L.Jones, Austral. Orchid Res. 2: 26, f.30 (1991) | |
| | Caladenia iridescens R.S.Rogers sensu Curtis (1979) = Caladenia transitoria | 4a:112 |
| | Caladenia latifolia R.Br., Prodr. Fl. Nov. Holland. 324 (1810) | 4a:105 |
| e | Caladenia lindleyana (Rchb.f.) M.A.Clem. & D.L.Jones, Austral. Orchid Res. 1: 27 (1989) | |
| | Caladenia longii R.S.Rogers = Caladenia angustata | 4a:108 |
| | Caladenia lyallii Hook.f. sensu Curtis (1979) = Caladenia cracens & C. alpina | 4a:111 |
| | Caladenia mentiensi D.L.Jones, Austral. Orchid Res. 3: 35 (1998) | 4a:106 |
| | Caladenia menziesii R.Br. = Leptoceras menziesii | 4a:99 |
| e | Caladenia pallida Lindl., Gen. Sp. Orchid. Pl. 421 (1840) | 4a:104 |
| | Caladenia patersonii R.Br., Prodr. Fl. Nov. Holland. 324 (1810) | 4a:100 |
| | Caladenia patersonii R.Br. var. dilatata (R.Br.) Benth. = Caladenia dilatata | |
| | Caladenia praecox Nicholls sensu Curtis (1979) = Caladenia angustata (?) | 4a:109 |
| | Caladenia prolata D.L.Jones, Austral. Orchid Res. 2: 30 (1991) | |
| | Caladenia pusilla W.M.Curtis, The Student's Flora of Tasmania 4a: 133 (1980) | 4a:107 |
| | Caladenia reticulata Fitzg. sensu Curtis (1979) = Caladenia lindleyana | |
| e | Caladenia saggicola D.L.Jones, Austral. Orchid Res. 3: 39 (1998) | |
| | Caladenia simulans G.W.Carr = Caladenia dilatata | |
| | Caladenia suaveolens (R.Br.) Rchb.f. = Lyperanthus suaveolens | 4a:94 |
| e | Caladenia sylvicola D.L.Jones, Austral. Orchid Res. 3: 40 (1998) | |
| | Caladenia testacea R.Br. sensu Rodway (1903) = Caladenia atrata | |
| e | Caladenia tonellii D.L.Jones, Austral. Orchid Res. 3: 41 (1998) | |
| | Caladenia transitoria D.L.Jones, Austral. Orchid Res. 3: 42 (1998) | 4a:112 |
| | Caladenia tutelata R.S.Rogers = Calassodia tutelata | 4a:113 |
| | Caladenia venusta G.W.Carr sensu Buchanan (1995) = Caladenia saggicola | |
| | Caladenia vulgaris D.L.Jones, Austral. Orchid Res. 2: 34 (1991) | |
| | Caleana major R.Br., Prodr. Fl. Nov. Holland. 329 (1810) | 4a:87 |
| | Caleana minor R.Br. = Paracaleana minor | 4a:88 |
| | Calochilus campestris R.Br., Prodr. Fl. Nov. Holland. 320 (1810) | 4a:53 |
| e | Calochilus herbaceus Lindl., Gen. Sp. Orchid. Pl. 459 (1840) | 4a:55 |
| | Calochilus imberbis R.S.Rogers, Trans. & Proc. Roy. Soc. South Australia 51: 4 (1927) | 4a:53 |
| | Calochilus paludosus R.Br., Prodr. Fl. Nov. Holland. 320 (1810) | 4a:56 |
| | Calochilus platytilus D.L.Jones, Orchadian 15: 547 (2008) | 4a:55 |
| | Calochilus robertsonii Benth. sensu Curtis (1979) = Calochilus platytilus | 4a:55 |
| | Calochilus saprophyticus R.S.Rogers sensu Curtis (1953) = Calochilus campestris | |
| | Calonema filamentosum (R.Br.) D.L.Jones & M.A.Clem. = Caladenia filamentosa | 4a:99 |
| | Calonemorhis filamentosa (R.Br.) Szlach. = Caladenia filamentosa | 4a:99 |
| | Chiloglottis cornuta Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 69 (1844) | 4a:85 |
| | Chiloglottis diphylla R.Br. sensu Rodway (1903) = Chiloglottis reflexa | 4a:86 |

| | | |
|---|--|--------|
| e | Chiloglottis grammata G.W.Carr, Indigenous Flora and Fauna Association Miscellaneous Paper 1: 20 (1991) | 4a:86 |
| e | Chiloglottis gunnii Lindl., Gen. Sp. Orchid. Pl. 387 (1840) | 4a:84 |
| | Chiloglottis × pescottiana R.S.Rogers sensu Curtis (1953, 1979) = Chiloglottis grammata | 4a:86 |
| | Chiloglottis platychila G.W.Carr = Chiloglottis gunnii | |
| | Chiloglottis reflexa (Labill.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916, Suppl. 2: 614 (1917) | 4a:86 |
| | Chiloglottis trapeziformis Fitzg., Austral. Orch. 1(3): t.9 (1877) | 4a:86 |
| e | Chiloglottis triceratops D.L.Jones, Austral. Orchid Res. 3: 66 (1998) | |
| | Chiloglottis trilabra Fitzg. = Chiloglottis reflexa (Tasmanian plants) | |
| | Chiloglottis valida D.L.Jones, Austral. Orchid Res. 2: 43 (1991) | |
| e | Chiloglottis sp. Wielangta (M.Wapstra 934) Tas Herbarium | |
| | Corunastylis archeri (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13: 460 (2002) | 4a:82 |
| e | Corunastylis brachystachya (Lindl.) D.L.Jones & M.A.Clem., Orchadian 13: 460 (2002) | 4a:79 |
| | Corunastylis despectans (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13: 460 (2002) | 4a:76 |
| e | Corunastylis firthii (Cady) D.L.Jones & M.A.Clem., Orchadian 13: 461 (2002) | 4a:80 |
| | Corunastylis morrisii (Nicholls) D.L.Jones & M.A.Clem., Orchadian 13: 461 (2002) | 4a:83 |
| | Corunastylis nuda (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13: 461 (2002) | 4a:82 |
| e | Corunastylis nudiscapa (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13: 461 (2002) | 4a:79 |
| | Corunastylis pumila (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13: 461 (2002) | 4a:80 |
| e | Corunastylis tasmanica (D.L.Jones) D.L.Jones & M.A.Clem., Orchadian 13: 462 (2002) | 4a:77 |
| | Corybas aconitiflorus Salisb., Parad. Lond. 1(2): t.83 (1806) | 4a:123 |
| | Corybas diemenicus (Lindl.) Rchb.f., Beitr. Syst. Pflanzenk. 42 (1871) | 4a:121 |
| | Corybas diemenicus (Lindl.) Rupp sensu Curtis (1979) = Corybas incurvus | 4a:121 |
| | Corybas dilatatus (Rupp & Nicholls) Rupp & Nicholls ex Rupp = Corybas diemenicus | 4a:121 |
| | Corybas fimbriatus (R.Br.) Rchb.f., Beitr. Syst. Pflanzenk. 42 (1871) | 4a:121 |
| | Corybas fordhamii (Rupp) Rupp, Vict. Naturalist 59: 61 (1942) | 4a:124 |
| | Corybas incurvus D.L.Jones & M.A.Clem., Kew Bull. 43: 135 (1988) | 4a:121 |
| | Corybas unguiculatus (R.Br.) Rchb.f., Beitr. Syst. Pflanzenk. 43 (1871) | 4a:124 |
| | Corysanthes bicalcarata R.Br. = Corybas aconitiflorus | 4a:123 |
| | Corysanthes diemenica Lindl. = Corybas diemenicus | 4a:121 |
| | Corysanthes dilatata Rupp & Nicholls sensu Nicholls & Olsen 1941 = Corybas diemenicus | |
| | Corysanthes fimbriata R.Br. = Corybas fimbriatus | 4a:121 |
| | Corysanthes incurva (D.L.Jones & M.A.Clem.) D.L.Jones & M.A.Clem. = Corybas incurvus | 4a:121 |
| | Corysanthes pruinosa R.Cunn. sensu Rodway (1903) = Corybas incurvus | 4a:121 |
| | Corysanthes unguiculata R.Br. = Corybas unguiculatus | |
| | Crangonorchis pedoglossa (Fitzg.) D.L.Jones & M.A.Clem. = Pterostylis pedoglossa | 4a:13 |
| | Cryptostylis leptochila F.Muell. ex Benth., Fl. Austral. 6: 334 (1873) | 4a:126 |
| | Cryptostylis longifolia R.Br. = Cryptostylis subulata | 4a:125 |
| | Cryptostylis subulata (Labill.) Rchb.f., Beitr. Syst. Pflanzenk. 15 (1871) | 4a:125 |
| | Cyanicula caerulea (R.Br.) Hopper & A.P.Br. recorded in error | |
| | Cyrtostylis reniformis R.Br., Prodr. Fl. Nov. Holland. 322 (1810) | 4a:119 |
| | Cyrtostylis robusta D.L.Jones & M.A.Clem., Lindleyana 2: 157 (1987) | |
| | Dendrobium striolatum Rchb.f. = Dockrillia striolata | 4a:129 |

| | | |
|---|---|--------|
| | Diplodium alatum (Labill.) D.L.Jones & M.A.Clem. = Pterostylis alata | 4a:20 |
| | Diplodium atrans (D.L.Jones) D.L.Jones & M.A.Clem. = Pterostylis atrans | 4a:22 |
| | Diplodium decurvum (R.S.Rogers) D.L.Jones & M.A.Clem. = Pterostylis decurva | 4a:22 |
| | Diplodium grandiflorum (R.Br.) D.L.Jones & M.A.Clem. = Pterostylis grandiflora | 4a:20 |
| | Dipodium punctatum (Sm.) R.Br. sensu Curtis (1979) = Dipodium roseum | 4a:131 |
| | Dipodium roseum D.L.Jones & M.A.Clem., Austral. Orchid Res. 2: 51 (1991) | 4a:131 |
| i | Disa bracteata Sw., Kongl. Vetensk. Acad. Nya Handl. 21: 211 (1800) | |
| | Disperis alata Labill. = Pterostylis alata | |
| | Diuris chryseopsis D.L.Jones, Austral. Orchid Res. 3: 74 (1998) | 4a:34 |
| | Diuris corymbosa Lindl. sensu Curtis (1979) = Diuris orientis | 4a:36 |
| | Diuris flavopurpurea Messmer sensu Curtis (1979) = Diuris pardina | 4a:35 |
| e | Diuris lanceolata Lindl., Gen. Sp. Orchid. Pl. 508 (1840) | 4a:34 |
| | Diuris longifolia R.Br. sensu Curtis (1979) = Diuris orientis | 4a:36 |
| | Diuris maculata Sm. sensu Curtis (1979) = Diuris pardina | 4a:34 |
| | Diuris monticola D.L.Jones, Austral. Orchid Res. 3: 76 (1998) | |
| | Diuris orientis D.L.Jones, Austral. Orchid Res. 3: 77 (1998) | 4a:36 |
| | Diuris palachila R.S.Rogers sensu Curtis (1979) = Diuris chryseopsis × D. pardina | 4a:35 |
| | Diuris palustris Lindl., Gen. Sp. Orchid. Pl. 507 (1840) | 4a:32 |
| | Diuris pardina Lindl., Gen. Sp. Orchid. Pl. 507 (1840) | 4a:34 |
| | Diuris pedunculata R.Br. sensu Curtis (1979) = Diuris chryseopsis p.p.maj. | 4a:34 |
| | Diuris sulphurea R.Br., Prodr. Fl. Nov. Holland. 316 (1810) | 4a:36 |
| e | Dockrillia striolata (Rchb.f.) Rauschert subsp. chrysantha D.L.Jones, Austral. Orchid Res. 3: 9 (1998) | |
| | Dockrillia striolata (Rchb.f.) Rauschert subsp. striolata , Feddes Repert. 94(7-8): 447 (1983) | |
| | Epipactis cucullata Labill. = Eriochilus cucullatus | |
| | Epipactis reflexa Labill. = Chiloglottis reflexa | |
| | Eriochilus autumnalis R.Br. sensu Rodway (1903) = Eriochilus cucullatus | 4a:91 |
| | Eriochilus cucullatus (Labill.) Rchb.f., Beitr. Syst. Pflanzenk. 27 (1871) | 4a:91 |
| | Eriochilus magenteus D.L.Jones sensu Baker & Duretto (2011) = Eriochilus cucullatus possibly reported in error | |
| | Gastrodia procera G.W.Carr, Indigenous Flora and Fauna Association Miscellaneous Paper 1: 22 (1991) | |
| | Gastrodia sesamoides R.Br., Prodr. Fl. Nov. Holland. 330 (1810) | 4a:127 |
| | Gastrodia surcula D.L.Jones, Orchadian 15: 554 (2008) | |
| | Genoplesium archeri (Hook.f.) D.L.Jones & M.A.Clem. = Corunastylis archeri | 4a:82 |
| | Genoplesium brachystachyum (Lindl.) D.L.Jones & M.A.Clem. = Corunastylis brachystachya | 4a:79 |
| | Genoplesium despectans (Hook.f.) D.L.Jones & M.A.Clem. = Corunastylis despectans | 4a:76 |
| | Genoplesium firthii (Cady) D.L.Jones = Corunastylis firthii | 4a:80 |
| | Genoplesium morrisii (Nicholls) D.L.Jones & M.A.Clem. = Corunastylis morrisii | 4a:83 |
| | Genoplesium nudiscapum (Hook.f.) D.L.Jones & M.A.Clem. = Corunastylis nudiscapa | 4a:79 |
| | Genoplesium nudum (Hook.f.) D.L.Jones & M.A.Clem. = Corunastylis nuda | 4a:82 |
| | Genoplesium pumilum (Hook.f.) D.L.Jones & M.A.Clem. = Corunastylis pumila | 4a:80 |
| | Genoplesium tasmanicum D.L.Jones = Corunastylis tasmanica | 4a:77 |
| | Glossodia major R.Br., Prodr. Fl. Nov. Holland. 326 (1810) | 4a:90 |
| | Gunnia australis Lindl. sensu Hooker (1860) = Sarcochilus australis | |

| | |
|---|--------|
| Hydrorchis orbicularis (R.S.Rogers) D.L.Jones & M.A.Clem., Orchadian 13: 462 (2002) | 4a:60 |
| Hymenochilus cycnocephalus (Fitzg.) D.L.Jones & M.A.Clem. sensu Buchanan (2005) = Pterostylis ziegeleri | 4a:26 |
| Hymenochilus muticus (R.Br.) D.L.Jones & M.A.Clem. = Pterostylis mutica | 4a:25 |
| Hymenochilus pratensis (D.L.Jones) D.L.Jones & M.A.Clem. = Pterostylis pratensis | |
| Hymenochilus rubenachii (D.L.Jones) D.L.Jones & M.A.Clem. = Pterostylis rubenachii | |
| Hymenochilus wapstrarum (D.L.Jones) D.L.Jones & M.A.Clem. = Pterostylis wapstrarum | |
| Hymenochilus ziegeleri (D.L.Jones) D.L.Jones & M.A.Clem. = Pterostylis ziegeleri | |
| Jonesiopsis filamentosa (R.Br.) D.L.Jones & M.A.Clem. = Caladenia filamentosa | 4a:99 |
| Leptoceras menziesii (R.Br.) Lindl., Gen. Sp. Orchid. Pl. 416 (1840) | 4a:99 |
| Linguella nana (R.Br.) D.L.Jones & M.A.Clem. = Pterostylis nana | 4a:14 |
| Lyperanthus nigricans R.Br. = Pyrorchis nigricans | 4a:93 |
| Lyperanthus suaveolens R.Br., Prodr. Fl. Nov. Holland. 325 (1810) | 4a:94 |
| Malaxis subulata Labill. = Cryptostylis subulata | |
| Microtidium atratum (Lindl.) D.L.Jones & M.A.Clem., Orchadian 13: 463 (2002) | 4a:60 |
| Microtis arenaria Lindl., Gen. Sp. Orchid. Pl. 396 (1840) | 4a:58 |
| Microtis atrata Lindl. = Microtidium atratum | 4a:60 |
| Microtis biloba Nicholls = Microtis arenaria | 4a:58 |
| Microtis oblonga R.S.Rogers, Trans. & Proc. Roy. Soc. South Australia 47: 339 (1923) | 4a:59 |
| Microtis orbicularis R.S.Rogers = Hydrorchis orbicularis | 4a:60 |
| Microtis parviflora R.Br., Prodr. Fl. Nov. Holland. 321 (1810) | 4a:59 |
| Microtis porrifolia (Sw.) R.Br. ex Spreng. var. parviflora (R.Br.) Rodway nom. illeg., sensu Rodway (1903) = Microtis parviflora | |
| Microtis pulchella R.Br. sensu Hooker (1860), misapplied to Microtis unifolia and M. arenaria | |
| Microtis rara R.Br. sensu Curtis (1979) = Microtis oblonga | 4a:59 |
| Microtis unifolia (G.Forst.) Rchb.f., Beitr. Syst. Pflanzenk. 62 (1871) | 4a:58 |
| Nemacianthus caudatus (R.Br.) D.L.Jones & M.A.Clem. = Acianthus caudatus | 4a:117 |
| Oligochaetochilus commutatus (D.L.Jones) Szlach. = Pterostylis commutata | |
| Oligochaetochilus pratensis (D.L.Jones) Szlach. = Pterostylis pratensis | |
| Oligochaetochilus rubenachii (D.L.Jones) Szlach. = Pterostylis rubenachii | |
| Oligochaetochilus squamatus (R.Br.) Szlach. = Pterostylis squamata | 4a:27 |
| Oligochaetochilus stenochilus (D.L.Jones) Szlach. = Pterostylis stenochila | |
| Oligochaetochilus wapstrarum (D.L.Jones) Szlach. = Pterostylis wapstrarum | |
| Oligochaetochilus williamsonii (D.L.Jones) Szlach. = Pterostylis williamsonii | |
| Oligochaetochilus ziegleri (D.L.Jones) Szlach. = Pterostylis ziegleri | |
| Orthoceras strictum R.Br., Prodr. Fl. Nov. Holland. 317 (1810) | 4a:30 |
| Paracaleana minor (R.Br.) Blaxell, Contr. New South Wales Natl. Herb. 4: 281 (1972) | 4a:88 |
| Petalochilus alatus (R.Br.) D.L.Jones & M.A.Clem. = Caladenia alata | |
| Petalochilus atrochilus (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia atrochila | |
| Petalochilus aurantiacus (R.S.Rogers) D.L.Jones & M.A.Clem. = Caladenia aurantiaca | |
| Petalochilus campbellii (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia campbellii | |
| Petalochilus carneus (R.Br.) D.L.Jones & M.A.Clem. = Caladenia carnea | 4a:106 |
| Petalochilus fuscatus (Rchb.f.) D.L.Jones & M.A.Clem. = Caladenia fuscata | |

| | | |
|---|---|--------|
| | Petalochilus mentiens (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia mentiens | |
| | Petalochilus prolatus (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia prolata | |
| | Petalochilus pusillus (W.M.Curtis) D.L.Jones & M.A.Clem. = Caladenia pusilla | 4a:107 |
| | Petalochilus sylvicola (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia sylvicola | |
| | Petalochilus tonellii (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia tonellii | |
| | Petalochilus vulgaris (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia vulgaris | |
| | Pheladenia deformis (R.Br.) D.L.Jones & M.A.Clem., Orchadian 13: 411 (2001) | 4a:113 |
| | Plumatichilos plumosum (Cady) Szlach. = Pterostylis plumosa | 4a:24 |
| | Plumatichilos tasmanicum (D.L.Jones) Szlach. = Pterostylis tasmanica | |
| | Prasophyllum abblittiorum P.A.Collier, Muelleria 36: 5 (2017) | |
| | Prasophyllum album R.S.Rogers sensu Curtis (1979) a name of uncertain application | 4a:69 |
| | Prasophyllum alpestre D.L.Jones = Prasophyllum mimulum (misapplied in Tasmania) | |
| e | Prasophyllum alpinum R.Br., Prodr. Fl. Nov. Holland. 318 (1810) | 4a:75 |
| e | Prasophyllum amoenum D.L.Jones, Austral. Orchid Res. 3: 99 (1998) | |
| e | Prasophyllum apoxychilum D.L.Jones, Austral. Orchid Res. 3: 100 (1998) | |
| | Prasophyllum archeri Hook.f. = Corunastylis archeri | 4a:82 |
| e | Prasophyllum sp. Arthurs Lake (R.Smith DLJ11363) Tas Herbarium | |
| e | Prasophyllum atratum D.L.Jones, Austral. Orchid Res. 5: 144 (2006) | |
| | Prasophyllum australe R.Br., Prodr. Fl. Nov. Holland. 318 (1810) | 4a:66 |
| | Prasophyllum beaugleholei Nicholls = Corunastylis nuda | 4a:82 |
| | Prasophyllum brachystachyum Lindl. = Corunastylis brachystachya | 4a:79 |
| | Prasophyllum brainei R.S.Rogers = Prasophyllum lindleyanum | 4a:74 |
| | Prasophyllum brevilabre (Lindl.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 11, t.110A (1858) | 4a:68 |
| | Prasophyllum buftonianum J.H.Willis = Corunastylis pumila | 4a:80 |
| e | Prasophyllum castaneum D.L.Jones, Austral. Orchid Res. 3: 102 (1998) | |
| e | Prasophyllum concinnum Nicholls, Vict. Naturalist 64: 232 (1948) | 4a:73 |
| | Prasophyllum correctum D.L.Jones = Prasophyllum incorrectum (misapplied in Tasmania) | |
| e | Prasophyllum crebriflorum D.L.Jones, Muelleria 18: 103 (2003) | |
| | Prasophyllum despectans Hook.f. = Corunastylis despectans | 4a:76 |
| | Prasophyllum elatum R.Br., Prodr. Fl. Nov. Holland. 318 (1810) | 4a:67 |
| e | Prasophyllum favonium D.L.Jones, Austral. Orchid Res. 3: 104 (1998) | |
| | Prasophyllum firthii Cady = Corunastylis firthii | 4a:80 |
| | Prasophyllum fitzgeraldii R.S.Rogers & Maiden recorded in error | |
| | Prasophyllum flavum R.Br., Prodr. Fl. Nov. Holland. 318 (1810) | 4a:66 |
| | Prasophyllum frenchii F.Muell. sensu Curtis (1979) a name of uncertain application | 4a:73 |
| | Prasophyllum fuscum R.Br. sensu Curtis (1979) a name of uncertain application | 4a:75 |
| | Prasophyllum gracile Lindl. sensu Curtis (1979) a name of uncertain application | 4a:72 |
| e | Prasophyllum incorrectum D.L.Jones, Muelleria 18: 107 (2003) | |
| e | Prasophyllum incurvum D.L.Jones, Austral. Orchid Res. 3: 106 (1998) | |
| | Prasophyllum intricatum C.Stuart ex Benth. = Corunastylis archeri | 4a:82 |
| e | Prasophyllum limnetes D.L.Jones, Austral. Orchid Res. 5: 151 (2006) | |
| | Prasophyllum lindleyanum Rchb.f., Beitr. Syst. Pflanzenk. 58 (1871) | 4a:74 |
| | Prasophyllum lutescens Lindl. sensu Hooker (1860) = Prasophyllum australe | |

| | | |
|---|---|-------|
| | Prasophyllum lutescens Lindl. var. brevilabre Lindl. = Prasophyllum brevilabre | |
| e | Prasophyllum milfordense D.L.Jones, Austral. Orchid Res. 3: 107 (1998) | |
| e | Prasophyllum mimulum D.L.Jones, Orchadian 14: 372 (2004) | 4a:69 |
| | Prasophyllum aff. montanum R.J.Bates & D.L.Jones sensu Buchanan (2009) = Prasophyllum sp. Arthurs Lake (R.Smith DLJI 1363) Tas Herbarium | |
| | Prasophyllum morrisii Nicholls = Corunastylis morrisii | 4a:83 |
| | Prasophyllum nigricans R.Br. sensu Curtis (1979) = Corunastylis tasmanica | 4a:77 |
| | Prasophyllum nudiscapum Hook.f. = Corunastylis nudiscapa | 4a:79 |
| | Prasophyllum nudum Hook.f. = Corunastylis nuda | |
| | Prasophyllum odoratum R.S.Rogers sensu Curtis (1979) a name of uncertain application | 4a:68 |
| e | Prasophyllum olidum D.L.Jones, Austral. Orchid Res. 3: 108 (1998) | |
| | Prasophyllum parviflorum (R.S.Rogers) Nicholls sensu Curtis (1979) a name of uncertain application | 4a:74 |
| | Prasophyllum patens R.Br. sensu Rodway (1903) = Prasophyllum truncatum | 4a:71 |
| | Prasophyllum patens R.Br. var. robustum Nicholls sensu Curtis (1953) = Prasophyllum robustum | |
| e | Prasophyllum perangustum D.L.Jones, Austral. Orchid Res. 3: 109 (1998) | 4a:72 |
| e | Prasophyllum pulchellum D.L.Jones, Austral. Orchid Res. 3: 110 (1998) | |
| | Prasophyllum pyriforme E.Coleman sensu Buchanan (2005) = Prasophyllum rostratum | |
| e | Prasophyllum robustum (Nicholls) M.A.Clem. & D.L.Jones, Austral. Orchid Res. 1: 117 (1989) | |
| | Prasophyllum rogersii Rupp sensu Curtis (1979) = Prasophyllum perangustum | 4a:72 |
| e | Prasophyllum rostratum Lindl., Gen. Sp. Orchid. Pl. 516 (1840) | |
| | Prasophyllum rufum R.Br. sensu Curtis (1979) = Corunastylis tasmanica | 4a:79 |
| e | Prasophyllum secutum D.L.Jones, Austral. Orchid Res. 3: 113 (1998) | |
| | Prasophyllum sphacelatum D.L.Jones, Muelleria 9: 59 (1996) | |
| e | Prasophyllum stellatum D.L.Jones, Austral. Orchid Res. 3: 115 (1998) | |
| | Prasophyllum suttonii R.S.Rogers & B.Rees sensu Curtis (1979) = Prasophyllum mimulum | 4a:69 |
| | Prasophyllum tadgellianum (R.S.Rogers) R.S.Rogers, Trans. & Proc. Roy. Soc. South Australia 47: 338 (1923) | |
| e | Prasophyllum taphanyx D.L.Jones, Orchadian 14: 373 (2004) | |
| e | Prasophyllum truncatum Lindl., Gen. Sp. Orchid. Pl. 513 (1840) | 4a:71 |
| e | Prasophyllum tunbridgense D.L.Jones, Austral. Orchid Res. 3: 117 (1998) | |
| | Prasophyllum uroglossum Rupp sensu Curtis (1979) a name of uncertain application | 4a:76 |
| | Pterostylis acuminata R.Br. sensu Curtis (1953) = Pterostylis × ingens | |
| e | Pterostylis alata (Labill.) Rchb.f., Beitr. Syst. Pflanzenk. 70 (1871) | 4a:20 |
| | Pterostylis alpina R.S.Rogers sensu Curtis (1979) = Pterostylis scabrada | 4a:19 |
| e | Pterostylis aphylla Lindl., Gen. Sp. Orchid. Pl. 392 (1840) | 4a:24 |
| | Pterostylis atrans D.L.Jones, Muelleria 8: 185 (1994) | 4a:22 |
| e | Pterostylis atriola D.L.Jones, Austral. Orchid Res. 3: 140 (1998) | |
| | Pterostylis barbata Lindl. sensu Rodway (1903) = Pterostylis plumosa | 4a:24 |
| e | Pterostylis commutata D.L.Jones, Muelleria 8: 186 (1994) | |
| | Pterostylis concinna R.Br., Prodr. Fl. Nov. Holland. 326 (1810) | 4a:11 |
| | Pterostylis cucullata R.Br. subsp. cucullata , Prodr. Fl. Nov. Holland. 327 (1810) | |
| | Pterostylis curta R.Br., Prodr. Fl. Nov. Holland. 326 (1810) | 4a:12 |
| | Pterostylis cycnocephala Fitzg. sensu Curtis (1979) = Pterostylis ziegeleri | 4a:26 |

| | | |
|---|--|--------|
| | Pterostylis decurva R.S.Rogers, Trans. & Proc. Roy. Soc. South Australia 47: 339, t.27 (1923) | 4a:22 |
| e | Pterostylis dubia R.Br., Prodr. Fl. Nov. Holland. 328 (1810) | 4a:16 |
| | Pterostylis falcata R.S.Rogers, Proc. Roy. Soc. Victoria 28: 106, t.9 (1915) | 4a:17 |
| | Pterostylis foliata Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 249 (1853) | 4a:19 |
| e | Pterostylis furcata Lindl., Gen. Sp. Orchid. Pl. 390 (1840) | 4a:17 |
| | Pterostylis grandiflora R.Br., Prodr. Fl. Nov. Holland. 327 (1810) | 4a:20 |
| | Pterostylis × ingens (Rupp) D.L.Jones, Orchadian 5: 54 (1976) | 4a:12 |
| | <i>Pterostylis longifolia</i> R.Br. sensu Curtis (1979) = Pterostylis melagramma , P. williamsonii , P. stenochila & P. tunstallii | 4a:27 |
| | Pterostylis lustra D.L.Jones, Austral. Orchid Res. 5: 87 (2006) | |
| | Pterostylis melagramma D.L.Jones, Austral. Orchid Res. 3: 145 (1998) | 4a:27 |
| | Pterostylis mutica R.Br., Prodr. Fl. Nov. Holland. 328 (1810) | 4a:25 |
| | Pterostylis nana R.Br., Prodr. Fl. Nov. Holland. 327 (1810) | 4a:14 |
| | Pterostylis nutans R.Br., Prodr. Fl. Nov. Holland. 327 (1810) | 4a:13 |
| | <i>Pterostylis obtusa</i> R.Br. sensu Curtis (1979) = Pterostylis atrans | 4a:22 |
| | Pterostylis parviflora R.Br., Prodr. Fl. Nov. Holland. 327 (1810) | 4a:23 |
| | Pterostylis pedoglossa Fitzg., Austral. Orchids 1(3): 5th plate (1877) | 4a:13 |
| | Pterostylis pedunculata R.Br., Prodr. Fl. Nov. Holland. 327 (1810) | 4a:14 |
| | Pterostylis plumosa Cady, Austral. Pl. 5: 138 (1969) | 4a:24 |
| | <i>Pterostylis praecox</i> Lindl. sensu Rodway (1903) = Pterostylis alata | 4a:20 |
| e | Pterostylis pratensis D.L.Jones, Austral. Orchid Res. 3: 149 (1998) | |
| | <i>Pterostylis pusilla</i> R.S.Rogers sensu Curtis (1953) = Pterostylis squamata | |
| e | Pterostylis rubenachii D.L.Jones, Austral. Orchid Res. 3: 150 (1998) | |
| | <i>Pterostylis rufa</i> R.Br. sensu Curtis (1979) = Pterostylis squamata | 4a:27 |
| | Pterostylis sanguinea D.L.Jones & M.A.Clem., Austral. Orchid Res. 1: 126 (1989) | 4a:28 |
| e | Pterostylis scabrida Lindl., Gen. Sp. Orchid. Pl. 389 (1840) | 4a:19 |
| | Pterostylis squamata R.Br., Prodr. Fl. Nov. Holland. 327 (1810) | 4a:27 |
| e | Pterostylis stenochila D.L.Jones, Austral. Orchid Res. 3: 153 (1998) | |
| | Pterostylis tasmanica D.L.Jones, Muellera 8: 190 (1994) | |
| | Pterostylis × toveyana Ewart & Sharman, Proc. Roy. Soc. Victoria 28: 235 (1916) | 4a:21 |
| | Pterostylis tunstallii D.L.Jones & M.A.Clem., Austral. Orchid Res. 1: 128 (1989) | |
| | Pterostylis uliginosa D.L.Jones, Austral. Orchid Res. 3: 155 (1998) | |
| | <i>Pterostylis vereenae</i> R.S.Rogers sensu Curtis (1953) = Pterostylis foliata | |
| | <i>Pterostylis vittata</i> Lindl. sensu Curtis (1979) = Pterostylis sanguinea | 4a:28 |
| e | Pterostylis wapstrarum D.L.Jones, Austral. Orchid Res. 3: 156 (1998) | |
| e | Pterostylis williamsonii D.L.Jones, Austral. Orchid Res. 3: 157 (1998) | 4a:27 |
| e | Pterostylis ziegelieri D.L.Jones, Austral. Orchid Res. 3: 158 (1998) | 4a:26 |
| | Pyrorchis nigricans (R.Br.) D.L.Jones & M.A.Clem., Phytologia 77: 449 (1994) | 4a:93 |
| | Sarcochilus australis (Lindl.) Rchb.f., Ann. Bot. Syst. (Walpers) 6: 501 (1863) | 4a:131 |
| | <i>Sarcochilus parviflorus</i> Lindl. sensu Rodway (1903) = Sarcochilus australis | 4a:131 |
| | <i>Simpliglottis cornuta</i> (Hook.f.) Szlach. = Chiloglottis cornuta | 4a:85 |
| | <i>Simpliglottis grammata</i> (G.W.Carr) Jeanes = Chiloglottis grammata | |
| | <i>Simpliglottis gunnii</i> (Lindl.) Szlach. = Chiloglottis gunnii | 4a:84 |

| | | |
|---|---|--------|
| | Simpliglottis triceratops (D.L.Jones) Jeanes = Chiloglottis triceratops | |
| | Simpliglottis valida (D.L.Jones) Szlach. = Chiloglottis valida | |
| | Specularantha aphylla (Lindl.) D.L.Jones & M.A.Clem. = Pterostylis aphylla | 4a:24 |
| | Specularantha atriola (D.L.Jones) D.L.Jones & M.A.Clem. = Pterostylis atriola | |
| | Specularantha parviflora (R.Br.) D.L.Jones & M.A.Clem. = Pterostylis parviflora | 4a:23 |
| | Specularantha uliginosa (D.L.Jones) D.L.Jones & M.A.Clem. = Pterostylis uliginosa | |
| | Spiculaea huntiana (F.Muell.) Schltr. = Thynninorchis huntiana | 4a:89 |
| | Spiranthes alticola D.L.Jones, Orchadian 15: 555 (2008) | |
| | Spiranthes australis (R.Br.) Lindl., Bot. Reg. 10: subt.823 (1824) | 4a:128 |
| | Spiranthes sinensis (Pers.) Ames subsp. australis (R.Br.) Kitam. = Spiranthes australis | 4a:128 |
| | Stegostyla alpina (R.S.Rogers) D.L.Jones & M.A.Clem. = Caladenia alpina | 4a:111 |
| | Stegostyla angustata (Lindl.) D.L.Jones & M.A.Clem. = Caladenia angustata | 4a:108 |
| | Stegostyla atrata (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia atrata | |
| | Stegostyla congesta (R.Br.) D.L.Jones & M.A.Clem. = Caladenia congesta | 4a:112 |
| | Stegostyla cracens (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia cracens | |
| | Stegostyla gracilis (R.Br.) D.L.Jones & M.A.Clem. = Caladenia gracilis | 4a:110 |
| | Stegostyla transitoria (D.L.Jones) D.L.Jones & M.A.Clem. = Caladenia transitoria | |
| | Sullivania minor (R.Br.) D.L.Jones & M.A.Clem. = Paracaleana minor | 4a:88 |
| | Taurantha concinna (R.Br.) D.L.Jones & M.A.Clem. = Pterostylis concinna | 4a:11 |
| e | Thelymitra aggericola D.L.Jones, The Orchadian 12: 517 (1999) | |
| | Thelymitra angustifolia R.Br. sensu Hooker (1860) a name of uncertain application | |
| | Thelymitra antennifera (Lindl.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 4, t.101A (1858) | 4a:51 |
| | Thelymitra arenaria Lindl., Gen. Sp. Orchid. Pl. 519 (1840) | |
| | Thelymitra aristata Lindl., Gen. Sp. Orchid. Pl. 521 (1840) | 4a:42 |
| | Thelymitra atronitida Jeanes, Muellera 14: 91 (2000) | |
| | Thelymitra azurea R.S.Rogers sensu Buchanan (1995) = Thelymitra jonesii | |
| | Thelymitra benthamiana Rchb.f., Beitr. Syst. Pflanzenk. 55 (1871) | |
| | Thelymitra bracteata J.Z.Weber ex Jeanes, Muellera 19: 43 (2004) | |
| | Thelymitra brevifolia Jeanes, Muellera 19: 30 (2004) | |
| | Thelymitra canaliculata R.Br. sensu Curtis (1979) = Thelymitra jonesii | 4a:46 |
| | Thelymitra carnea R.Br., Prodr. Fl. Nov. Holland. 314 (1810) | 4a:50 |
| | Thelymitra chasmogama R.S.Rogers sensu Curtis (1979) a plant of uncertain hybrid origin | 4a:47 |
| | Thelymitra circumsepta Fitzg., Austral. Orchids 1: t.1 (1878) | 4a:48 |
| | Thelymitra cyanea (Lindl.) Benth., Fl. Austral. 6: 323 (1873) | 4a:49 |
| | Thelymitra decora Cheeseman sensu Buchanan (1995) = Thelymitra simulata | |
| | Thelymitra erosa D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 184 (1998) | |
| | Thelymitra exigua Jeanes, Muellera 19: 28 (2004) | |
| | Thelymitra flexuosa Endl., Nov. Stirp. Dec. 3: 23 (1839) | 4a:51 |
| | Thelymitra grandiflora Fitzg. sensu Curtis (1953) = Thelymitra aristata | |
| | Thelymitra holmesii Nicholls, Vict. Naturalist 49: 263 (1933) | 4a:43 |
| e | Thelymitra imbricata D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 186 (1998) | 4a:43 |
| | Thelymitra improcera D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 187 (1998) | 4a:46 |
| | Thelymitra inflata Jeanes, Muellera 19: 71 (2004) | |

| | | |
|----------------------------|---|--------|
| | Thelymitra × irregularis Nicholls, Vict. Naturalist 63: 126 (1946) | 4a:48 |
| | Thelymitra ixioides Sw., Ksvenska Vet. Akad. Handl. 21: 253 (1800) | 4a:44 |
| e | Thelymitra jonesii Jeanes, Muellera 15: 81 (2001) | 4a:46 |
| | Thelymitra juncifolia Lindl., Gen. Sp. Orchid. Pl. 522 (1840) | |
| | <i>Thelymitra longifolia</i> J.R.Forst. & G.Forst. sensu Bentham (1873) = Thelymitra nuda | |
| | Thelymitra longiloba D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 191 (1998) | 4a:46 |
| | Thelymitra lucida Jeanes, Muellera 19: 70 (2004) | |
| | <i>Thelymitra luteocilium</i> Fitzg. sensu Curtis (1979) = Thelymitra rubra | 4a:47 |
| | <i>Thelymitra × macmillanii</i> F.Muell. sensu Curtis (1979) = a hybrid involving Thelymitra nuda | 4a:49 |
| | Thelymitra malvina M.A.Clem., D.L.Jones & Molloy, Austral. Orchid Res. 1: 141 (1989) | |
| | <i>Thelymitra media</i> R.Br. sensu Curtis (1979) = Thelymitra improcera & T. longiloba | 4a:46 |
| | <i>Thelymitra megcalyptra</i> Fitzg. sensu Curtis (1979) = Thelymitra imbricata | 4a:43 |
| | Thelymitra × merraniae Nicholls, Vict. Naturalist 46: 139 (1929) | |
| | Thelymitra mucida Fitzg., Gard. Chron. 17: 495 (1882) | 4a:46 |
| | Thelymitra nuda R.Br., Prodr. Fl. Nov. Holland. 314 (1810) | 4a:44 |
| | <i>Thelymitra nuda</i> R.Br. var. <i>grandiflora</i> Lindl. = Thelymitra imbricata | |
| | Thelymitra pauciflora R.Br., Prodr. Fl. Nov. Holland. 314 (1810) | 4a:42 |
| | <i>Thelymitra pauciflora</i> R.Br. var. <i>holmesii</i> (Nicholls) Nicholls = Thelymitra holmesii | 4a:43 |
| | Thelymitra peniculata Jeanes, Muellera 19: 50 (2004) | |
| e | Thelymitra polychroma D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 193 (1998) | |
| | <i>Thelymitra pulchella</i> Hook.f. recorded in error | |
| | <i>Thelymitra retecta</i> Rupp = Thelymitra circumsepta | 4a:48 |
| | Thelymitra rubra Fitzg., Gard. Chron. 17: 495 (1882) | 4a:50 |
| e | Thelymitra silena D.L.Jones, The Orchadian 12: 518 (1999) | |
| | Thelymitra simulata D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 195 (1998) | |
| | <i>Thelymitra smithiana</i> (Gunn ex Lindl.) Hook.f. = Thelymitra flexuosa | |
| e | Thelymitra spadicea D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 196 (1998) | |
| e | Thelymitra sparsa D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 197 (1998) | |
| | Thelymitra × truncata R.S.Rogers, Trans. & Proc. Roy. Soc. South Australia 41: 343, t.17 (1917) | 4a:45 |
| | <i>Thelymitra venosa</i> R.Br. sensu Curtis (1979) = Thelymitra cyanea | 4a:48 |
| | <i>Thelymitra versicolor</i> Lindl. = Thelymitra nuda | |
| e | Thelymitra viridis Jeanes, Muellera 19: 36 (2004) | |
| x | Thynninorchis huntiana (F.Muell.) D.L.Jones & M.A.Clem., Orchadian 13: 457 (2002) | 4a:89 |
| e | Thynninorchis nothofagicola (D.L.Jones) D.L.Jones & M.A.Clem., Orchadian 13: 457 (2002) | |
| t | Townsonia viridis (Hook.f.) Schltr., Repert. Spec. Nov. Regni Veg. 9: 250 (1911) | 4a:118 |
| | <i>Urochilus sanguineus</i> (D.L.Jones & M.A.Clem.) D.L.Jones & M.A.Clem. = Pterostylis sanguinea | 4a:28 |
| | ×Calassodia tutelata (R.S.Rogers) M.A.Clem., Austral. Orchid Res. 1: 33 (1989) | 4a:113 |
| | ×Glossadenia tutelata (R.S.Rogers) Kavulak nom. illeg. = ×Calassodia tutelata | |
| | ×Taurodium toveyanum (Ewart & Sharman) D.L.Jones & M.A.Clem. sensu Buchanan (2005) = Pterostylis toveyana | |
| POACEAE (GRAMINEAE) | | |
| | <i>Achnatherum caudatum</i> (Trin.) S.W.L.Jacobs & J.Everett = Amelichloa caudata | 4b:192 |
| | <i>Agropyron elongatum</i> (Host) P.Beauv. = Thinopyrum elongatum | 4b:295 |

| | | |
|---|---|--------|
| | Agropyron junceiforme (Á.Löve & D.Löve) Á.Löve & D.Löve = Thinopyrum junceiforme | 4b:293 |
| | Agropyron junceum (L.) P.Beauv. = Thinopyrum junceiforme | 4b:293 |
| | Agropyron pectinatum (Labill.) P.Beauv. = Australopyrum pectinatum | 4b:301 |
| | Agropyron repens (L.) P.Beauv. = Elytrigia repens | 4b:295 |
| | Agropyron scabrum (R.Br.) P.Beauv. = Anthosachne scabra | 4b:296 |
| | Agropyron velutinum Nees = Australopyrum velutinum | 4b:301 |
| | Agrostis aemula R.Br. = Lachnagrostis aemula | |
| | Agrostis aemula R.Br. var. aemula = Lachnagrostis aemula | 4b:258 |
| | Agrostis aemula R.Br. var. setifolia (Hook.f.) Vickery = Lachnagrostis punicea subsp. punicea | 4b:259 |
| | Agrostis aequata Nees = Lachnagrostis rudis | 4b:262 |
| | Agrostis australiensis Mez, Feddes Repert. Spec. Nov. Regni Veg. 17: 302 (1921) | 4b:257 |
| | Agrostis australiensis Mez sensu Curtis & Morris (1994) = Lachnagrostis rudis | 4b:257 |
| | Agrostis sp. aff. australiensis Mez sensu Curtis & Morris (1994) = Agrostis diemenica | 4b:257 |
| | Agrostis avenacea J.F.Gmel. = Lachnagrostis filiformis | 4b:259 |
| | Agrostis bettyae S.W.L.Jacobs, Telopea 9: 679 (2001) | |
| | Agrostis billardierei R.Br. var. billardierei = Lachnagrostis billardierei subsp. billardierei | 4b:260 |
| | Agrostis billardierei R.Br. var. collicola D.I.Morris = Lachnagrostis collicola | 4b:261 |
| | Agrostis billardierei R.Br. var. filifolia Vickery = Lachnagrostis punicea subsp. filifolia | 4b:261 |
| | Agrostis billardierei R.Br. var. robusta Vickery = Lachnagrostis robusta | 4b:261 |
| | Agrostis billardierei R.Br. var. tenuiseta D.I.Morris = Lachnagrostis billardierei subsp. tenuiseta | 4b:261 |
| i | Agrostis capillaris L. var. aristata (Parn.) Druce, Fl. Oxfordshire, ed. 2: 474 (1927) | 4b:264 |
| i | Agrostis capillaris L. var. capillaris , Sp. Pl. 1: 62 (1753) | 4b:264 |
| | Agrostis contracta F.Muell. ex Hook.f. = Deyeuxia contracta | |
| | Agrostis crinita (L.f.) R.Br. = Dichelachne crinita | |
| e | Agrostis diemenica D.I.Morris, Telopea 10: 765 (2004) | 4b:257 |
| i | Agrostis gigantea Roth, Tent. Fl. Germ. 1: 31 (1788) | 4b:264 |
| | Agrostis sp. aff. hiemalis (Walter) Britton, Sterns & Poggenb. sensu Curtis & Morris (1994) = Agrostis propinqua | 4b:258 |
| | Agrostis joyceae S.W.L.Jacobs, Telopea 9: 680 (2001) | |
| | Agrostis lacunarum D.I.Morris = Lachnagrostis lacunarum | 4b:260 |
| | Agrostis lobata R.Br. = Deyeuxia quadriseta | |
| | Agrostis montana R.Br. = Deyeuxia monticola | |
| | Agrostis muelleriana Vickery, Contr. New South Wales Natl. Herb. 1: 103 (1941) | 4b:256 |
| | Agrostis ovata G.Forst. sensu Labillardiere (1805) = Echinopogon ovatus | |
| | Agrostis parviflora R.Br., Prodr. Fl. Nov. Holland. 170 (1810) | 4b:256 |
| | Agrostis sp. aff. parviflora R.Br. sensu Curtis & Morris (1994) = Agrostis thompsoniae | 4b:257 |
| | Agrostis propinqua S.W.L.Jacobs, Telopea 9: 681 (2001) | 4b:258 |
| | Agrostis quadrifida Labill. = Pentapogon quadrifidus | |
| | Agrostis quadriseta (Labill.) R.Br. = Deyeuxia quadriseta | |
| | Agrostis rudis Roem. & Schult. sensu Curtis & Morris (1994) = Lachnagrostis morrisii | 4b:261 |
| | Agrostis scabra R.Br. sensu Rodway (1903) = Agrostis parviflora & A. propinqua | |
| | Agrostis sp. aff. scabra R.Br. sensu Curtis & Morris (1994) = Agrostis propinqua | 4b:258 |
| | Agrostis sciurea R.Br. = Dichelachne micrantha | |

| | | |
|------|--|--------|
| | <i>Agrostis semiverticillata</i> (Forssk.) C.Chr. = Agrostis viridis | 4b:283 |
| i | Agrostis stolonifera L., Sp. Pl. 1: 62 (1753) | 4b:262 |
| | <i>Agrostis tenuis</i> Sibth. = Agrostis capillaris | 4b:264 |
| | Agrostis thompsoniae S.W.L.Jacobs, Telopea 9: 682 (2001) | 4b:257 |
| | Agrostis venusta Trin., Mém. Acad. Imp. Sci. St.-Pétersbourg, Ser. 6 Sci. Math., Seconde Pt. Sci. Nat. 4: 340 (1841) | 4b:256 |
| | <i>Agrostis virginica</i> L. sensu Labillardiere (1805) = Sporobolus virginicus | |
| i | Agrostis viridis Gouan, Hortus Monsp. 546 (1762) | 4b:283 |
| | <i>Agrostis vulgaris</i> With. sensu Spicer (1878) = Agrostis capillaris | 4b:264 |
| i | Aira caryophyllea L. subsp. caryophyllea , Sp. Pl. 1: 66 (1753) | 4b:246 |
| i * | <i>Aira cupaniana</i> Guss., Fl. Sicul. Syn. 1: 148 (1843) | |
| | <i>Aira elegans</i> Willd. ex Gaudin = Aira elegantissima | 4b:247 |
| i | Aira elegantissima Schur, Verh. Mitth. Siebenbürg. Vereins Naturwiss. Hermannstadt 4: 85 (1853) | 4b:247 |
| i | Aira praecox L., Sp. Pl. 1: 65 (1753) | 4b:246 |
| | <i>Alopecurus agrestis</i> L. = Alopecurus myosuroides | |
| i | Alopecurus geniculatus L., Sp. Pl. 1: 60 (1753) | 4b:284 |
| i | Alopecurus myosuroides Huds., Fl. Angl. (Hudson): 23 (1762) | 4b:285 |
| i | Alopecurus pratensis L. subsp. pratensis , Sp. Pl. 1: 60 (1753) | 4b:284 |
| i | Amelichloa caudata (Trin.) Arriaga & Barkworth, Sida 22: 148 (2006) | 4b:192 |
| i | Ammophila arenaria (L.) Link subsp. arenaria , Hort. Berol. [Link] 1: 105 (1827) | 4b:274 |
| | Amphibromus archeri (Hook.f.) P.Morris, Vict. Naturalist 51: 146 (1934) | 4b:232 |
| ?i # | Amphibromus fluitans Kirk, Trans. & Proc. New Zealand Inst. 16: 374 (1884) | |
| | Amphibromus macrorhinus S.W.L.Jacobs & Lapinpuro, Telopea 2: 723 (1986) | 4b:233 |
| | Amphibromus neesii Steud., Syn. Pl. Glumac. 1: 328 (1854) | 4b:234 |
| | Amphibromus nervosus (Hook.f.) Baill., Hist. Pl. 12: 203 (1894) | |
| | Amphibromus recurvatus Swallen, Amer. J. Bot. 18: 415 (1931) | 4b:232 |
| | Amphibromus sinuatus S.W.L.Jacobs & Lapinpuro, Telopea 2: 727 (1986) | 4b:233 |
| | <i>Anthistiria australis</i> R.Br. = Themeda triandra | |
| | <i>Anthistiria ciliata</i> L.f. sensu Rodway (1903) = Themeda triandra | 4b:358 |
| i | Anthosachne kingiana (Endl.) Govaerts subsp. multiflora (Banks & Sol. ex Hook.f.) Govaerts, J. Adelaide Bot. Gard. 27: 24 (2014) | |
| | Anthosachne scabra (R.Br.) Nevski, Trudy Sredne-Aziatsk. Gosud. Univ., Ser. 8b, Bot. 17: 65 (1934) | 4b:296 |
| | <i>Anthoxanthum crinitum</i> L.f. sensu Brown (1810) = Dichelachne crinita | |
| i | Anthoxanthum odoratum L., Sp. Pl. 1: 28 (1753) | 4b:250 |
| i | Aristida benthamii Henrard var. benthamii , Meded. Rijks-Herb. 58A: 246, t.117 (1932) | 4b:323 |
| | <i>Aristida jerichoensis</i> (Domin) Henrard sensu Buchanan et al. (1989) = Aristida benthamii var. benthamii | 4b:323 |
| | <i>Arrhenatherum avenaceum</i> (Scop.) P.Beauv. sensu Rodway (1903) = Arrhenatherum elatius | 4b:234 |
| i | Arrhenatherum elatius (L.) P.Beauv. ex J.Presl & C.Presl var. bulbosum (Willd.) Spenn., Fl. Friburg. 1: 113 (1825) | 4b:234 |
| | <i>Arundo penicillatum</i> Labill. = Rytidosperma penicillatum | |
| | <i>Arundo phragmites</i> L. sensu Brown (1810) = Phragmites australis | |
| | <i>Arundo poiformis</i> Labill. sensu Labillardiere (1805) = Poa poiformis | |

| | | |
|-----|--|--------|
| | Arundo semiannularis Labill. = Rytidosperma semiannulare | |
| | Australopyrum pectinatum (Labill.) Á.Löve, Feddes Repert. 95: 443 (1984) | 4b:301 |
| | Australopyrum velutinum (Nees) B.K.Simon, Austrobaileya 2: 241 (1986) | 4b:301 |
| | Austrodanthonia alpicola (Vickery) H.P.Linder sensu Fl. Australia 44B: 51 = Rytidosperma oreophilum (Tasmanian plants) | |
| | Austrodanthonia caespitosa (Gaudich.) H.P.Linder = Rytidosperma caespitosum | 4b:312 |
| | Austrodanthonia carphoides (F.Muell. ex Benth.) H.P.Linder = Rytidosperma carphoides | 4b:306 |
| | Austrodanthonia diemenica (D.I.Morris) H.P.Linder = Rytidosperma diemenicum | 4b:311 |
| | Austrodanthonia geniculata (J.M.Black) H.P.Linder = Rytidosperma geniculatum | 4b:308 |
| | Austrodanthonia induta (Vickery) H.P.Linder = Rytidosperma indutum | 4b:312 |
| | Austrodanthonia laevis (Vickery) H.P.Linder = Rytidosperma laevis | 4b:312 |
| | Austrodanthonia penicillata (Labill.) H.P.Linder = Rytidosperma penicillatum | 4b:313 |
| | Austrodanthonia pilosa (R.Br.) H.P.Linder = Rytidosperma pilosum | 4b:313 |
| | Austrodanthonia popinensis (D.I.Morris) H.P.Linder = Rytidosperma fulvum | 4b:316 |
| | Austrodanthonia procera (Vickery) S.W.L.Jacobs = Rytidosperma indutum | 4b:312 |
| | Austrodanthonia racemosa (R.Br.) H.P.Linder = Rytidosperma racemosum | 4b:314 |
| | Austrodanthonia racemosa (R.Br.) H.P.Linder var. racemosa = Rytidosperma racemosum var. racemosum | 4b:314 |
| | Austrodanthonia remota (D.I.Morris) H.P.Linder = Rytidosperma remotum | 4b:316 |
| | Austrodanthonia setacea (R.Br.) H.P.Linder = Rytidosperma setaceum | 4b:311 |
| | Austrodanthonia tenuior (Steud.) H.P.Linder = Rytidosperma tenuius | 4b:310 |
| i t | Austroderia richardii (Endl.) N.P.Barker & H.P.Linder, Ann. Missouri Bot. Gard. 97: 344 (2010) | 4b:319 |
| | Austrofestuca hookeriana (F.Muell. ex Hook.f.) S.W.L.Jacobs = Hookerchloa hookeriana | 4b:223 |
| | Austrofestuca littoralis (Labill.) E.B.Alexeev, Byull. Moskovsk. Obshch. Isp. Prir., Otd. Biol. 81(5): 55 (1976) | 4b:223 |
| e | Austrostipa aphylla (Rodway) S.W.L.Jacobs & J.Everett, Telopea 6: 584 (1996) | 4b:192 |
| | Austrostipa bigeniculata (Hughes) S.W.L.Jacobs & J.Everett, Telopea 6: 584 (1996) | 4b:190 |
| | Austrostipa blackii (C.E.Hubb.) S.W.L.Jacobs & J.Everett, Telopea 6: 584 (1996) | 4b:191 |
| | Austrostipa flavescens (Labill.) S.W.L.Jacobs & J.Everett, Telopea 6: 585 (1996) | 4b:190 |
| | Austrostipa mollis (R.Br.) S.W.L.Jacobs & J.Everett, Telopea 6: 587 (1996) | 4b:189 |
| | Austrostipa nodosa (S.T.Blake) S.W.L.Jacobs & J.Everett, Telopea 6: 587 (1996) | 4b:188 |
| | Austrostipa pubinodis (Trin. & Rupr.) S.W.L.Jacobs & J.Everett, Telopea 6: 588 (1996) | 4b:187 |
| | Austrostipa rudis (Spreng.) S.W.L.Jacobs & J.Everett subsp. australis (J.Everett & S.W.L.Jacobs) S.W.L.Jacobs & J.Everett, Telopea 6: 588 (1996) | 4b:186 |
| | Austrostipa scabra (Lindl.) S.W.L.Jacobs & J.Everett subsp. falcata (Hughes) S.W.L.Jacobs & J.Everett, Telopea 6: 588 (1996) | 4b:188 |
| | Austrostipa scabra (Lindl.) S.W.L.Jacobs & J.Everett subsp. scabra , Telopea 6: 588 (1996) | 4b:188 |
| | Austrostipa semibarbata (R.Br.) S.W.L.Jacobs & J.Everett, Telopea 6: 588 (1996) | 4b:189 |
| | Austrostipa stipoides (Hook.f.) S.W.L.Jacobs & J.Everett, Telopea 6: 589 (1996) | 4b:186 |
| | Austrostipa stuposa (Hughes) S.W.L.Jacobs & J.Everett, Telopea 6: 589 (1996) | 4b:191 |
| i * | Avellinia michelii (Savi) Parl., Pl. Nov. 61 (1842) | |
| | Avena alba Vahl sensu Townrow (1969) = Avena barbata | 4b:236 |
| i | Avena barbata Pott ex Link, J. Bot. (Shrader) 2: 314,315 (1799) | 4b:236 |

| | | |
|-----|--|--------|
| i | Avena fatua L., Sp. Pl. 1: 80 (1753) | 4b:237 |
| | <i>Avena filiformis</i> G.Forst. sensu Labillardiere (1805) = Lachnagrostis filiformis | |
| i | Avena ludoviciana Durieu, Actes Soc. Linn. Bordeaux 20: 41 (1855) | 4b:238 |
| | <i>Avena quadriseta</i> Labill. = Deyeuxia quadriseta | |
| i | Avena sativa L., Sp. Pl. 1: 79 (1753) | 4b:237 |
| | <i>Avena sterilis</i> L. subsp. <i>ludoviciana</i> (Durieu) Gillet & Magne = Avena ludoviciana | 4b:238 |
| i | Avena strigosa Schreb., Spic. Fl. Lips. 52 (1771) | 4b:237 |
| i | Bothriochloa macra (Steud.) S.T.Blake, Proc. Roy. Soc. Queensland 80: 64 (1969) | 4b:355 |
| i | Brachypodium distachyon (L.) P.Beauv., Ess. Agrostogr. 101, 155, 156 (1812) | 4b:292 |
| i | Briza maxima L., Sp. Pl. 1: 70 (1753) | 4b:208 |
| i | Briza minor L., Sp. Pl. 1: 70 (1753) | 4b:208 |
| i | Bromus alopecuroides Poir., Voy. Barbarie 2: 100 (1789) | 4b:288 |
| | Bromus arenarius Labill., Nov. Holl. Pl. 1: 23, t.28 (1805) | 4b:288 |
| i | Bromus brevis Nees ex Steud., Syn. Pl. Glumac. 1: 326 (1854) | 4b:290 |
| i | Bromus catharticus Vahl, Symb. Bot. (Vahl) 2: 22 (1791) | 4b:290 |
| i | Bromus cecadilla Steud., Syn. Pl. Glumac. 1: 321 (1854) | 4b:290 |
| i | Bromus diandrus Roth, Bot. Abh. Beobacht. 44 (1787) | 4b:288 |
| | <i>Bromus fonkii</i> Phil. = Bromus lithobius | 4b:290 |
| i | Bromus hordeaceus L., Sp. Pl. 1: 77 (1753) | 4b:287 |
| | <i>Bromus japonicus</i> Murray var. <i>vestitus</i> sensu Buchanan (2005) = Bromus arenarius | 4b:288 |
| i | Bromus lithobius Trin., Linnaea 10: 303 (1836) | 4b:290 |
| | <i>Bromus macrostachys</i> Desf. sensu Townrow (1969) = Bromus alopecuroides | 4b:288 |
| i | Bromus madritensis L. var. ciliatus Guss., Fl. Sicul. Syn. 1: 78 (1843) | 4b:289 |
| | <i>Bromus mollis</i> L. = Bromus hordeaceus | 4b:287 |
| i | Bromus sterilis L., Sp. Pl. 1: 77 (1753) | 4b:289 |
| | <i>Bromus thominii</i> Hardouin sensu Townrow (1969) = Bromus hordeaceus | 4b:287 |
| | <i>Bromus unioloides</i> Kunth = Bromus catharticus | 4b:290 |
| | <i>Bromus willdenowii</i> Kunth = Bromus catharticus | 4b:290 |
| i x | Calamagrostis epigejos (L.) Roth, Tent. Fl. Germ. 1: 34 (1788) | 4b:273 |
| i | Catapodium marinum (L.) C.E.Hubb., Kew Bull. 9: 375 (1954) | 4b:225 |
| i | Catapodium rigidum (L.) C.E.Hubb., Fl. Bedfordshire: 437 (1953) | 4b:224 |
| i | Cenchrus clandestinus (Hochst. ex Chiov.) Morrone, Ann. Bot. (Oxford) 106(1): 127 (2010) | 4b:348 |
| i | Cenchrus longisetus M.C.Johnst., Sida 1(3): 182 (1963) | 4b:349 |
| i | Cenchrus macrourus (Trin.) Morrone, Ann. Bot. (Oxford) 106(1): 128 (2010) | 4b:351 |
| | <i>Cenchrus purpurascens</i> Thunb. recorded in error | 4b:351 |
| | <i>Ceratochloa unioloides</i> DC. = Bromus catharticus | 4b:290 |
| i | Chloris truncata R.Br., Prodr. Fl. Nov. Holland. 186 (1810) | |
| i | Cortaderia jubata (Lemoine) Stapf, Bot. Mag. 124, t.7607 (1898) | 4b:320 |
| | <i>Cortaderia richardii</i> (Endl.) Zotov = Austroderia richardii | 4b:319 |
| i | Cortaderia selloana (Schult. & Schult.f.) Asch. & Graebn., Syn. Mitteleur. Fl. [Ascherson & Graebner] 2: 325 (1900) | 4b:320 |
| | <i>Critesion marinum</i> (Huds.) Á.Löve = Hordeum marinum | 4b:300 |
| | <i>Critesion murinum</i> (L.) Á.Löve = Hordeum murinum | 4b:298 |

| | | |
|---|---|--------|
| i | Cynodon dactylon (L.) Pers. var. dactylon , Syn. Pl. (Persoon) 1: 85 (1805) | 4b:330 |
| i | Cynosurus cristatus L., Sp. Pl. 1: 72 (1753) | 4b:207 |
| i | Cynosurus echinatus L., Sp. Pl. 1: 72 (1753) | 4b:207 |
| i | Dactylis glomerata L., Sp. Pl. 1: 71 (1753) | 4b:221 |
| | Danthonia archeri Hook.f. = Amphibromus archeri | |
| | Danthonia caespitosa Gaudich. = Rytidosperma caespitosum | 4b:312 |
| | Danthonia carphoides F.Muell. ex Benth. var. angustior Vickery = Rytidosperma carphoides | 4b:306 |
| i | Danthonia decumbens (L.) DC., Lam., Fl. Franc. (DC. & Lamarck), ed. 3. 3: 33 (1805) | 4b:317 |
| | Danthonia diemenica D.I.Morris = Rytidosperma diemenicum | 4b:311 |
| | Danthonia dimidiata Vickery = Rytidosperma dimidiatum | 4b:317 |
| | Danthonia eriantha Lindl. sensu Townrow (1969) a name of uncertain application | |
| | Danthonia fortunae-hibernae Renvoize = Rytidosperma fortunae-hibernae | 4b:310 |
| | Danthonia geniculata J.M.Black = Rytidosperma geniculatum | 4b:308 |
| | Danthonia gracilis Hook.f. = Rytidosperma gracile | 4b:309 |
| | Danthonia laevis Vickery = Rytidosperma laeve | 4b:312 |
| | Danthonia longifolia R.Br. sensu Townrow (1969) = Rytidosperma tenuius | 4b:310 |
| | Danthonia nervosa Hook.f. = Amphibromus nervosus | |
| | Danthonia nitens D.I.Morris = Rytidosperma nitens | 4b:315 |
| | Danthonia nivicola Vickery = Rytidosperma nivicola | 4b:315 |
| | Danthonia nudiflora P.Morris = Rytidosperma nudiflorum | 4b:314 |
| | Danthonia pallida R.Br. sensu Bentham (1878) = <i>Rytidosperma pallidum</i> | |
| | Danthonia pauciflora R.Br. = Rytidosperma pauciflorum | 4b:309 |
| | Danthonia penicillata (Labill.) R.Br. ex P.Beauv. = Rytidosperma penicillatum | 4b:313 |
| | Danthonia penicillata (Labill.) R.Br. ex P.Beauv. var. pallida (R.Br.) Rodway sensu Rodway (1903) = <i>Rytidosperma pallidum</i> | |
| | Danthonia penicillata (Labill.) R.Br. ex P.Beauv. var. pilosa (R.Br.) Rodway = Rytidosperma pilosum | |
| | Danthonia penicillata (Labill.) R.Br. ex P.Beauv. var. racemosa (R.Br.) Rodway = Rytidosperma racemosum | |
| | Danthonia penicillata (Labill.) R.Br. ex P.Beauv. var. semiannularis (Labill.) Rodway = Rytidosperma semiannulare | |
| | Danthonia penicillata (Labill.) R.Br. ex P.Beauv. var. setacea (R.Br.) Rodway = Rytidosperma setaceum | |
| | Danthonia pilosa R.Br. = Rytidosperma pilosum | 4b:313 |
| | Danthonia popinensis D.I.Morris = Rytidosperma fulvum | 4b:316 |
| | Danthonia procera Vickery = Rytidosperma indutum | 4b:312 |
| | Danthonia pulvinorum D.I.Morris = Rytidosperma nivicola | 4b:315 |
| | Danthonia purpurascens Vickery sensu Townrow (1969) = Rytidosperma tenuius | 4b:310 |
| | Danthonia racemosa R.Br. = Rytidosperma racemosum | 4b:314 |
| | Danthonia racemosa R.Br. var. penicillata (Labill.) Benth. nom. illeg. = Rytidosperma penicillatum | |
| | Danthonia remota D.I.Morris = Rytidosperma remotum | 4b:316 |
| | Danthonia semiannularis (Labill.) R.Br. var. gracilis (Hook.f.) Hook.f. = Rytidosperma gracile | 4b:308 |
| | Danthonia semiannularis (Labill.) R.Br. var. semiannularis = Rytidosperma semiannulare | 4b:309 |
| | Danthonia setacea R.Br. = Rytidosperma setaceum | 4b:311 |

| | | |
|-----|--|--------|
| | Danthonia subulata Hook.f. = Rytidosperma setaceum | |
| | Danthonia tenuior (Steud.) Conert = Rytidosperma tenuius | 4b:310 |
| | Deschampsia cespitosa (L.) P.Beauv., Ess. Agrostogr. 91: 160, t.18, fig.3 (1812) | 4b:242 |
| t | Deschampsia gracillima Kirk, J. Bot. 29: 237 (1891) | 4b:243 |
| | Deyeuxia accedens Vickery = Deyeuxia frigida | 4b:271 |
| | Deyeuxia aequata (Nees) Benth. = Lachnagrostis rudis | 4b:261 |
| e | Deyeuxia apsleyensis D.I.Morris, Muelleria 7: 160 (1990) | 4b:273 |
| | Deyeuxia benthamiana Vickery = Deyeuxia scaberula | 4b:270 |
| | Deyeuxia billardierei (R.Br.) Kunth = Lachnagrostis billardierei | 4b:260 |
| | Deyeuxia brachyathera (Stapf) Vickery, Contr. New South Wales Natl. Herb. 1: 68 (1940) | 4b:272 |
| | Deyeuxia carinata Vickery, Contr. New South Wales Natl. Herb. 1: 58 (1940) | 4b:272 |
| | Deyeuxia contracta (F.Muell. ex Hook.f.) Vickery, Contr. New South Wales Natl. Herb. 1: 65 (1940) | 4b:269 |
| | Deyeuxia decipiens (R.Br.) Vickery, Contr. New South Wales Natl. Herb. 1: 70 (1940) | 4b:269 |
| | Deyeuxia densa Benth., Fl. Austral. 7: 582 (1878) | 4b:270 |
| | Deyeuxia forsteri Kunth = Lachnagrostis filiformis | 4b:259 |
| | Deyeuxia frigida F.Muell. ex Benth., Fl. Austral. 7: 583 (1878) | 4b:271 |
| | Deyeuxia gunniana (Nees) Benth., Fl. Austral. 7: 584 (1878) | 4b:267 |
| | Deyeuxia innominata D.I.Morris, Muelleria 7: 164 (1990) | 4b:268 |
| e x | Deyeuxia lawrencei Vickery, Contr. New South Wales Natl. Herb. 1: 48 (1940) | 4b:266 |
| | Deyeuxia minor F.Muell. ex Benth., Fl. Austral. 7: 582 (1878) | 4b:270 |
| | Deyeuxia montana (R.Br.) Benth. = Deyeuxia monticola | 4b:268 |
| | Deyeuxia monticola (Roem. & Schult.) Vickery, Contr. New South Wales Natl. Herb. 1: 56 (1940) | 4b:268 |
| | Deyeuxia parviseta Vickery sensu Townrow (1969) = Deyeuxia innominata | 4b:268 |
| | Deyeuxia quadriseta (Labill.) Benth., Fl. Austral. 7: 581 (1878) | 4b:267 |
| | Deyeuxia rodwayi Vickery, Contr. New South Wales Natl. Herb. 1: 60 (1940) | 4b:271 |
| | Deyeuxia scaberula Vickery, Contr. New South Wales Natl. Herb. 1: 64 (1940) | 4b:270 |
| | Deyeuxia scabra Kunth sensu Rodway (1903) = Deyeuxia contracta & D. scaberula | |
| | Dichanthium sericeum (R.Br.) A.Camus subsp. sericeum previously listed as naturalised but insufficient evidence exists to support this | 4b:355 |
| | Dichelachne crinita (L.f.) Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 293 (1853) | 4b:277 |
| | Dichelachne inaequiglumis (Hack. ex Cheeseman) Edgar & Connor, New Zealand J. Bot. 20: 307 (1982) | 4b:278 |
| | Dichelachne micrantha (Cav.) Domin, Biblioth. Bot. 20(85): 353 (1915) | 4b:278 |
| | Dichelachne montana Endl., Prodr. Fl. Norfolk. 20 (1833) | |
| | Dichelachne parva B.K.Simon, Austrobaileya 1: 462 (1982) | |
| | Dichelachne rara (R.Br.) Vickery, Contr. New South Wales Natl. Herb. 1: 337 (1951) | 4b:279 |
| | Dichelachne sciurea (R.Br.) Hook.f. sensu Rodway (1903) = Dichelachne spp. excluding Dichelachne crinita | |
| | Dichelachne sieberiana Trin. & Rupr., Sp. Gram. Stipac. 2–3 (1842) | |
| | Dichelachne stipoides Hook.f. = Austrostipa stipoides | |
| | Digitaria ciliaris (Retz.) Koeler previously listed as naturalised but insufficient evidence exists to support this | 4b:347 |
| i | Digitaria sanguinalis (L.) Scop., Fl. Carniol., ed. 2. 2: 52 (1772) | 4b:346 |

| | | |
|-----|--|--------|
| | Digitaria ternata (Hochst. ex A.Rich.) Stapf previously listed as naturalised but insufficient evidence exists to support this | 4b:346 |
| | Diplax tasmanica Hook.f. = Microlaena tasmanica | |
| | Disarrenum antarcticum Labill. = Hierochloe redolens | |
| | Distichlis distichophylla (Labill.) Fassett, Rhodora 27: 71 (1925) | 4b:324 |
| | Distichlis maritima Raf. sensu Rodway (1903) = Distichlis distichophylla | 4b:324 |
| | Dryopoa dives (F.Muell.) Vickery, Contr. New South Wales Natl. Herb. 3: 196 (1963) | 4b:199 |
| i | Echinochloa crus-galli (L.) P.Beauv., Ess. Agrostogr. 53 (1812) | 4b:336 |
| | Echinochloa crus-galli (L.) P.Beauv. var. frumentacea W.Wright = Echinochloa frumentacea | 4b:338 |
| i | Echinochloa esculenta (A.Braun) H.Scholz, Taxon 41: 523 (1992) | 4b:336 |
| i | Echinochloa frumentacea Link, Hort. Berol. [Link] 1: 204 (1827) | 4b:338 |
| | Echinochloa oryzoides (Ard.) Fritsch previously listed as naturalised but insufficient evidence exists to support this | |
| | Echinochloa utilis Ohwi & Yabuno = Echinochloa esculenta | 4b:336 |
| | Echinopogon ovatus (G.Forst.) P.Beauv., Ess. Agrostogr. 42, 161, t.9 fig.5 (1812) | 4b:275 |
| | Ehrharta acuminata (R.Br.) Spreng. = Tetrarrhena acuminata | 4b:182 |
| i | Ehrharta calycina Sm., Pl. Icon. Ined. 2: t.33 (1790) | 4b:178 |
| | Ehrharta distichophylla Labill. = Tetrarrhena distichophylla | 4b:183 |
| i | Ehrharta erecta Lam. var. erecta , Encycl. (Lamarck) 2: 347 (1786) | 4b:179 |
| | Ehrharta juncea (R.Br.) Spreng. = Tetrarrhena juncea | 4b:183 |
| i | Ehrharta longiflora Sm., Pl. Icon. Ined. 2: t.32 (1790) | 4b:179 |
| | Ehrharta oreophila (D.I.Morris) L.P.M.Willemse var. minor (D.I.Morris) L.P.M.Willemse = Tetrarrhena oreophila var. minor | 4b:182 |
| | Ehrharta oreophila (D.I.Morris) L.P.M.Willemse var. oreophila = Tetrarrhena oreophila var. oreophila | 4b:182 |
| | Ehrharta stipoides Labill. = Microlaena stipoides | 4b:181 |
| | Ehrharta tasmanica (Hook.f.) L.P.M.Willemse var. subalpina (F.Muell. ex Benth.) L.P.M.Willemse = Microlaena tasmanica var. subalpina | 4b:182 |
| | Ehrharta tasmanica (Hook.f.) L.P.M.Willemse var. tasmanica = Microlaena tasmanica var. tasmanica | 4b:181 |
| i | Ehrharta villosa Schult.f., Syst. Veg., ed. 15 bis [Roemer & Schultes] 7: 1374 (1830) | 4b:184 |
| i # | Eleusine indica (L.) Gaertn., Fruct. Sem. Pl. 1: 8 (1788) | |
| i | Eleusine tristachya (Lam.) Lam., Tabl. Encycl. 1: 57 (1791) | |
| | Elymus elongatus (Host) Runemark = Thinopyrum elongatum | 4b:295 |
| | Elymus farctus (Viv.) Runemark ex Melderis subsp. boreali-atlanticus (Simonet & Guin.) Melderis = Thinopyrum junceiforme | |
| | Elymus multiflorus (Banks & Sol. ex Hook.f.) Á.Löve & Connor = Anthosachne kingiana subsp. multiflora | |
| | Elymus repens (L.) Gould = Elytrigia repens | 4b:295 |
| | Elymus scaber (R.Br.) Á.Löve = Anthosachne scabra | 4b:296 |
| i | Elytrigia repens (L.) Desv. ex Nevski, Trudy Bot. Inst. Akad. Nauk S.S.S.R., Ser. I, Fl. Sist. Vyssh. Rast. 1: 14 (1933) | 4b:295 |
| i | Enneapogon nigricans (R.Br.) P.Beauv., Ess. Agrostogr. 82, 161, 171 (1812) | |
| ?i | Eragrostis brownii (Kunth) Nees, Cat. Ind. Pl. 2: 105 (1834) | 4b:326 |
| i | Eragrostis cilianensis (All.) Vignolo ex Janch., Mitt. Naturwiss. Vereins Univ. Wien 5(9): 110 (1907) | 4b:326 |

| | | |
|-------|---|--------|
| i | Eragrostis curvula (Schrad.) Nees, Fl. Afr. Austral. III. 397 (1841) | 4b:327 |
| i | Eragrostis minor Host, Icon. Descr. Gram. Austriac. 4: 15 (1809) | |
| | <i>Eragrostis molybdea</i> Vickery = Eragrostis brownii | 4b:326 |
| i | Eragrostis parviflora (R.Br.) Trin., Mém. Acad. Imp. Sci. St.Pétersbourg Hist. Acad. II: 411 (1830) | |
| i | Eragrostis pilosa (L.) P.Beauv., Ess. Agrostogr. 71, 162, 175 (1812) | |
| n i # | <i>Eragrostis tenuifolia</i> (A.Rich.) Hochst. ex Steud., Syn. Pl. Glumac. I: 268 (1854) | 4b:327 |
| | <i>Erythranthera australis</i> (Petrie) Zotov = Rytidosperma australe | 4b:318 |
| | <i>Festuca archeri</i> E.B.Alexeev = Festuca arundinacea | 4b:428 |
| i | Festuca arundinacea Schreb., Spic. Fl. Lips. 57 (1771) | 4b:197 |
| | <i>Festuca asperula</i> Vickery sensu Townrow (1969) = Festuca plebeia | 4b:196 |
| | <i>Festuca bromoides</i> L. = Vulpia bromoides | 4b:203 |
| | <i>Festuca distichophylla</i> (Labill.) Hook.f. nom. illeg. = Distichlis distichophylla | |
| | <i>Festuca dives</i> F.Muell. = Dryopoa dives | 4b:199 |
| | <i>Festuca duriuscula</i> L. sensu Bentham (1878) = Festuca plebeia (misapplied in Tasmania) | |
| | <i>Festuca elatior</i> L. subsp. <i>arundinacea</i> (Schreb.) Hack. = Festuca arundinacea | 4b:197 |
| | <i>Festuca hookeriana</i> F.Muell. ex Hook.f. = Hookerchloa hookeriana | 4b:223 |
| | <i>Festuca littoralis</i> Labill. = Austrofestuca littoralis | 4b:223 |
| i | Festuca nigrescens Lam., Encycl. (Lamarck) 2: 460 (1788) | 4b:197 |
| | <i>Festuca ovina</i> L. sensu Rodway (1903) = Festuca plebeia | 4b:196 |
| | <i>Festuca pectinata</i> Labill. = Australopyrum pectinatum | |
| e | Festuca plebeia R.Br., Prodr. Fl. Nov. Holland. 178 (1810) | 4b:196 |
| i | Festuca rubra L., Sp. Pl. I: 74 (1753) | 4b:197 |
| | <i>Festuca scabra</i> Labill. nom. illeg., non Vahl = Anthosachne scabra | |
| | <i>Gastridium lendigerum</i> (L.) Gaudin sensu Rodway (1903) = Gastridium ventricosum | 4b:280 |
| i t | Gastridium ventricosum (Gouan) Schinz & Thell., Vierteljahrsschr. Naturf. Ges. Zürich 58: 39 (1913) | 4b:280 |
| i | Gaudinia fragilis (L.) P.Beauv., Ess. Agrostogr. 95 (1812) | 4b:238 |
| | Glyceria australis C.E.Hubb., Bull. Misc. Inform. Kew. 1934: 450 (1934) | 4b:228 |
| i | Glyceria declinata Bréb., Fl. Normandie, ed. 3: 354 (1859) | 4b:229 |
| i t | Glyceria fluitans (L.) R.Br., Prodr. Fl. Nov. Holland. 179 (1810) | 4b:229 |
| | <i>Glyceria fluitans</i> (L.) R.Br. sensu Rodway (1903) = Glyceria australis | 4b:228 |
| i | Glyceria maxima (Hartm.) Holmb., Bot. Not. 72: 97 (1919) | 4b:229 |
| | <i>Glyceria notata</i> Chevall. sensu Buchanan (2005) = Glyceria plicata | |
| i * | <i>Glyceria plicata</i> (Fr.) Fr., Novitiae Florae Suecicae Mantissa 3: 176 (1845) | 4b:229 |
| | <i>Glyceria stricta</i> Hook.f. = Puccinellia stricta | 4b:205 |
| i | Hainardia cylindrica (Willd.) Greuter, Boissiera 13: 177 (1967) | 4b:227 |
| | <i>Hemarthria compressa</i> (L.f.) R.Br. sensu Bentham (1878) = Hemarthria uncinata (misapplied in Tasmania) | |
| | Hemarthria uncinata R.Br. var. <i>uncinata</i> , Prodr. Fl. Nov. Holland. 207 (1810) | 4b:359 |
| | <i>Hierochloe antarctica</i> (Labill.) R.Br. = Hierochloe redolens | |
| | <i>Hierochloe borealis</i> Roem. & Schult. sensu Hooker (1860) = Hierochloe fraseri (misapplied in Tasmania) | |
| e | Hierochloe fraseri Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) I: 93 (1845) | 4b:248 |
| | Hierochloe rariflora Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) I: 93 (1845) | 4b:249 |

| | | |
|-----|--|--------|
| | Hierochloe redolens (Vahl) R.Br. ex Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 2: 514 (1817) | 4b:247 |
| | Hierochloe redolens (Vahl) R.Br. ex Roem. & Schult. var. <i>fraseri</i> (Hook.f.) Benth. = Hierochloe fraseri | |
| i | Holcus lanatus L., Sp. Pl. 2: 1048 (1753) | 4b:245 |
| i * | Holcus mollis L., Syst. Nat., ed. 10, 2: 1305 (1759) | 4b:245 |
| | Hookerchloa hookeriana (F.Muell. ex Hook.f.) E.B.Alexeev, Byull. Moskovsk. Obshch. Isp. Prir., Otd. Biol. 90: 106 (1985) | 4b:223 |
| i | Hordeum distichon L., Sp. Pl. 1: 85 (1753) | 4b:297 |
| i | Hordeum glaucum Steud., Syn. Pl. Glumac. 1: 352 (1854) | 4b:298 |
| i | Hordeum hystrix Roth, Catal. Bot. 1: 23 (1797) | 4b:300 |
| i | Hordeum leporinum Link, Linnaea 9: 133 (1834) | 4b:298 |
| i | Hordeum marinum Huds., Fl. Angl. (Hudson) (ed. 2) 1: 57 (1778) | 4b:300 |
| | Hordeum marinum Huds. subsp. <i>gussoneanum</i> (Parl.) Thell. = Hordeum hystrix | 4b:300 |
| i | Hordeum murinum L., Sp. Pl. 1: 85 (1753) | 4b:298 |
| | Hordeum murinum L. subsp. <i>glaucum</i> (Steud.) Tzvelev = Hordeum glaucum | 4b:298 |
| | Hordeum murinum L. subsp. <i>leporinum</i> (Link) Arcang. = Hordeum leporinum | 4b:298 |
| | Hordeum nodosum L. sensu Bentham (1878) a name of uncertain application | |
| i | Hordeum vulgare L., Sp. Pl. 1: 84 (1753) | 4b:297 |
| | Imperata arundinacea Cirillo sensu Rodway (1903) = Imperata cylindrica var. major | 4b:353 |
| | Imperata cylindrica (L.) P.Beauv. var. major (Nees) C.E.Hubb., Grasses of Mauritius & Rodriguez: 96 (1940) | 4b:353 |
| | Isachne globosa (Thunb. ex Murray) Kuntze, Revis. Gen. Pl. 2: 778 (1891) | |
| | Koeleria cristata Pers. = Rostraria cristata | 4b:241 |
| | Koeleria cristata (L.) Bertol. sensu Rodway (1903) = Koeleria macrantha | 4b:240 |
| i x | Koeleria macrantha (Ledeb.) Schult., Mant. 2 (Schultes): 345 (1824) | 4b:240 |
| | Koeleria phleoides (Vill.) Pers. sensu Rodway (1903) = Rostraria cristata | 4b:241 |
| | Lachnagrostis aemula (R.Br.) Trin., Fund. Agrost. (Trinius) 128 (1820) | 4b:258 |
| | Lachnagrostis aequata (Nees) S.W.L.Jacobs = Lachnagrostis rudis | 4b:261 |
| | Lachnagrostis billardierei (R.Br.) Trin. subsp. <i>billardierei</i> , Fund. Agrost. (Trinius) 128 (1820) | 4b:260 |
| e | Lachnagrostis billardierei (R.Br.) Trin. subsp. <i>tenuisetata</i> (D.I.Morris) S.W.L.Jacobs, Telopea 9: 445 (2001) | 4b:261 |
| e | Lachnagrostis collicola (D.I.Morris) S.W.L.Jacobs, Telopea 9: 445 (2001) | 4b:261 |
| | Lachnagrostis filiformis (G.Forst.) Trin., Fund. Agrost. (Trinius) 128 (1820) | 4b:259 |
| e | Lachnagrostis lacunarum (D.I.Morris) S.W.L.Jacobs, Telopea 9: 446 (2001) | 4b:260 |
| e | Lachnagrostis morrisii A.J.Br., Muellera 24: 127 (2006) | 4b:261 |
| | Lachnagrostis punicea (A.J.Br. & N.G.Walsh) S.W.L.Jacobs subsp. <i>filifolia</i> (Vickery) S.W.L.Jacobs, Telopea 10: 840 (2004) | 4b:261 |
| | Lachnagrostis punicea (A.J.Br. & N.G.Walsh) S.W.L.Jacobs subsp. <i>punicea</i> , Telopea 9: 837 (2002) | 4b:259 |
| | Lachnagrostis robusta (Vickery) S.W.L.Jacobs, Telopea 9: 447 (2001) | 4b:261 |
| | Lachnagrostis rudis (Roem. & Schult.) Trin. subsp. <i>nana</i> A.J.Br., Muellera 33: 93 (2015) | |
| | Lachnagrostis rudis (Roem. & Schult.) Trin. subsp. <i>rudis</i> , Fund. Agrost. 128 (1820) | |
| | Lachnagrostis scabra (P.Beauv.) Nees ex Steud. sensu de Salas & Baker 2014 = Lachnagrostis rudis | 4b:261 |
| i | Lagurus ovatus L., Sp. Pl. 1: 81 (1753) | 4b:281 |

| | | |
|-----|---|--------|
| | Lepturus cylindricus (Willd.) Trin. = Hainardia cylindrica | 4b:227 |
| | Lepturus incurvatus Trin. = Parapholis incurva | 4b:226 |
| i t | Leymus arenarius (L.) Hochst., Flora 31: 118 (1848) | |
| i | Lolium loliaceum (Bory & Chaub. ex Fauché) Hand.-Mazz., Ann. K. K. Naturhist. Hofmus. 28: 32 (1914) | 4b:202 |
| i | Lolium multiflorum Lam., Fl. Franç. (Lamarck) 3: 621 (1779) | 4b:201 |
| i | Lolium perenne L., Sp. Pl. 1: 83 (1753) | 4b:201 |
| | <i>Lolium perenne</i> L. var. <i>italicum</i> (A.Braun) Rodway = Lolium multiflorum | |
| i | Lolium rigidum Gaudin, Agrost. Helv. 1: 334 (1811) | 4b:201 |
| i | Lolium temulentum L. f. <i>arvense</i> (With.) Junge, Jahrb. Hamburg. Wiss. Anst. Beih. 30: 314 (1913) | 4b:200 |
| | <i>Lophochloa cristata</i> (L.) Hyl. = Rostraria cristata | 4b:241 |
| | <i>Lophopyrum elongatum</i> (Host) Á.Löve = Thinopyrum elongatum | 4b:295 |
| | <i>Lophopyrum ponticum</i> (Podp.) Á.Löve sensu Buchanan (2005) = Thinopyrum elongatum | 4b:295 |
| | <i>Microlaena gunnii</i> Hook.f. = Microlaena stipoides | |
| | Microlaena stipoides (Labill.) R.Br. var. <i>stipoides</i> , Prodr. Fl. Nov. Holland. 210 (1810) | 4b:181 |
| e | Microlaena tasmanica (Hook.f.) Benth. var. <i>subalpina</i> F.Muell. ex Benth., Fl. Austral. 7: 553 (1878) | 4b:182 |
| e | Microlaena tasmanica (Hook.f.) Benth. var. <i>tasmanica</i> , Fl. Austral. 7: 552 (1878) | 4b:181 |
| i * | <i>Molineriella minuta</i> (L.) Rouy, Fl. France [Rouy & Foucaud] 14: 102 (1913) | |
| | <i>Monerma cylindrica</i> (Willd.) Coss. & Durieu = Hainardia cylindrica | 4b:227 |
| i t | Nardus stricta L., Sp. Pl. 1: 53 (1753) | 4b:184 |
| i | Nassella leucotricha (Trin. & Rupr.) R.W.Pohl, Taxon 39: 610 (1990) | |
| i | Nassella neesiana (Trin. & Rupr.) Barkworth, Taxon 39: 611 (1990) | |
| i | Nassella trichotoma (Nees) Hack. ex Arechav., Anales Mus. Nac. Montevideo 1: 366 (1896) | 4b:193 |
| | <i>Notodanthonia caespitosa</i> (Gaudich.) Zotov = Rytidosperma caespitosum | 4b:312 |
| | <i>Notodanthonia carphoides</i> (F.Muell. ex Benth.) Zotov = Rytidosperma carphoides | 4b:306 |
| | <i>Notodanthonia diemenica</i> (D.I.Morris) H.P.Linder = Rytidosperma diemenicum | 4b:311 |
| | <i>Notodanthonia geniculata</i> (J.M.Black) Zotov = Rytidosperma geniculatum | 4b:308 |
| | <i>Notodanthonia gracilis</i> (Hook.f.) Zotov = Rytidosperma gracile | 4b:309 |
| | <i>Notodanthonia laevis</i> (Vickery) Zotov = Rytidosperma laeve | 4b:312 |
| | <i>Notodanthonia penicillata</i> (Labill.) Zotov = Rytidosperma penicillatum | 4b:313 |
| | <i>Notodanthonia pilosa</i> (R.Br.) Zotov = Rytidosperma pilosum | 4b:313 |
| | <i>Notodanthonia popinensis</i> (D.I.Morris) H.P.Linder = Rytidosperma fulvum | 4b:316 |
| | <i>Notodanthonia racemosa</i> (R.Br.) Zotov = Rytidosperma racemosum | 4b:314 |
| | <i>Notodanthonia remota</i> (D.I.Morris) H.P.Linder = Rytidosperma remotum | 4b:316 |
| | <i>Notodanthonia semiannularis</i> (Labill.) Zotov = Rytidosperma semiannulare | 4b:308 |
| | <i>Notodanthonia setacea</i> (R.Br.) Veldkamp = Rytidosperma setaceum | 4b:311 |
| | <i>Notodanthonia tenuior</i> (Steud.) Conert = Rytidosperma tenuius | 4b:310 |
| | <i>Oryzopsis miliacea</i> (L.) Benth. & Hook.f. ex Asch. & Schweinf. = Piptatherum miliaceum | 4b:194 |
| | <i>Panicum capillare</i> L. previously listed as naturalised but insufficient evidence exists to support this | 4b:335 |
| | <i>Panicum capillare</i> L. var. <i>occidentale</i> Rydb. = <i>Panicum capillare</i> | 4b:335 |
| i * | <i>Panicum gilvum</i> Launert, Mit. Bot. Staatssamml. München 8: 153 (1970) | 4b:335 |
| | <i>Panicum gracile</i> R.Br. sensu Rodway (1903) = <i>Paspalidium gracile</i> | |
| i | Panicum hillmanii Chase, J. Wash. Acad. Sci. 14: 345 (1934) | 4b:334 |

| | | |
|---|---|--------|
| i | Panicum miliaceum L., Sp. Pl. 1: 58 (1753) | 4b:333 |
| | <i>Panicum sanguinale</i> L. = Digitaria sanguinalis | 4b:346 |
| | <i>Panicum schinzii</i> Hack. sensu Curtis & Morris (1994) = Panicum gilvum | 4b:335 |
| i | Parapholis incurva (L.) C.E.Hubb., Blumea Suppl. 3: 14 (1946) | 4b:226 |
| i | Parapholis strigosa (Dumort.) C.E.Hubb., Blumea suppl. 3: 14 (1946) | 4b:226 |
| | <i>Paspalidium gracile</i> (R.Br.) Hughes sensu Townrow (1969) a name of uncertain application | |
| i | Paspalum dilatatum Poir., Encycl. (Lamarck) 5: 35 (1804) | 4b:340 |
| i | Paspalum distichum L., Syst. Nat., ed. 10. 2: 855 (1759) | 4b:340 |
| i | Paspalum urvillei Steud., Syn. Pl. Glumac. 1: 24 (1853) | 4b:340 |
| | <i>Pennisetum alopecuroides</i> (L.) Spreng. = <i>Cenchrus purpurascens</i> | |
| | <i>Pennisetum clandestinum</i> Hochst. ex Chiov. = Cenchrus clandestinus | 4b:348 |
| | <i>Pennisetum macrourum</i> Trin. = Cenchrus macrourus | 4b:351 |
| | <i>Pennisetum villosum</i> R.Br. ex Fresen. = Cenchrus longisetus | 4b:349 |
| | <i>Pentapogon billardierei</i> R.Br. sensu Rodway (1903) = Pentapogon quadrifidus | 4b:275 |
| | <i>Pentapogon billardierei</i> R.Br. var. <i>parviflorus</i> Benth. = Pentapogon quadrifidus var. parviflorus | |
| e | Pentapogon quadrifidus (Labill.) Baill. var. parviflorus (Benth.) D.I.Morris, Muelleria 7: 167 (1990) | 4b:276 |
| | Pentapogon quadrifidus (Labill.) Baill. var. quadrifidus , Hist. Pl. (Baillon) 12: 280 (1893) | 4b:275 |
| i | Phalaris aquatica L., Cent. Pl. 1: 4 (1755) | 4b:252 |
| i | Phalaris arundinacea L. var. arundinacea , Sp. Pl. 1: 55 (1753) | 4b:251 |
| i | Phalaris canariensis L., Sp. Pl. 1: 54 (1753) | 4b:251 |
| i | Phalaris minor Retz., Observ. Bot. (Retzius) 3: 8 (1783) | 4b:252 |
| i | Phalaris paradoxa L., Sp. Pl., ed. 2. 2: 1665 (1763) | 4b:251 |
| | <i>Phalaris tuberosa</i> L. = Phalaris aquatica | 4b:252 |
| i | Phleum pratense L. subsp. pratense , Sp. Pl. 1: 59 (1753) | 4b:286 |
| | Phragmites australis (Cav.) Trin. ex Steud., Nomencl. Bot. [Steudel], ed. 2. 2: 324 (1841) | 4b:322 |
| | <i>Phragmites communis</i> Trin. sensu Rodway (1903) = Phragmites australis | 4b:322 |
| i | Piptatherum miliaceum (L.) Coss., Notes sur quelques Plantes Critiques, Rares ou Nouvelles: 129 (1851) | 4b:194 |
| | <i>Poa affinis</i> R.Br. sensu Hooker (1860) a name of uncertain application | |
| i | Poa annua L., Sp. Pl. 1: 68 (1753) | 4b:211 |
| | <i>Poa australis</i> R.Br. = Poa poiformis | |
| | <i>Poa billardierei</i> (Spreng.) St.-Yves sensu Rodway (1903) = Poa poiformis var. poiformis | 4b:220 |
| i | Poa bulbosa L., Sp. Pl. 1: 70 (1753) | 4b:213 |
| | <i>Poa caespitosa</i> G.Forst. ex Spreng. var. <i>alpina</i> F.Muell. ex Benth. sensu Rodway (1903) = Poa gunnii | 4b:218 |
| | <i>Poa caespitosa</i> G.Forst. ex Spreng. var. <i>australis</i> (R.Br.) Benth. sensu Rodway (1903) = Poa rodwayi | 4b:215 |
| | <i>Poa caespitosa</i> G.Forst. ex Spreng. var. <i>laevis</i> (R.Br.) Benth. = Poa poiformis | |
| | <i>Poa caespitosa</i> G.Forst. ex Spreng. var. <i>tenera</i> (F.Muell.) Benth. sensu Rodway (1903) = Poa tenera | 4b:217 |
| | Poa clelandii Vickery, Contr. New South Wales Natl. Herb.4: 193 (1970) | 4b:221 |
| | Poa clivicola Vickery, Contr. New South Wales Natl. Herb. 4: 213 (1970) | |
| i | Poa compressa L., Sp. Pl. 1: 69 (1753) | 4b:214 |
| | Poa costiniana Vickery, Contr. New South Wales Natl. Herb. 4: 214 (1970) | 4b:218 |
| | <i>Poa distichophylla</i> (Labill.) R.Br. = Distichlis distichophylla | |
| | Poa fawcettiae Vickery, Contr. New South Wales Natl. Herb. 4: 232 (1970) | 4b:217 |

| | | |
|----|---|--------|
| e | Poa gunnii Vickery, Contr. New South Wales Natl. Herb. 4: 217 (1970) | 4b:218 |
| | Poa halmaturina J.M.Black, Trans. Roy. Soc. South Australia 66: 248 (1943) | 4b:220 |
| | Poa hiemata Vickery, Contr. New South Wales Natl. Herb. 4: 230 (1970) | 4b:217 |
| | Poa hookeri Vickery, Contr. New South Wales Natl. Herb. 4: 222 (1970) | 4b:216 |
| i | Poa infirma Kunth, Bonpl. & Kunth, Nov. Gen. Sp. [H.B.K.]1: 158 (1816) | 4b:213 |
| e | Poa jugicola D.I.Morris, Muelleria 7: 167 (1990) | 4b:219 |
| | Poa labillardierei Steud. var. acris Vickery, Contr. New South Wales Natl. Herb. 4: 205 (1970) | 4b:220 |
| | Poa labillardierei Steud. var. labillardierei , Syn. Pl. Glumac. 1: 262 (1854) | 4b:219 |
| | <i>Poa laevis</i> R.Br. = Poa poiformis | |
| e | Poa mollis Vickery, Contr. New South Wales Natl. Herb. 4: 241 (1970) | 4b:215 |
| | Poa poiformis (Labill.) Druce var. poiformis , Rep. Bot. Soc. Exch. Club Brit. Isles 1916, Suppl. 2: 640 (1917) | 4b:220 |
| | Poa poiformis (Labill.) Druce var. ramifer D.I.Morris, Muelleria 7: 169 (1990) | 4b:220 |
| i | Poa pratensis L., Sp. Pl. 1: 67 (1753) | 4b:214 |
| | <i>Poa pubinervis</i> (Vickery) S.W.L.Jacobs sensu Baker & Duretto (2011) = Austrofestuca littoralis | |
| | <i>Poa rigida</i> L. = Catapodium rigidum | 4b:224 |
| | Poa rodwayi Vickery, Contr. New South Wales Natl. Herb. 4: 235 (1970) | 4b:215 |
| | <i>Poa saxicola</i> R.Br. = Saxipoa saxicola | 4b:214 |
| | Poa sieberiana Spreng. var. hirtella Vickery, Contr. New South Wales Natl. Herb. 4: 228 (1970) | |
| | Poa sieberiana Spreng. var. sieberiana , Syst. Veg. (ed. 16) [Sprengel] 4(2, Cur. Post.): 35 (1827) | 4b:216 |
| | Poa tenera F.Muell. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 124, t.164a (1858) | 4b:217 |
| i | Poa trivialis L., Sp. Pl. 1: 67 (1753) | 4b:213 |
| | <i>Polypogon littoralis</i> Sm. = Polypogon lutosus | 4b:283 |
| i | Polypogon lutosus (Poir.) Hitchc., U.S.D.A. Bull. (1915–23) 772: 138 (1920) | 4b:283 |
| i | Polypogon maritimus Willd. var. subspatheaceus (Req.) Parl., Fl. Ital. (Parlatore) 1: 200 (1850) | 4b:282 |
| i | Polypogon monspeliensis (L.) Desf., Fl. Atlant. 1: 67 (1798) | 4b:282 |
| | <i>Polypogon viridis</i> (Gouan) Breistr. = Agrostis viridis | 4b:283 |
| | <i>Psamma arenaria</i> (L.) P.Beauv. = Ammophila arenaria | 4b:274 |
| i | Puccinellia distans (Jacq.) Parl., Fl. Ital. (Parlatore) 1: 366 (1850) | 4b:205 |
| e | Puccinellia harcusiana A.R.Williams, Fl. Australia 44A: 386 (2009) | |
| | Puccinellia perlaxa (Stapf ex N.G.Walsh) N.G.Walsh & A.R.Williams, Nuytsia 16: 464 (2007) | 4b:205 |
| | Puccinellia stricta (Hook.f.) C.H.Blom, Acta Horti Gothob. 5: 89 (1930) | 4b:205 |
| | <i>Puccinellia stricta</i> (Hook.f.) C.H.Blom var. <i>perlaxa</i> Stapf ex N.G.Walsh = Puccinellia perlaxa | 4b:205 |
| i | Rostraria cristata (L.) Tzvelev, Novosti Sist. Vyssh. Rast. 7: 47 (1971) | 4b:241 |
| | Rytidosperma australe (Petrie) Clayton & Renvoize ex Connor & Edgar, New Zealand J. Bot. 25: 166 (1987) | 4b:318 |
| | Rytidosperma caespitosum (Gaudich.) Connor & Edgar, New Zealand J. Bot. 17: 325 (1979) | 4b:312 |
| | Rytidosperma carphoides (F.Muell. ex Benth.) Connor & Edgar, New Zealand J. Bot. 17: 331 (1979) | 4b:306 |
| e | Rytidosperma diemenicum (D.I.Morris) A.M.Humphreys & H.P.Linder, Ann. Missouri Bot. Gard. 97: 358 (2010) | 4b:311 |
| | Rytidosperma dimidiatum (Vickery) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979) | 4b:317 |
| e | Rytidosperma fortunae-hibernae (Renvoize) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979) | 4b:310 |
| ?i | Rytidosperma fulvum (Vickery) A.M.Humphreys & H.P.Linder, Ann. Missouri Bot. Gard. 97: 358 (2010) | |

| | | |
|----|--|--------|
| | Rytidosperma geniculatum (J.M.Black) Connor & Edgar, New Zealand J. Bot. 17: 323 (1979) | 4b:308 |
| t | Rytidosperma gracile (Hook.f.) Connor & Edgar, New Zealand J. Bot. 17: 330 (1979) | 4b:309 |
| | Rytidosperma indutum (Vickery) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979) | 4b:312 |
| | Rytidosperma laeve (Vickery) Connor & Edgar, New Zealand J. Bot. 17: 325 (1979) | 4b:312 |
| e | Rytidosperma nitens (D.I.Morris) H.P.Linder, Telopea 6: 614 (1996) | 4b:315 |
| | Rytidosperma nivicola (Vickery) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979) | 4b:315 |
| | Rytidosperma nudiflorum (P.Morris) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979) | 4b:314 |
| | Rytidosperma oreophilum H.P.Linder & N.G.Walsh, Muelleria 8: 283 (1995) | |
| | Rytidosperma pallidum (R.Br.) A.M.Humphreys & H.P.Linder recorded in error | |
| e | Rytidosperma pauciflorum (R.Br.) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979) | 4b:309 |
| | Rytidosperma penicillatum (Labill.) Connor & Edgar, New Zealand J. Bot. 17: 327 (1979) | 4b:313 |
| | Rytidosperma pilosum (R.Br.) Connor & Edgar, New Zealand J. Bot. 17: 326 (1979) | 4b:313 |
| | Rytidosperma popinensis (D.I.Morris) A.M.Humphreys & H.P.Linder = Rytidosperma fulvum | 4b:316 |
| | Rytidosperma procerum (Vickery) Connor & Edgar = Rytidosperma indutum | 4b:312 |
| | Rytidosperma racemosum (R.Br.) Connor & Edgar var. racemosum , New Zealand J. Bot. 17: 327 (1979) | 4b:314 |
| e | Rytidosperma remotum (D.I.Morris) A.M.Humphreys & H.P.Linder, Ann. Missouri Bot. Gard. 97: 359 (2010) | 4b:316 |
| | Rytidosperma semiannulare (Labill.) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979) | 4b:308 |
| | Rytidosperma setaceum (R.Br.) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979) | 4b:311 |
| | Rytidosperma tenuius (Steud.) A.Hansen & Sunding, Fl. Macaronesia, ed. 2, 1: 93 (1979) | 4b:310 |
| | Saxipoa saxicola (R.Br.) Soreng, L.J.Gillespie & S.W.L.Jacobs, Austral. Syst. Bot. 22: 407 (2010) | 4b:214 |
| | Schedonorus hookerianus (F.Muell. ex Hook.f.) P.Beauv. = Hookerochloa hookeriana | 4b:223 |
| | Schedonorus littoralis (Labill.) P.Beauv. = Austrofestuca littoralis | 4b:223 |
| | Schedonorus phoenix (Scop.) Holub = Festuca arundinacea | 4b:197 |
| i | Sclerochloa dura (L.) P.Beauv., Ess. Agrostogr. 98, 177, t.19 fig.4 (1812) | 4b:225 |
| i | Secale cereale L., Sp. Pl. 1: 84 (1753) | 4b:302 |
| | Setaria geniculata P.Beauv. var. pauciseta Desv. = Setaria parviflora | 4b:344 |
| | Setaria gracilis Kunth var. pauciseta (Desv.) B.K.Simon = Setaria parviflora | 4b:344 |
| i | Setaria italica (L.) P.Beauv., Ess. Agrostogr. 51, 170, 178 (1812) | 4b:343 |
| i | Setaria parviflora (Poir.) Kerguelen, Lejeunia 120: 161 (1987) | 4b:344 |
| | Setaria pumila (Poir.) Roem. & Schult. subsp. pumila previously listed as naturalised but insufficient evidence exists to support this | |
| i | Setaria verticillata (L.) P.Beauv., Ess. Agrostogr. 51, 171, 178 (1812) | 4b:342 |
| i | Setaria viridis (L.) P.Beauv., Ess. Agrostogr. 51, 171, 178 (1812) | 4b:343 |
| | Sieglingia decumbens (L.) Bernh. = Danthonia decumbens | 4b:317 |
| | Sorghum bicolor (L.) Moench previously listed as naturalised but insufficient evidence exists to support this | |
| i* | Sorghum halepense (L.) Pers., Syn. Pl. (Persoon) 1: 101 (1805) | |
| i | Spartina anglica C.E.Hubb., Bot. J. Linn. Soc. 76: 364 (1978) | 4b:330 |
| | Spartina townsendii H.Groves & J.Groves sensu Townrow (1969) = Spartina anglica | 4b:330 |
| | Spinifex hirsutus Labill. sensu Rodway (1903) = Spinifex sericeus | 4b:351 |
| | Spinifex sericeus R.Br., Prodr. Fl. Nov. Holland. 198 (1810) | 4b:351 |
| i | Sporobolus africanus (Poir.) Robyns & Tournay, Bull. Jard. Bot. État Bruxelles 25: 242 (1955) | 4b:328 |

| | | |
|---|--|--------|
| | Sporobolus capensis Kunth = Sporobolus africanus | 4b:328 |
| | Sporobolus indicus (L.) R.Br. = Sporobolus africanus | 4b:328 |
| | Sporobolus virginicus (L.) Kunth, Révis. Gramin. 1: 67 (1829) | 4b:328 |
| | Stenotaphrum americanum Schrank = Stenotaphrum secundatum | 4b:344 |
| i | Stenotaphrum secundatum (Walter) Kuntze, Revis. Gen. Pl. 2: 794 (1891) | 4b:344 |
| | Stipa aphanoneura Hughes = Austrostipa flavescens | 4b:190 |
| | Stipa aphylla (Rodway) J.Townrow = Austrostipa aphylla | 4b:192 |
| | Stipa bigeniculata Hughes = Austrostipa bigeniculata | 4b:190 |
| | Stipa blackii C.E.Hubb. = Austrostipa blackii | 4b:191 |
| | Stipa caudata Trin. = Amelichloa caudata | 4b:192 |
| | Stipa compacta Hughes = Austrostipa flavescens | 4b:190 |
| | Stipa elatior (Benth.) Hughes = Austrostipa flavescens | 4b:190 |
| | Stipa elegantissima Labill. sensu Labillardiere (1805) and Bentham (1878) recorded in error | |
| | Stipa eremophila Reader sensu Townrow (1969) a name of uncertain application | |
| | Stipa falcata Hughes sensu Townrow (1978) = Austrostipa scabra p.p.maj. | 4b:188 |
| | Stipa flavescens Labill. = Austrostipa flavescens | 4b:190 |
| | Stipa mollis R.Br. = Austrostipa mollis | 4b:189 |
| | Stipa neesiana Trin. & Rupr. = Nassella neesiana | |
| | Stipa nervosa Vickery var. neutralis sensu Townrow (1978) = Austrostipa rudis subsp. australis | 4b:186 |
| | Stipa nodosa S.T.Blake = Austrostipa nodosa | 4b:188 |
| | Stipa pubescens R.Br. sensu Rodway (1903) = Austrostipa pubinodis | 4b:192 |
| | Stipa pubescens R.Br. var. aphylla Rodway = Austrostipa aphylla | 4b:187 |
| | Stipa pubinodis Trin. & Rupr. = Austrostipa pubinodis | 4b:187 |
| | Stipa rudis Spreng. = Austrostipa rudis | 4b:186 |
| | Stipa scabra Lindl. = Austrostipa scabra | 4b:188 |
| | Stipa semibarbata R.Br. = Austrostipa semibarbata | 4b:189 |
| | Stipa setacea R.Br. sensu Rodway (1903) = Austrostipa stuposa | 4b:191 |
| | Stipa stipoides (Hook.f.) Veldkamp = Austrostipa stipoides | 4b:186 |
| | Stipa stuposa Hughes = Austrostipa stuposa | 4b:191 |
| | Stipa teretifolia Steud. = Austrostipa stipoides | 4b:186 |
| | Stipa variabilis Hughes sensu Townrow (1969) = Austrostipa nodosa | 4b:188 |
| | Tetrarrhena acuminata R.Br., Prodr. Fl. Nov. Holland. 210 (1810) | 4b:182 |
| | Tetrarrhena distichophylla (Labill.) R.Br., Prodr. Fl. Nov. Holland. 210 (1810) | 4b:183 |
| | Tetrarrhena juncea R.Br., Prodr. Fl. Nov. Holland. 210 (1810) | 4b:183 |
| e | Tetrarrhena oreophila D.I.Morris var. minor D.I.Morris, Records of the Queen Victoria Museum, n.ser., 55: 4 (1977) | 4b:182 |
| e | Tetrarrhena oreophila D.I.Morris var. oreophila , Records of the Queen Victoria Museum, n.ser., 55: 4 (1977) | 4b:182 |
| | Tetrarrhena tenacissima Nees sensu Hooker (1860) = Tetrarrhena juncea | |
| | Themeda australis (R.Br.) Stapf = Themeda triandra | 4b:358 |
| | Themeda triandra Forssk., Fl. Aegypt.-Arab. 178 (1775) | 4b:358 |
| i | Thinopyrum elongatum (Host) D.R.Dewey, Gene Manipulation Pl. Improv. 274 (1984) | 4b:295 |
| i | Thinopyrum junceiforme (Á.Löve & D.Löve) Á.Löve, Taxon 29: 351 (1980) | 4b:293 |

| | | |
|-------------------------|--|--------|
| | Thonandia gracilis (Hook.f.) H.P.Linder = Rytidosperma gracile | 4b:309 |
| | Thonandia semiannularis (Labill.) H.P.Linder = Rytidosperma semiannulare | 4b:308 |
| | Trisetum spicatum (L.) K.Richt. subsp. australiense Hultén ex Veldkamp, Gard. Bull. Singapore 36: 135 (1983) | 4b:239 |
| | Trisetum subspicatum (L.) P.Beauv. sensu Rodway (1903) = Trisetum spicatum subsp. australiense | 4b:239 |
| i | Triticum aestivum L., Sp. Pl. 1: 85 (1753) | 4b:303 |
| | Triticum pectinatum (Labill.) R.Br. = Australopyrum pectinatum | |
| | Triticum scabrum R.Br. = Anthosachne scabra | |
| | Triticum velutinum (Nees) Hook.f. nom. illeg., nom. superfl. = Australopyrum velutinum | |
| | Uniola distichophylla Labill. = Distichlis distichophylla | |
| i | Vulpia bromoides (L.) Gray, Nat. Arr. Brit. Pl. 2: 124 (1821) | 4b:203 |
| i | Vulpia fasciculata (Forssk.) Fritsch, Exkursionsfl. Oesterreich (Ed. 2): 74 (1909) | 4b:204 |
| | Vulpia megalura (Nutt.) Rydb. = Vulpia myuros f. megalura | 4b:204 |
| i | Vulpia muralis (Kunth) Nees, Linnaea 19: 694 (1847) | |
| i | Vulpia myuros (L.) C.C.Gmel. f. megalura (Nutt.) Stace & R.Cotton, Watsonia 11: 72 (1976) | 4b:204 |
| i | Vulpia myuros (L.) C.C.Gmel. f. myuros , Fl. Bad. 1: 8 (1805) | |
| | Zoysia macrantha Desv. subsp. walshii Night., Fl. Australia 43: 376 (2002) | 4b:332 |
| | Zoysia pungens Willd. sensu Rodway (1903) = Zoysia macrantha subsp. walshii | 4b:332 |
| | ×Agropogon littoralis (Sm.) C.E.Hubb. = Polypogon lutosus | 4b:283 |
| POSIDONIACEAE | | |
| | Caulinia antarctica R.Br. = Amphibolis antarctica (Cymodoceaceae) | |
| | Caulinia oceanica R.Br. = Potamogeton australiensis (Potamogetonaceae) | |
| | Posidonia australis Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 43 (1858) | 4b:25 |
| POTAMOGETONACEAE | | |
| x | Lepilaena australis J.Drumm. ex Harv., Hooker's J. Bot. Kew Gard. Misc. 7: 58 (1855) | 4b:22 |
| | Lepilaena bilocularis Kirk, Trans. & Proc. New Zealand Inst. 28: 500 (1896) | 4b:24 |
| | Lepilaena cylindrocarpa (Körn. ex Müll.Stuttg) Benth., Fl. Austral. 7: 180 (1878) | 4b:22 |
| | Lepilaena marina E.L.Robertson, Marine Benthic Fl. South. Australia 1: 80 (1984) | 4b:23 |
| | Lepilaena patentifolia E.L.Robertson, Fl. S. Australia 4: 1736 (1986) | 4b:23 |
| | Lepilaena preissii (Lehm.) F.Muell., Fragm. (Mueller) 8: 217 (1874) | 4b:24 |
| | Potamogeton australiensis A.Benn., J. Bot. 48: 149 (1910) | 4b:19 |
| | Potamogeton cheesemanii A.Benn., J. Bot. 66: 66 (1883) | 4b:19 |
| | Potamogeton crispus L., Sp. Pl. 1: 126 (1753) | 4b:18 |
| | Potamogeton drummondii Benth., Fl. Austral. 7: 171 (1878) | |
| | Potamogeton gramineus L. sensu Hooker (1860) = Potamogeton ochreatus identity somewhat uncertain | |
| | Potamogeton heterophyllus Schreb. sensu Hooker (1860) = Potamogeton perfoliatus identity somewhat uncertain | |
| | Potamogeton natans L. sensu Rodway (1903) misapplied to P. cheesemanii , P. australiensis and P. drummondii 4b:19 | |
| | Potamogeton obtusifolius Mert. & W.D.J.Koch sensu Rodway (1903) = Potamogeton crispus | 4b:18 |
| | Potamogeton ochreatus Raoul, Ann. Sci. Nat., Bot., sér. 3, 2: 117 (1844) | 4b:18 |
| | Potamogeton pectinatus L. = Stuckenia pectinata | 4b:18 |

| | | |
|---------------------|---|-------|
| | Potamogeton perfoliatus L., Sp. Pl. 1: 126 (1753) | 4b:18 |
| | Potamogeton praelongus Wulfen sensu Rodway (1903) = Potamogeton ochreatus | 4b:18 |
| | Potamogeton tricarinatus F.Muell. & A.Benn. ex A.Benn. sensu Baker & de Salas (2013) misapplied to P. australiensis , P. cheesemanii and P. drummondii | 4b:19 |
| | Stuckenia pectinata (L.) Börner, Fl. Deut. Volk. 713 (1912) | 4b:18 |
| | Zannichellia palustris L. sensu Hooker (1860) probably misapplied to Lepilaena spp. | |
| RESTIONACEAE | | |
| | Acion hookeri (D.I.Morris) B.G.Briggs & L.A.S.Johnson = Chordifex hookeri | 4b:46 |
| | Acion monocephalum (R.Br.) B.G.Briggs & L.A.S.Johnson = Chordifex monocephalus | 4b:47 |
| | Apodasmia brownii (Hook.f.) B.G.Briggs & L.A.S.Johnson, Telopea 7: 371 (1998) | 4b:39 |
| | Baloskion australe (R.Br.) B.G.Briggs & L.A.S.Johnson, Telopea 8: 23 (1998) | 4b:46 |
| | Baloskion tetraphyllum (Labill.) B.G.Briggs & L.A.S.Johnson subsp. tetraphyllum , Telopea 8: 23 (1998) | 4b:45 |
| | Calorophus ater L.A.S.Johnson & B.G.Briggs = Calorophus erostris | 4b:42 |
| | Calorophus elongatus Labill., Nov. Holl. Pl. 2: 78, t.228 (1806) | 4b:41 |
| e | Calorophus erostris (C.B.Clarke) L.A.S.Johnson & B.G.Briggs, The Student's Flora of Tasmania 4b: 425 (1994) | 4b:42 |
| | Calorophus lateriflorus (R.Br.) F.Muell. = Empodisma minus | 4b:43 |
| e | Chordifex hookeri (D.I.Morris) B.G.Briggs, Telopea 10: 685 (2004) | 4b:46 |
| e | Chordifex monocephalus (R.Br.) B.G.Briggs, Telopea 10: 685 (2004) | 4b:47 |
| | Empodisma minus (Hook.f.) L.A.S.Johnson & D.F.Cutler, Kew Bull. 28: 383 (1974) | 4b:43 |
| | Eurychorda complanata (R.Br.) B.G.Briggs & L.A.S.Johnson, Telopea 7: 359 (1998) | 4b:46 |
| | Hypolaena exsulca R.Br. recorded in error | |
| | Hypolaena fastigiata R.Br., Prodr. Fl. Nov. Holland. 251 (1810) | 4b:40 |
| | Hypolaena lateriflora (R.Br.) Benth. = Empodisma minus | 4b:43 |
| | Hypolaena longissima Benth. sensu Rodway (1903) = Calorophus elongatus | 4b:41 |
| | Leptocarpus brownii Hook.f. = Apodasmia brownii | 4b:39 |
| | Leptocarpus tenax (Labill.) R.Br., Prodr. Fl. Nov. Holland. 250 (1810) | 4b:39 |
| | Lepyrodia muelleri Benth., Fl. Austral. 7: 215 (1878) | 4b:43 |
| | Lepyrodia tasmanica Hook.f. = Sporadanthus tasmanicus | 4b:44 |
| | Restio australis R.Br. = Baloskion australe | 4b:46 |
| | Restio complanatus R.Br. = Eurychorda complanata | 4b:46 |
| | Restio glaber (Rodway) L.A.S.Johnson & B.G.Briggs = Chordifex monocephalus | 4b:47 |
| | Restio gracilis R.Br. sensu Rodway (1903) = Baloskion australe | 4b:46 |
| | Restio hookeri D.I.Morris = Chordifex hookeri | 4b:46 |
| | Restio lateriflorus R.Br. nom. illeg. = Calorophus elongatus | |
| | Restio monocephalus R.Br. = Chordifex monocephalus | 4b:47 |
| | Restio oligocephalus F.Muell. = Chordifex monocephalus | 4b:46 |
| | Restio oligocephalus F.Muell. var. glabrum Rodway = Chordifex monocephalus | |
| | Restio oligocephalus F.Muell. var. intermedius Rodway = Chordifex hookeri | |
| | Restio tetraphyllum Labill. = Baloskion tetraphyllum | 4b:45 |
| | Schoenodum tenax Labill. = Leptocarpus tenax | |
| | Sporadanthus tasmanicus (Hook.f.) B.G.Briggs & L.A.S.Johnson, Telopea 8: 31 (1998) | 4b:44 |

- e **Winifredia sola** L.A.S.Johnson & B.G.Briggs, *Telopea* 2: 738 (1986) 4b:47
- RUPPIACEAE**
- Ruppia antarctica* Labill. = **Amphibolis antarctica** (Cymodoceaceae)
- Ruppia maritima* L. sensu Rodway (1903) = **Ruppia polycarpa** (misapplied in Tasmania) 4b:21
- Ruppia megacarpa** R.Mason, *New Zealand J. Bot.* 5: 525 (1967) 4b:21
- Ruppia polycarpa** R.Mason, *New Zealand J. Bot.* 5: 524 (1967) 4b:21
- Ruppia tuberosa** J.S.Davis & Toml., *J. Arnold Arbor.* 55: 60 (1974) 4b:21
- TYPHACEAE**
- Typha angustifolia* L. sensu Rodway (1903) = **Typha domingensis** & **T. orientalis**
- Typha domingensis** Pers., *Syn. Pl. (Persoon)* 2: 532 (1807) 4b:364
- i **Typha latifolia** L., *Sp. Pl.* 2: 971 (1753) 4b:363
- Typha orientalis** C.Presl, *Epimel. Bot.* 239 (1851) 4b:363
- XANTHORRHOEACEAE**
- e **Xanthorrhoea arenaria** D.J.Bedford, *Fl. Australia* 46: 225 (1986) 4b:421
- Xanthorrhoea australis** R.Br., *Prodr. Fl. Nov. Holland.* 288 (1810) 4b:419
- e **Xanthorrhoea bracteata** R.Br., *Prodr. Fl. Nov. Holland.* 288 (1810) 4b:421
- Xanthorrhoea hastilis* R.Br. sensu Hooker (1860) = **Xanthorrhoea australis** (misapplied in Tasmania)
- Xanthorrhoea minor* R.Br. sensu Rodway (1903) = **Xanthorrhoea arenaria** & **X. bracteata** 4b:421
- XYRIDACEAE**
- Xyris gracilis* R.Br. sensu Bentham (1878), misapplied to **X. marginata**, **X. muelleri**, and **X. tasmanica**
- Xyris gracilis* R.Br. subsp. *tasmanica* D.I.Morris = **Xyris tasmanica** 4b:35
- Xyris gracilis* R.Br. var. *bracteata* Benth. = **Xyris muelleri**
- e **Xyris marginata** Rendle, *J. Bot.* 37: 503 (1899) 4b:35
- e **Xyris muelleri** Malme, *Svensk. Bot. Tidskr.* 21: 381 (1928) 4b:35
- Xyris operculata** Labill., *Nov. Holl. Pl.* 1: 14, t.10 (1805) 4b:34
- e **Xyris tasmanica** (D.I.Morris) Doust & B.J.Conn, *Austral. Syst. Bot.* 10(2): 212 (1997) 4b:35
- ZOSTERACEAE**
- Heterozostera nigricaulis** J.Kuo, *Aquatic Bot.* 81(2): 110 (2005)
- Heterozostera tasmanica** (G.Martens ex Asch.) Hartog, *Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Sect. 2.* 59: 116 (1970) 4b:28
- Nanozostera muelleri* (Irmisch ex Asch.) Toml. & Posl. = **Zostera muelleri** subsp. **muelleri** 4b:29
- Zostera marina* R.Br. = **Heterozostera nigricaulis**
- Zostera muelleri** Irmisch ex Asch. subsp. **muelleri**, *Linnaea* 35: 168 (1868) 4b:29
- Zostera nana* Mert. ex Roth sensu Rodway (1903) = **Zostera muelleri** subsp. **muelleri** 4b:29
- Zostera tasmanica* G.Martens ex Asch. = **Heterozostera tasmanica** 4b:28
- GYMNOSPERMS**
- CUPRESSACEAE**
- e **Athrotaxis cupressoides** D.Don, *Ann. Nat. Hist.* 1: 234 (1838) 1:6
- e **Athrotaxis** × **laxifolia** Hook., *Icon. Pl.* 6: t.573 (1843) = **A. cupressoides** × **A. selaginoides** 1:6
- e **Athrotaxis selaginoides** D.Don, *Ann. Nat. Hist.* 1: 234 (1838) 1:7
- Athrotaxis tetragona* Hook. = **Microcachrys tetragona** (Podocarpaceae)

- Callitris cupressiformis F.Muell. var. tasmanica (Benth.) Maiden = **Callitris rhomboidea**
- e **Callitris oblonga** Rich. & A.Rich. subsp. **oblonga**, Comm. Bot. Conif. Cycad. 49, t.18 fig.2 (1826) 1:5
- Callitris rhomboidea** R.Br. ex Rich. & A.Rich., Comm. Bot. Conif. Cycad. 47, t.18 fig.1 (1826) 1:5
- Callitris rhomboidea R.Br. ex Rich. & A.Rich. var. tasmanica (Benth.) Ewart, B.Rees & B.Wood =
Callitris rhomboidea
- Callitris tasmanica (Benth.) R.T.Baker & H.G.Sm. = **Callitris rhomboidea**
- e **Diselma archeri** Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 353 t.98 (1857) 1:5
- Fitzroya archeri (Hook.f.) Benth. & Hook.f. = **Diselma archeri** 1:5
- Frenela australis R.Br. ex Benth. nom. illeg., sensu Bentham (1878) = **Callitris oblonga**
- Frenela rhomboidea (R.Br. ex Rich. & A.Rich.) Endl. var. tasmanica Benth. = **Callitris rhomboidea**
- i **Hesperocyparis macrocarpa** (Hartw. ex Gordon) Bartel, Phytologia 91: 182 (2009)

PINACEAE

- i **Pinus pinaster** Aiton, Hortus Kew. (W. Aiton) 3: 367 (1789)
- i **Pinus radiata** D.Don, Trans. Linn. Soc. London 17: 442 (1837) 1:8
- i # **Pinus wallichiana** A.B.Jacks., Bull. Misc. Inform. Kew 1938(2): 85 (1938)

PODOCARPACEAE

- Dacrydium franklinii Hook.f. = **Lagarostrobos franklinii** 1:3
- e **Lagarostrobos franklinii** (Hook.f.) Quinn, Austral. J. Bot. 30: 316 (1982) 1:3
- e **Microcachrys tetragona** (Hook.) Hook.f., London J. Bot. 4: 150 (1845) 1:3
- Microstrobos niphophilus J.Garden & L.A.S.Johnson = **Pherosphaera hookeriana** 1:2
- e **Pherosphaera hookeriana** W.Archer bis, Hooker's J. Bot. Kew Gard. Misc. 2: 52 (1850) 1:2
- e **Phyllocladus aspleniifolius** (Labill.) A.Rich. ex Hook.f., London J. Bot. 4: 151 (1845) 1:2
- Phyllocladus rhomboidalis A.Rich. = **Phyllocladus aspleniifolius** 1:2
- Podocarpus alpinus Hook.f. = **Podocarpus lawrencei** 1:4
- Podocarpus aspleniifolia Labill. = **Phyllocladus aspleniifolius**
- Podocarpus lawrencei** Hook.f., London J. Bot. 4: 151 (1845) 1:4

PTERIDOPHYTES

ADIANTACEAE

- Adiantum aethiopicum** L., Syst. Nat., ed. 10, 2: 1329 (1759)
- Adiantum assimile Sw. sensu Brown (1810) = **Adiantum aethiopicum**
- Adiantum trigonum Labill. = **Adiantum aethiopicum**
- Anogramma leptophylla** (L.) Link, Fil. Spec. 137 (1841)
- Cheilanthes austrotenuifolia** H.M.Quirk & T.C.Chambers, Austral. J. Bot. 31: 510 (1983)
- Cheilanthes distans** (R.Br.) Mett., Abh. Senckenberg. Naturf. Ges. 3: 69 (1859)
- Cheilanthes sieberi** Kunze subsp. **sieberi**, Pl. Preiss. [J.G.C.Lehman] 2: 112 (1846)
- Cheilanthes tenuifolia (Burm.f.) Sw. sensu Rodway (1903) = **Cheilanthes austrotenuifolia**
- Pellaea calidirupium** Brownsey & Lovis, New Zealand J. Bot. 28: 197 (1990)
- Pellaea falcata** (R.Br.) Fée, Mem. Foug., 5. Gen. Filic. 129 (1850)
- Platyloma falcata (R.Br.) J.Sm. = **Pellaea falcata**

ASPLENIACEAE

- Asplenium appendiculatum** (Labill.) C.Presl subsp. **appendiculatum**, Tent. Pterid. 106 (1836)

- Asplenium brownii* J.Sm. sensu Hooker (1860) = **Diplazium australe**
- Asplenium bulbiferum*** G.Forst. subsp. ***gracillimum*** (Colenso) Brownsey, New Zealand J. Bot. 15: 60 (1977)
- Asplenium bulbiferum* G.Forst. var. *hookeriana* (Colenso) Rodway = ***Asplenium hookerianum***
- Asplenium flabellifolium*** Cav., Descr. Pl. (Cavanilles): 257 (1801)
- Asplenium flaccidum*** G.Forst. subsp. ***flaccidum***, Fl. Ins. Austr. 80 (1786)
- Asplenium hookerianum*** Colenso, Tasmanian J. Nat. Sci. 2: 169 (1844)
- Asplenium laxum* R.Br. nom. illeg. = ***A. appendiculatum*** × ***A. bulbiferum*** subsp. ***gracillimum***
- Asplenium lucidum* G.Forst. sensu Hooker (1860) = ***Asplenium obtusatum***
- Asplenium obliquum* G.Forst. sensu Labillardiere (1805) = ***Asplenium obtusatum*** subsp. ***northlandicum*** (misapplied in Tasmania)
- Asplenium obtusatum*** G.Forst. subsp. ***northlandicum*** Brownsey, New Zealand J. Bot. 15: 49, t.6 (1977)
- Asplenium terrestre* Brownsey = ***Asplenium appendiculatum***
- Asplenium trichomanes*** L. subsp. ***quadrivalens*** D.E.Mey., Ber. Deutsch. Bot. Ges. 74: 456 (1961)
- Asplenium trichomanes*** L. subsp. ***trichomanes***, Sp. Pl. 2: 1080 (1753)
- Asplenium umbrosum* J.Sm. sensu Bentham (1878) = ***Diplazium australe*** (Athyraceae) misapplied in Tasmania
- Caenopteris appendiculata* Labill. = ***Asplenium appendiculatum***
- Pleurosorus rutifolius*** (R.Br.) Fée, Mem. Foug., 5. Gen. Filic. 179, t.16c (1852)

ATHYRIACEAE

- Allantodia australis* R.Br. = ***Diplazium australe***
- Athyrium australe* (R.Br.) C.Presl = ***Diplazium australe***
- Cystopteris fragilis* (L.) Bernh. var. *tasmanica* (Hook.) Hook.f. = ***Cystopteris tasmanica***
- Cystopteris tasmanica*** Hook., Sp. Fil. 1: 199 (1846)
- Diplazium australe*** (R.Br.) N.A.Wakef., Vict. Naturalist 58: 142 t.3 (1942)

AZOLLACEAE

- Azolla filiculoides* Lam. = ***Azolla rubra***
- Azolla rubra*** R.Br., Prodr. Fl. Nov. Holland. 167 (1810)

BLECHNACEAE

- Blechnum aggregatum* (Colenso) Tindale sensu Willis (1973) = ***Blechnum chambersii***
- Blechnum cartilagineum*** Sw., Syn. Fil. (Swartz): 114 (1806)
- Blechnum chambersii*** Tindale, Flora of the Sydney Region, ed. 2: 86 (1972)
- Blechnum fluviatile*** (R.Br.) E.J.Lowe ex Salomon, Nomencl. Gefässkrypt. 115 (1883)
- Blechnum minus*** (R.Br.) Ettingsh., Denkschr. Kaiserl. Akad. Wiss. Math.-Naturwiss. Kl. 23: 63, t.8 (1864)
- Blechnum neohollandicum*** Christenh., Phytotaxa 19: 20 (2011)
- Blechnum nudum*** (Labill.) Mett. ex Luerss., lora oder Allgemeinen Botanischer Zeitung 59: 292 (1876)
- Blechnum parrissiae*** Christenh., Phytotaxa 19: 20 (2011)
- Blechnum patersonii*** (R.Br.) Mett. subsp. ***patersonii***, Fil. Hort. Bot. Lips.: 64, t.4 figs. 4-10 (1856)
- Blechnum pennamarina*** (Poir.) Kuhn subsp. ***alpina*** (R.Br.) T.C.Chambers & P.A.Farrant, Fern Gaz. 15: 96 (1996)
- Blechnum procerum* (G.Forst.) Sw. sensu Labillardiere (1805) = ***Blechnum wattsii***
- Blechnum rupestre* (Kaulf. ex Link) Christenh. (cited in error) = ***Blechnum spinulosum***
- Blechnum spinulosum*** Poir., Encycl. (Lamarck) Suppl. 1: 644 (1810)

- Blechnum vulcanicum** (Blume) Kuhn, Ann. Mus. Bot. Lugduno-Batavi4: 284 (1869)
Blechnum wattsii Tindale, Contr. New South Wales Natl. Herb. 3: 247 (1963)
Doodia aspera R.Br. = **Blechnum neohollandicum**
Doodia australis (Parris) Parris = **Blechnum parrissiae**
Doodia caudata (Cav.) R.Br. = **Blechnum spinulosum**
Doodia media R.Br. = **Blechnum parrissiae** (Tasmanian plants)
Lomaria alpina Spreng. = **Blechnum pennamarina** subsp. **alpina**
Lomaria capensis (L.) Willd. sensu Bentham (1878) = **Blechnum wattsii** (misapplied in Tasmania)
Lomaria discolor (G.Forst.) Willd. sensu Rodway (1903) = **Blechnum nudum**
Lomaria fluviatilis Spreng. = **Blechnum fluviatile**
Lomaria lanceolata Spreng. = **Blechnum chambersii**
Lomaria patersonii Spreng. = **Blechnum patersonii**
Lomaria procera Spreng. sensu Rodway (1903) = **Blechnum wattsii**
Lomaria procera Spreng. var. paludosa Rodway = **Blechnum minus**
Lomaria vulcanica Blume = **Blechnum vulcanicum**
Onoclea nuda Labill. = **Blechnum nudum**
Stegania alpina R.Br. = **Blechnum pennamarina** subsp. **alpina**
Stegania falcata R.Br. = **Blechnum nudum**
Stegania fluviatilis R.Br. = **Blechnum fluviatile**
Stegania lanceolata R.Br. = **Blechnum chambersii**
Stegania minor R.Br. = **Blechnum minus**
Stegania nuda R.Br. = **Blechnum nudum**
Stegania patersonii R.Br. = **Blechnum patersonii**
Stegania procera R.Br. = **Blechnum wattsii**

CULCITACEAE

- Calochlaena dubia** (R.Br.) M.D.Turner & R.A.White, Amer. Fern J. 78: 92 (1988)
Culcita dubia (R.Br.) Maxon = **Calochlaena dubia**
Davallia dubia R.Br. = **Calochlaena dubia**

CYATHEACEAE

- Alsophila australis R.Br. = **Cyathea australis**
Cyathea australis (R.Br.) Domin subsp. **australis**, Pteridophyta 262 (1929)
Cyathea cunninghamii Hook.f., Icon. Pl. 10: t.985 (1854)
Cyathea × marcescens N.A.Wakef., Vict. Naturalist 59: 33, figs.1-5 (1942) = **C. australis** × **C. cunninghamii**
Cyathea medullaris (G.Forst.) Sw. sensu Bentham (1878) = **Cyathea cunninghamii** (misapplied in Tasmania)

DENNSTAEDTIACEAE

- Histiopteris incisa** (Thunb.) J.Sm., Hist. Fil. 295 (1875)
Hypolepis amaurochis (Kunze) Hook., Sp. Fil. 2: 62 (1852)
Hypolepis australis N.A.Wakef. = **Hypolepis amaurochis**
t **Hypolepis distans** Hook., Sp. Fil. 2: 70, t.95c (1852)
Hypolepis glandulifera Brownsey & Chinnock, J. Adelaide Bot. Gard. 10: 16, fig.8 (1987)
Hypolepis muelleri N.A.Wakef., Vict. Naturalist 60: 42 (1943)

- Hypolepis punctata (Thunb.) Mett. ex Kuhn sensu Willis (1973) = **Hypolepis glandulifera**
Hypolepis rugosula (Labill.) J.Sm., Bot. Mag., 72 (Companion): 8 (1846)
Hypolepis tenuifolia (G.Forst.) Bernh. ex C.Presl sensu Rodway (1903) = **Hypolepis amaurorachis** &
H. rugosula
Pteridium esculentum (G.Forst.) Cockayne = **Pteridium esculentum** subsp. **esculentum**
Pteridium esculentum (G.Forst.) Cockayne subsp. **esculentum**, Report of the Botanical Survey of
Tongariro National Park: 34 (1908)

DICKSONIACEAE

- Dicksonia antarctica** Labill., Nov. Holl. Pl. 2: 100, t.249 (1807)
Dicksonia dubia (R.Br.) Gaudich. sensu Hooker (1860) = **Calochlaena dubia**

DRYOPTERIDACEAE

- Aspidium aculeatum (L.) Sw. sensu Bentham (1878) = **Polystichum proliferum** (misapplied in
Tasmania)
Aspidium capense Willd. nom. illeg. = **Rumohra adiantiformis**
Aspidium coriaceum (Sw.) Sw. sensu Brown (1810) = **Rumohra adiantiformis**
Aspidium decompositum (R.Br.) Spreng. sensu Bentham (1878) = **Lastreopsis acuminata** (misapplied
in Tasmania)
Aspidium proliferum R.Br. = **Polystichum proliferum**
Lastreopsis acuminata (Houlston) C.V.Morton, Contr. U.S. Natl. Herb. 30: 245 (1973)
Lastreopsis hispida (Sw.) Tindale, Vict. Naturalist 73: 183 (1957)
Lastreopsis shepherdii (Kuntze ex Mett.) Tindale = **Lastreopsis acuminata**
Nephrodium decompositum R.Br. sensu Rodway (1903) = **Lastreopsis acuminata**
Nephrodium hispidum (Sw.) Hook. = **Lastreopsis hispida**
Polystichum coriaceum Schott = **Rumohra adiantiformis**
Polystichum proliferum (R.Br.) C.Presl, Tent. Pterid. 83 (1836)
Polystichum vestitum (G.Forst.) C.Presl sensu Rodway (1903) = **Polystichum proliferum**
Rumohra adiantiformis (G.Forst.) Ching, Sinensia 5: 70 (1934)

EQUISETACEAE

- i **Equisetum hyemale** L., Sp. Pl. 2: 1062 (1753)

GLEICHENIACEAE

- Ctenopteris heterophylla (Labill.) Tindale = **Notogrammitis heterophylla**
e **Gleichenia abscida** Rodway, The Tasmanian Flora: 289 (1903)
e **Gleichenia alpina** R.Br., Prodr. Fl. Nov. Holland. 161 (1810)
Gleichenia circinata Sw. sensu Rodway (1903) = **Gleichenia microphylla**
Gleichenia dicarpa R.Br., Prodr. Fl. Nov. Holland. 161 (1810)
Gleichenia flabellata R.Br. sensu Rodway (1903) = **Sticherus tener** & **S. urceolatus**
Gleichenia microphylla R.Br., Prodr. Fl. Nov. Holland. 161 (1810)
Gleichenia tenera R.Br. = **Sticherus tener**
Grammitis sp. A sensu Garrett 1996 = **Notogrammitis garrettii**
Grammitis armstrongii Tindale = **Notogrammitis crassior**
Grammitis australis R.Br. = **Notogrammitis billardierei**
Grammitis sp. B sensu Garrett 1996 = **Notogrammitis gunnii**
Grammitis billardierei Willd. = **Notogrammitis billardierei**

- Grammitis garrettii Parris = **Notogrammitis garrettii**
 Grammitis gunnii Parris = **Notogrammitis gunnii**
 Grammitis heterophylla Labill. = **Notogrammitis heterophylla**
 Grammitis leptophylla (L.) Sw. sensu Bentham (1878) = **Anogramma leptophylla** (Adiantaceae)
 Grammitis magellanica Desv. subsp. nothofagei Parris = **Notogrammitis angustifolia** subsp. **nothofagei**
 Grammitis meridionalis Parris = **Notogrammitis angustifolia** subsp. **nothofagei** p.p.maj.
 Grammitis poeppigiana (Mett.) Pic.Serm. = **Notogrammitis crassior**
 Grammitis pseudociliata Parris = **Notogrammitis pseudociliata**
 Grammitis rutifolia R.Br. = **Pleurosorus rutifolius** (Aspleniaceae)
Sticherus lobatus N.A.Wakef., Vict. Naturalist 60: 110 (1943)
Sticherus tener (R.Br.) Ching, Sunyatsenia 5: 285 (1940)
 Sticherus tener form A sensu Garrett (1996) = **Sticherus urceolatus**
 Sticherus tener form B sensu Garrett (1996) = **Sticherus tener**
Sticherus urceolatus M.Garrett & Kantvilas, Muelleria 11: 103 (1998)

GRAMMITIDACEAE

- Notogrammitis angustifolia** (Jacq.) Parris subsp. **nothofagei** (Parris) Parris, New Zealand J. Bot. 50: 466 (2012)
Notogrammitis billardierei (Willd.) Parris, New Zealand J. Bot. 50: 466 (2012)
Notogrammitis crassior (Kirk) Parris, New Zealand J. Bot. 50: 467 (2012)
 e **Notogrammitis garrettii** (Parris) Parris, New Zealand J. Bot. 50: 468 (2012)
 t **Notogrammitis gunnii** (Parris) Parris, New Zealand J. Bot. 50: 468 (2012)
Notogrammitis heterophylla (Labill.) Parris, New Zealand J. Bot. 50: 469 (2012)
 t **Notogrammitis pseudociliata** (Parris) Parris, New Zealand J. Bot. 50: 469 (2012)

HEMIONITIDIACEAE

- Gymnogramma leptophylla (L.) Desv. = **Anogramma leptophylla** (Adiantaceae)
 Gymnogramma rutaefolia Hook. = **Pleurosorus rutifolius** (Aspleniaceae)

HYMENOPHYLLACEAE

- Apteropteris applanata A.M.Gray & R.G.Williams = **Hymenophyllum applanatum**
 Crepidomanes venosum (R.Br.) Bostock = **Polyphlebiium venosum**
 e **Hymenophyllum applanatum** (A.M.Gray & R.G.Williams) Ebihara & K.Iwats., Bull. Natl. Mus. Nat. Sci., Tokyo, B. 36: 43 (2010)
Hymenophyllum australe Willd., Sp. Pl., ed. 4 [Willdenow], 5: 527 (1810)
 Hymenophyllum crispatum Hook. & Grev. sensu Hooker (1860) = **Hymenophyllum australe**
Hymenophyllum cupressiforme Labill., Nov. Holl. Pl. 2: 102, t.250 fig.2 (1807)
Hymenophyllum flabellatum Labill., Nov. Holl. Pl. 2: 101, t.250 fig.1 (1807)
 Hymenophyllum javanicum Spreng. sensu Rodway (1903) = **Hymenophyllum australe**
 Hymenophyllum malingii (Hook.) Mett. sensu Rodway (1903) = **Hymenophyllum applanatum**
Hymenophyllum marginatum Hook. & Grev., Icon. Filic. 1: t.34 (1827)
 Hymenophyllum nitens R.Br. = **Hymenophyllum flabellatum**
Hymenophyllum peltatum (Poir.) Desv., Mém. Soc. Linn. Paris 6: 333 (1827)
Hymenophyllum rarum R.Br., Prodr. Fl. Nov. Holland. 159 (1810)
 Hymenophyllum tunbridgense (L.) Sm. sensu Rodway (1903) = **Hymenophyllum cupressiforme**

- Hymenophyllum unilaterale Willd. sensu Hooker (1860) = **Hymenophyllum peltatum**
 Hymenophyllum wilsoni Hook. sensu Rodway (1903) = **Hymenophyllum peltatum**
 Mecodium australe (Willd.) Copel. = **Hymenophyllum australe**
 Mecodium flabellatum (Labill.) Copel. = **Hymenophyllum flabellatum**
 Mecodium rarum (R.Br.) Copel. = **Hymenophyllum rarum**
Polyphlebium venosum (R.Br.) Copel., Philipp. J. Sci. 67: 55 (1938)
 Sphaerocionium applanatum (A.M.Gray & R.G.Williams) K.Iwats. = **Hymenophyllum applanatum**
 Trichomanes venosum R.Br. = **Polyphlebium venosum**

ISOETACEAE

- Isoetes sp. A sensu Garrett 1996 = **Isoetes sp. Maxwell River**
Isoetes drummondii A.Braun subsp. **drummondii**, Monatsber. Königl. Preuss. Akad. Wiss. Berlin 1868: 542 (1869)
 e **Isoetes elatior** F.Muell. ex A.Braun, Linnaea 25: 722 (1853)
 e **Isoetes gunnii** A.Braun, Monatsber. K. Akad. Wiss. Berlin: 535 (1868)
 e **Isoetes humilior** F.Muell. ex A.Braun, Linnaea 25: 722 (1853)
 Isoetes lacustris L. sensu Rodway (1903) = **Isoetes elatior, I. humilior & I. gunnii**
 e **Isoetes sp. Maxwell River (S.J.Jarman HO314082) Tas Herbarium**
Isoetes muelleri A.Braun, Monatsber. K. Akad. Wiss. Berlin: 541 (1868)

LINDSAEACEAE

- Lindsaea cuneata (G.Forst.) C.Chr. = **Lindsaea trichomanoides**
Lindsaea linearis Sw., J. Bot. (Schr.) 1800(2): 78 (1801)
Lindsaea trichomanoides Dryand., Trans. Linn. Soc. London 3: 43, t.11 (1797)

LYCOPODIACEAE

- Huperzia australiana** (Herter) Holub, Folia Geobot. Phytotax. 20: 70 (1985)
 Huperzia varia (R.Br.) Trevis. = **Phlegmariurus varius**
 t **Lycopodiella diffusa** (R.Br.) B.Ollg., Opera Bot. 92: 176 (1987)
Lycopodiella lateralis (R.Br.) B.Ollg., Opera Bot. 92: 176 (1987)
Lycopodiella serpentina (Kunze) B.Ollg., Opera Bot. 92: 176 (1987)
 Lycopodium australianum Herter = **Huperzia australiana**
 Lycopodium carolinianum L. sensu Benth. (1878) = **Lycopodiella serpentina** (misapplied in Tasmania)
 Lycopodium clavatum L. var. fastigiatum (R.Br.) Benth. nom. illeg. = **Lycopodium fastigiatum**
 Lycopodium decurrens R.Br. = **Lycopodium scariosum**
 Lycopodium densum Labill. = **Lycopodium deuterodensum**
Lycopodium deuterodensum Herter, Estud. Bot. (Montevideo) 20: 15 (1949)
 Lycopodium diffusum R.Br. = **Lycopodiella diffusa**
Lycopodium fastigiatum R.Br., Prodr. Fl. Nov. Holland. 165 (1810)
 Lycopodium laterale R.Br. = **Lycopodiella lateralis**
 Lycopodium myrtifolium G.Forst. nom. illeg., non L. = **Phlegmariurus varius**
Lycopodium scariosum G.Forst., Fl. Ins. Austr. 87 (1786)
 Lycopodium selago L. sensu Rodway (1903) = **Huperzia australiana**
 Lycopodium serpentinum Kunze = **Lycopodiella serpentina**
 Lycopodium uliginosum Labill. = **Selaginella uliginosa**

Lycopodium varium R.Br. = **Phlegmariurus varius**

Phlegmariurus varius (R.Br.) A.R.Field & Bostock, *Phytokeys* 20: 49 (2013)

Phylloglossum drummondii Kunze, *Bot. Zeitung (Berlin)* 1: 721 (1843)

MARSILEACEAE

i **Marsilea mutica** Mett., *Ann. Sci. Nat., Bot., sér. 4*, 15: 88 (1861)

Pilularia globulifera L. sensu Rodway (1903) = **Pilularia novae-hollandiae**

Pilularia novae-hollandiae A.Braun, *Monatsber. Königl. Preuss. Akad. Wiss. Berlin* 1863: 435 (1863)

OPHIOGLOSSACEAE

x **Botrychium australe** R.Br., *Prodr. Fl. Nov. Holland.* 164 (1810)

Botrychium lunaria (L.) Sw., *J. Bot. (Schrader)* 1800(2): 110 (1801)

Botrychium ternatum (Thunb.) Sw. sensu Rodway (1903) = **Botrychium australe**

Botrychium virginianum (L.) Sw. sensu Hooker (1860) = **Botrychium australe**

Ophioglossum lusitanicum L. subsp. **coriaceum** (A.Cunn.) R.T.Clausen, *Mem. Torrey Bot. Club* 19: 161 (1938)

Ophioglossum vulgatum L. sensu Bentham (1878) = **Ophioglossum lusitanicum** (misapplied in Tasmania)

OSMUNDACEAE

Todea africana Willd. ex Bernh. nom. illeg. = **Todea barbara**

Todea barbara (L.) T.Moore, *Index Fil. (T. Moore)*: 119 (1857)

POLYPODIACEAE

Microsorium diversifolium (Willd.) Copel. = **Microsorium pustulatum** subsp. **pustulatum**

Microsorium pustulatum (G.Forst.) Copel. subsp. **pustulatum**, *Gen. Fil. [Copeland]*: 196 (1947)

Phymatodes billardierei (R.Br.) C.Presl = **Microsorium pustulatum** subsp. **pustulatum**

Phymatosorus pustulatus (G.Forst.) Large, Braggins & P.S.Green = **Microsorium pustulatum** subsp. **pustulatum**

Polypodium australe (R.Br.) Mett. = **Notogrammitis billardierei** (Grammitidaceae)

Polypodium billardierei R.Br. = **Microsorium pustulatum**

Polypodium grammitidis R.Br. = **Notogrammitis heterophylla** (Grammitidaceae)

Polypodium punctatum Thunb. sensu Rodway (1903) = **Hypolepis rugosula** & **H. amaurorachis** (Dennstaedtiaceae)

Polypodium pustulatum G.Forst. = **Microsorium pustulatum**

Polypodium rugosulum Labill. = **Hypolepis rugosula** (Dennstaedtiaceae)

Polypodium scandens Labill. nom. illeg. = **Microsorium pustulatum** subsp. **pustulatum**

PSILOTACEAE

Psilotum truncatum R.Br. = **Tmesipteris obliqua** (misapplied in Tasmania)

Tmesipteris billardierei Endl. = **Tmesipteris obliqua**

Tmesipteris elongata P.A.Dang., *Botaniste* 2: 213 (1890-91)

Tmesipteris forsteri Endl. nom. illeg. sensu Hooker (1860) = **Tmesipteris obliqua**

Tmesipteris obliqua Chinnock, *Muelleria* 8: 60 (1993)

Tmesipteris parva N.A.Wakef., *Vict. Naturalist* 60: 143 (1944)

Tmesipteris tannensis (Spreng.) Bernh. sensu Rodway (1903) = **Tmesipteris obliqua** p.p.maj.

Tmesipteris tugana H.N.Barber = **Tmesipteris elongata**

PTERIDACEAE

Litobrochia comans (G.Forst.) C.Presl = **Pteris comans**

Litobrochia incisa (Thunb.) C.Presl = **Histiopteris incisa** (Dennstaedtiaceae)

Pteris aquilina L. sensu Rodway (1903) = **Pteridium esculentum** subsp. **esculentum**
(Dennstaedtiaceae)

Pteris comans G.Forst., Fl. Ins. Austr. 79 (1786)

Pteris endlicheriana J.Agarth sensu Hooker (1860) = **Pteris comans**

Pteris esculenta G.Forst. sensu Labillardiere (1805) = **Pteridium esculentum** subsp. **esculentum**
(Dennstaedtiaceae)

Pteris falcata R.Br. = **Pellaea falcata** (Adiantaceae)

Pteris incisa Thunb. = **Histiopteris incisa** (Deddstaedtiaceae)

Pteris tremula R.Br., Prodr. Fl. Nov. Holland. 154 (1810)

Pteris vespertilionis Labill. = **Histiopteris incisa** (Deddstaedtiaceae)

SCHIZAEACEAE

Schizaea asperula N.A.Wakef., Vict. Naturalist 59: 89 (1942)

Schizaea bifida Willd., Abh. Kurfurstl.-Mainz. Akad. Nützl. Wiss. Erfurt 5: 30, t.3 fig.3 (1802)

Schizaea fistulosa Labill., Nov. Holl. Pl. 2: 103, t.250 fig.3 (1807)

SELAGINELLACEAE

Selaginella gracillima (Kunze) Spring ex Salomon, Nomencl. Gefässkrypt. 353 (1883)

Selaginella preissiana Spring sensu Rodway (1903) = **Selaginella gracillima**

Selaginella uliginosa (Labill.) Spring, Mém. Acad. Roy. Sci. Belgique 24: 60 (1849)

THELYPTERIDACEAE

Cyclosorus penniger (G.Forst.) Ching = **Pneumatopteris pennigera**

Pneumatopteris pennigera (G.Forst.) Holttum, Blumea 21: 305 (1973)

Macquarie Island Vascular Plant Census

EUDICOTS

APIACEAE (UMBELLIFERAE)

- e **Azorella macquariensis** Orchard, *Muelleria* 7: 16 (1989)
Azorella selago Hook.f. (in Macquarie Island) = **Azorella macquariensis**

ARALIACEAE

- t **Hydrocotyle novae-zeelandiae** DC., *Prodr.* [A. P. de Candolle] 4: 67 (1830)
t **Stilbocarpa polaris** (Homb. & Jacquinot ex Hook.f.) A.Gray, *U.S. Expl. Exped.*, *Phan.* 15: 714 (1854)

ASTERACEAE (COMPOSITAE)

- Cotula plumosa* (Hook.f.) Hook.f. = **Leptinella plumosa**
t **Leptinella plumosa** Hook.f., *Bot. Antarct. Voy. I.* (Fl. Antarct.) 1: 26 (1844)
t **Pleurophyllum hookeri** Buchanan, *Trans. & Proc. New Zealand Inst.* 16: 395 (1884)

BRASSICACEAE (CRUCIFERAE)

- Cardamine corymbosa** Hook.f., *Bot. Antarct. Voy. I.* (Fl. Antarct.) 1: 6 (1844)
Cardamine depressa Hook.f. var. *stellata* (Hook.f.) Hook.f. = **Cardamine corymbosa**
Cardamine glacialis DC. var. *subcarnosa* (Hook.f.) O.E.Schulz sensu Cheeseman (1919) = **Cardamine corymbosa**

CARYOPHYLLACEAE

- i **Cerastium fontanum** Baumg. subsp. **fontanum**, *Enum. Stirp. Transsilv.* 1: 425 (1816)
n i **Cerastium glomeratum** Thuill., *Fl. Env. Paris*, ed. 2: 226 (1799)
Cerastium triviale Link sensu Cheeseman (1919) = **Cerastium fontanum** subsp. **fontanum**
(misapplied in Tasmania)
Colobanthus affinis (Hook.) Hook.f., *Bot. Antarct. Voy. III.* (Fl. Tasman.) 1: 45 (1855) 1:74
t **Colobanthus apetalus** (Labill.) Druce var. **alpinus** (Kirk) L.B.Moore, *Fl. New Zealand* 1: 214 (1961)
Colobanthus billardierei Fenzl sensu Cheeseman (1919) = **Colobanthus apetalus** var. **alpinus**
(misapplied in Tasmania)
t **Colobanthus muscoides** Hook.f., *Bot. Antarct. Voy. I.* (Fl. Antarct.) 1: 14 (1844)
Stellaria decipiens Hook.f. sensu Cheeseman (1919) = **Stellaria parviflora** (misapplied in Tasmania)
i **Stellaria media** (L.) Vill., *Hist. Pl. Dauphiné* (Villars) 3: 615 (1789) 1:71
t **Stellaria parviflora** Banks & Sol. ex Hook.f., *Bot. Antarct. Voy. II.* (Fl. Nov.-Zel.) 1: 25 (1852)

CRASSULACEAE

- t **Crassula moschata** G.Forst., *Commentat. Soc. Regiae Sci. Gott.* 9: 26 (1789)
Tillaea moschata (G.Forst.) DC. = **Crassula moschata**

HALORAGACEAE

- t **Myriophyllum triphyllum** Orchard, *Brunonia* 2: 259 (1980)

ONAGRACEAE

- t **Epilobium brunnescens** (Cockayne) P.H.Raven & Engelhorn subsp. **brunnescens**, *New Zealand J. Bot.* 9: 350 (1971)
Epilobium linnaeoides Hook.f. = **Epilobium pedunculare**
Epilobium nummulariifolium A.Cunn. var. *nerteroides* Hook.f. sensu Cheeseman (1919) = **Epilobium brunnescens** subsp. **brunnescens** (misapplied in MI)

t **Epilobium pedunculare** A.Cunn., Ann. Nat. Hist. 3: 31 (1839)

PLANTAGINACEAE

Callitriche antarctica Engelm. ex Hegelm., Verh. Bot. Vereins Prov. Brandenburg 9: 20 (1867)

POLYGONACEAE

FTO 95

i # x **Rumex crispus** L., Sp. Pl. 1: 335 (1753)

PORTULACACEAE

FTO 100

Montia fontana L. subsp. **fontana**, Sp. Pl. 1: 87 (1753)

RANUNCULACEAE

FTO 47

Ranunculus biternatus Sm. sensu Cheeseman (1919) = **Ranunculus crassipes**

Ranunculus crassipes Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 2: 224 (1845)

ROSACEAE

Acaena adscendens Vahl Sensu Hooker (1844) = **Acaena magellanica**

Acaena magellanica (Lam.) Vahl, Enum. Pl. [Vahl] 1: 297 (1804)

t **Acaena minor** (Hook.f.) Allan, Fl. New Zealand 1: 363 (1961)

Acaena sanguisorbae Vahl var. minor Hook.f. = **Acaena minor**

RUBIACEAE

t **Coprosma perpusilla** Colenso subsp. **subantarctica** Orchard, Brunonia 9: 133 (1987)

t **Galium antarcticum** Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 303 bis (1846)

MONOCOTS

CYPERACEAE

t **Carex trifida** Cav., Icon. [Cavanilles] 5: 41, t.465 (1799)

Isolepis aucklandica Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 88 (1844), pl.50 (1845)

4b:102

Scirpus aucklandicus (Hook.f.) Boeckeler = **Isolepis aucklandica**

4b:102

t **Uncinia divaricata** Boott, Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 286 (1853)

t **Uncinia hookeri** Boott, Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 91 (1844); pl.51 (1845)

Uncinia riparia R.Br. var. hookeri (Boott) Kük. = **Uncinia hookeri**

JUNCACEAE

t **Juncus scheuchzerioides** Gaudich., Ann. Sci. Nat. (Paris) 5: 100 (1825)

Luzula campestris DC. var. crinita (Hook.f.) Buchenau = **Luzula crinita**

t **Luzula crinita** Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 84, pl.48 (1844)

ORCHIDACEAE

e **Corybas dienemus** D.L.Jones, Fl. Australia 50: 572 (1993)

4a:121

Corybas macranthus (Hook.f.) Rchb.f. = **Corybas dienemus**

e **Corybas sulcatus** (M.A.Clem. & D.L.Jones) G.N.Backh., Vict. Naturalist 127: 56 (2010)

Nematoceras dienemum (D.L.Jones) D.L.Jones, M.A.Clem. & Molloy = **Corybas dienemus**

Nematoceras sulcatum M.A.Clem. & D.L.Jones = **Corybas sulcatus**

POACEAE (GRAMINEAE)

i # **Agrostis capillaris** L., Sp. Pl. 1: 62 (1753)

t **Agrostis magellanica** Lam., Tabl. Encycl. 1: 160 (1791)

i # **Agrostis stolonifera** L., Sp. Pl. 1: 62 (1753)

- i # x **Anthoxanthum odoratum** L., L., Sp. Pl. 1: 28 (1753)
Deschampsia cespitosa (L.) P.Beauv., Ess. Agrostogr. 91: 160, t.18, fig.3 (1812) 4b:242
- t **Deschampsia chapmanii** Petrie, Trans. & Proc. New Zealand Inst. 23: 401 (1891)
 Deschampsia penicillata Kirk = **Deschampsia cespitosa**
- t **Festuca contracta** Kirk, Trans. & Proc. New Zealand Inst. 27: 353 (1895)
 Festuca erecta d'Urv. nom. illeg. = **Festuca contracta**
- i **Poa annua** L., Sp. Pl. 1: 68 (1753) 4b:211
Poa cookii (Hook.f.) Hook.f., Philos. Trans. 168: 22 (1879)
- t **Poa foliosa** (Hook.f.) Hook.f., Handb. N. Zeal. Fl. 338 (1864)
 Poa hamiltonii Kirk = **Poa cookii**
- t **Poa litorosa** Cheeseman, Man. New Zealand Fl. 1156 (1906)
- e **Puccinellia macquariensis** (Cheeseman) Allan & Jansen, Trans. & Proc. Roy. Soc. New Zealand 69: 268 (1939)
 Triodia macquariensis Cheeseman = **Puccinellia macquariensis**

PTERIDOPHYTES

BLECHNACEAE

Blechnum pennamarina (Poir.) Kuhn subsp. **alpina** (R.Br.) T.C.Chambers & P.A.Farrant, Fern Gaz. 15: 96 (1996)

Lomaria penna-marina Trevis. = **Blechnum pennamarina** subsp. **alpina**

DRYOPTERIDACEAE

Aspidium vestitum (G.Forst.) Sw. = **Polystichum vestitum**

- t **Polystichum vestitum** (G.Forst.) C.Presl, Tent. Pterid. 83 (1836)

GLEICHENIACEAE

Grammitis poeppigiana (Mett.) Pic.Serm. = **Notogrammitis crassior**

GRAMMITIDACEAE

Notogrammitis crassior (Kirk) Parris, New Zealand J. Bot. 50: 467 (2012)

Polypodium australe (R.Br.) Mett. nom. illeg. sensu Cheeseman (1919) = **Notogrammitis billardierei**

HYMENOPHYLLACEAE

- t **Hymenophyllum falklandicum** Baker, Syn. Fil. (Hooker & Baker), ed. 2: 68 (1874)

LYCOPODIACEAE

Huperzia australiana (Herter) Holub, Folia Geobot. Phytotax. 20: 70 (1985)

Lycopodium varium R.Br. sensu Cheeseman (1919) = **Huperzia australiana**

APPENDIX I: CHANGES IN THE CENSUS BETWEEN THE 2018 AND 2019 EDITIONS

IA: TAXA NEW TO THE TASMANIAN CENSUS

MACQUARIE ISLAND EUDICOTS

CARYOPHYLLACEAE

i **Cerastium glomeratum** Thuill.

APPENDIX I: CHANGES IN THE CENSUS BETWEEN THE 2018 AND 2019 EDITIONS

IB: TAXA THAT HAVE CHANGED NAMES BETWEEN 2018 AND 2019

EUDICOTS

ASTERACEAE

Carpobrotus edulis = **Carpobrotus edulis** subsp.
edulis

CONVOLVULACEAE

Calystegia sepium = **Calystegia sepium** subsp.
sepium

Calystegia silvatica = **Calystegia silvatica** subsp.
silvatica

ERICACEAE

Astroloma pinifolium = **Stenantha pinifolium**
(*Aust. Syst. Bot.* 28: 384)

VALERIANACEAE

Centranthus ruber = **Centranthus ruber** subsp.
ruber

APPENDIX I: CHANGES IN THE CENSUS BETWEEN THE 2018 AND 2019 EDITIONS

IC: TAXA THAT HAVE CHANGED STATUS BETWEEN 2018 AND 2019

EUDICOTS

CRASSULACEAE

i **Aeonium haworthii**

ERICACEAE

i # **Arbutus unedo**

MONOCOTYLEDONEAE

CYPERACEAE

e **Isolepis alpina** (name misapplied to
mainland populations of *I. montivaga*)

POACEAE

i # **Eragrostis tenuifolia**

APPENDIX I: CHANGES IN THE CENSUS BETWEEN THE 2018 AND 2019 EDITIONS

ID: TAXA NOW NOT CONSIDERED TO BE PART OF THE TASMANIAN FLORA

EUDICOTS

VIOLACEAE

Viola sieberiana (misapplied to various other
Viola species in Tasmania)

APPENDIX 2. FAMILY PLACEMENT OF GENERA IN THIS CENSUS

(see Introduction for details regarding this appendix. Names not in current use in Tasmania are indented and in italics)

| | | | | | |
|--------------------|------------------|---|--------------------|------------------|---|
| Abrotanella | Asteraceae | E | Alopecurus | Poaceae | M |
| Acacia | Fabaceae | E | <i>Alsophila</i> | Cyatheaceae | P |
| Acaena | Rosaceae | E | Alstroemeria | Alstroemeriaceae | M |
| Acanthus | Acanthaceae | E | Alternanthera | Amaranthaceae | E |
| Acer | Sapindaceae | E | <i>Alyssum</i> | Brassicaceae | E |
| Acetosa | Polygonaceae | E | Alyxia | Apocynaceae | E |
| Acetosella | Polygonaceae | E | Amaranthus | Amaranthaceae | E |
| Achillea | Asteraceae | E | Amelichloa | Poaceae | M |
| <i>Achnatherum</i> | Poaceae | M | Ammi | Apiaceae | E |
| Acianthus | Orchidaceae | M | Ammobium | Asteraceae | E |
| <i>Acion</i> | Restionaceae | M | Ammophila | Poaceae | M |
| <i>Aciphylla</i> | Apiaceae | E | Amperea | Euphorbiaceae | E |
| Acradenia | Rutaceae | E | Amphibolis | Cymodoceaceae | M |
| Acrothamnus | Ericaceae | E | Amphibromus | Poaceae | M |
| Acrotriche | Ericaceae | E | Amsinckia | Boraginaceae | E |
| Actinotus | Apiaceae | E | <i>Anagallis</i> | Myrsinaceae | E |
| Actites | Asteraceae | E | Anchusa | Boraginaceae | E |
| Adiantum | Adiantaceae | P | Androstoma | Ericaceae | E |
| Adonis | Ranunculaceae | E | Anemone | Ranunculaceae | E |
| <i>Adriana</i> | Euphorbiaceae | E | Angianthus | Asteraceae | E |
| <i>Aegopodium</i> | Apiaceae | E | <i>Anguillaria</i> | Colchicaceae | M |
| Aeonium | Crassulaceae | E | Anigozanthos | Haemodoraceae | M |
| Agapanthus | Agapanthaceae | M | Anisodonteia | Malvaceae | E |
| Agastachys | Proteaceae | E | Anisotome | Apiaceae | E |
| Agave | Asparagaceae | M | Anodopetalum | Cunoniaceae | E |
| <i>Agropyron</i> | Poaceae | M | Anogramma | Adiantaceae | P |
| Agrostemma | Caryophyllaceae | E | Anopterus | Escalloniaceae | E |
| Agrostis | Poaceae | M | Anredera | Basellaceae | E |
| Aira | Poaceae | M | Anthemis | Asteraceae | E |
| Ajuga | Lamiaceae | E | <i>Anthericum</i> | Asphodelaceae | M |
| <i>Albizia</i> | Fabaceae | E | <i>Anthistria</i> | Poaceae | M |
| <i>Alchemilla</i> | Rosaceae | E | <i>Anthocercis</i> | Solanaceae | E |
| <i>Alepyrum</i> | Centrolepidaceae | M | Anthosachne | Poaceae | M |
| <i>Aletris</i> | Melanthiaceae | M | Anthoxanthum | Poaceae | M |
| Alisma | Alismataceae | M | Anthriscus | Apiaceae | E |
| <i>Allantodia</i> | Athyriaceae | P | <i>Anthyllis</i> | Fabaceae | E |
| Allittia | Asteraceae | E | Antirrhinum | Plantaginaceae | E |
| Allium | Alliaceae | M | <i>Anzybas</i> | Orchidaceae | M |
| Allocasuarina | Casuarinaceae | E | Aotus | Fabaceae | E |
| Almaleea | Fabaceae | E | Apalochlamys | Asteraceae | E |
| Alnus | Betulaceae | E | Aphanes | Rosaceae | E |
| Aloe | Asphodelaceae | M | Aphelia | Centrolepidaceae | M |

| | | | | | |
|--------------------------|--------------------|----|----------------------|-----------------|---|
| Apium | Apiaceae | E | Austrostipa | Poaceae | M |
| Apodasmia | Restionaceae | M | Avellinia | Poaceae | M |
| Aponogeton | Aponogetonaceae | M | Avena | Poaceae | M |
| <i>Aptenia</i> | Aizoaceae | E | Azolla | Azollaceae | P |
| <i>Apteropteris</i> | Hymenophyllaceae | P | Azorella | Apiaceae | E |
| Aquilegia | Ranunculaceae | E | Baeckea | Myrtaceae | E |
| Arabidopsis | Brassicaceae | E | Ballantinia | Brassicaceae | E |
| <i>Arachnorchis</i> | Orchidaceae | M | Baloskion | Restionaceae | M |
| Arbutus | Ericaceae | E | Banksia | Proteaceae | E |
| Archeria | Ericaceae | E | Barbarea | Brassicaceae | E |
| Arctium | Asteraceae | E | Bartsia | Orobanchaceae | E |
| Arctotheca | Asteraceae | E | Bassia | Amaranthaceae | E |
| Arctotis | Asteraceae | E | <i>Batrachium</i> | Ranunculaceae | E |
| Arenaria | Caryophyllaceae | E | Bauera | Cunoniaceae | E |
| Argemone | Papaveraceae | E | Baumea | Cyperaceae | M |
| Argentina | Rosaceae | E | Bedfordia | Asteraceae | E |
| Argentipallium | Asteraceae | E | <i>Bellardia</i> | Orobanchaceae | E |
| Argyrotegium | Asteraceae | E | Bellendena | Proteaceae | E |
| Aristida | Poaceae | M | Bellis | Asteraceae | E |
| Aristotelia | Elaeocarpaceae | E | Berberis | Berberidaceae | E |
| Arrhenatherum | Poaceae | M | Berkheya | Asteraceae | E |
| <i>Arthrochilus</i> | Orchidaceae | M | Bertya | Euphorbiaceae | E |
| <i>Arthrocnemum</i> | Amaranthaceae | E | Beta | Amaranthaceae | E |
| Arthropodium | Laxmanniaceae | M | Betula | Betulaceae | E |
| <i>Arundo</i> | Poaceae | M | Beyeria | Euphorbiaceae | E |
| <i>Ascyrum</i> | Hypericaceae | E | Billardiera | Pittosporaceae | E |
| Asparagus | Asparagaceae | M | Blandfordia | Blandfordiaceae | M |
| Asperula | Rubiaceae | E | Blechnum | Blechnaceae | P |
| Asphodelus | Asphodelaceae | M | <i>Boisduvalia</i> | Onagraceae | E |
| <i>Aspidium</i> | Dryopteridaceae | P | Bolboschoenus | Cyperaceae | M |
| Asplenium | Aspleniaceae | P | Borago | Boraginaceae | E |
| Astelia | Asteliaceae | M | Boronia | Rutaceae | E |
| <i>Aster</i> | Asteraceae | E | Bossiaea | Fabaceae | E |
| Asterotrichion | Malvaceae | E | Bothriochloa | Poaceae | M |
| Astroloma | Ericaceae | E | Botrychium | Ophioglossaceae | P |
| Atherosperma | Atherospermataceae | MG | <i>Brachyglottis</i> | Asteraceae | E |
| Athrotaxis | Cupressaceae | G | Brachyloma | Ericaceae | E |
| <i>Athyrium</i> | Athyriaceae | P | Brachypodium | Poaceae | M |
| Atriplex | Amaranthaceae | E | Brachyscome | Asteraceae | E |
| Australina | Urticaceae | E | <i>Bracteantha</i> | Asteraceae | E |
| Australopyrum | Poaceae | M | Brassica | Brassicaceae | E |
| <i>Austrocynoglossum</i> | Boraginaceae | E | Briza | Poaceae | M |
| <i>Austrodanthonia</i> | Poaceae | M | Bromus | Poaceae | M |
| Austroderia | Poaceae | M | <i>Brossaea</i> | Ericaceae | E |
| Austrofestuca | Poaceae | M | Brunonia | Goodeniaceae | E |

| | | | | | |
|----------------------|-------------------|---|----------------------|-------------------|----|
| Buddleja | Scrophulariaceae | E | Carpobrotus | Aizoaceae | E |
| Buglossoides | Boraginaceae | E | <i>Carpodontos</i> | Cunoniaceae | E |
| Bulbine | Asphodelaceae | M | <i>Carrichtera</i> | Brassicaceae | E |
| <i>Bunochilus</i> | Orchidaceae | M | Carthamus | Asteraceae | E |
| Burchardia | Colchicaceae | M | Cassinia | Asteraceae | E |
| Burnettia | Orchidaceae | M | Cassytha | Lauraceae | MG |
| Bursaria | Pittosporaceae | E | <i>Casuarina</i> | Casuarinaceae | E |
| Byblis | Droseraceae | E | Catapodium | Poaceae | M |
| <i>Cacalia</i> | Asteraceae | E | <i>Caucalis</i> | Apiaceae | E |
| <i>Caenopteris</i> | Aspleniaceae | P | <i>Caulinia</i> | Posidoniaceae | M |
| Caesia | Hemerocallidaceae | M | Caustis | Cyperaceae | M |
| Cakile | Brassicaceae | E | Cedronella | Lamiaceae | E |
| Caladenia | Orchidaceae | M | Celmisia | Asteraceae | E |
| Calamagrostis | Poaceae | M | <i>Celsia</i> | Scrophulariaceae | E |
| Calandrinia | Portulacaceae | E | Cenarrhenes | Proteaceae | E |
| <i>Caldasia</i> | Apiaceae | E | Cenchrus | Poaceae | M |
| <i>Calea</i> | Asteraceae | E | Centaurea | Asteraceae | E |
| Caleana | Orchidaceae | M | Centaurium | Gentianaceae | E |
| Calendula | Asteraceae | E | Centella | Apiaceae | E |
| Callistachys | Fabaceae | E | Centipeda | Asteraceae | E |
| <i>Callistemon</i> | Myrtaceae | E | Centranthus | Valerianaceae | E |
| Callitriche | Plantaginaceae | E | Centrolepis | Centrolepidaceae | M |
| Callitris | Cupressaceae | G | Centropappus | Asteraceae | E |
| Calluna | Ericaceae | E | Cerastium | Caryophyllaceae | E |
| Calocephalus | Asteraceae | E | <i>Ceratochloa</i> | Poaceae | M |
| Calochilus | Orchidaceae | M | <i>Chaetospora</i> | Cyperaceae | M |
| Calochlaena | Culcitaceae | P | Chamaecytisus | Fabaceae | E |
| <i>Calonema</i> | Orchidaceae | M | Chamaemelum | Asteraceae | E |
| <i>Calonemorchis</i> | Orchidaceae | M | Chamaescilla | Laxmanniaceae | M |
| Calorophus | Restionaceae | M | <i>Chamaesyce</i> | Euphorbiaceae | E |
| <i>Caltha</i> | Ranunculaceae | E | Chasmanthe | Iridaceae | M |
| <i>Calycotrix</i> | Myrtaceae | E | <i>Cheesemania</i> | Brassicaceae | E |
| Calystegia | Convolvulaceae | E | Cheilanthes | Adiantaceae | P |
| Calytrix | Myrtaceae | E | Chenopodium | Amaranthaceae | E |
| Camelina | Brassicaceae | E | Chiloglottis | Orchidaceae | M |
| Campanula | Campanulaceae | E | Chionogentias | Gentianaceae | E |
| Campynema | Campynemataceae | M | <i>Chionohebe</i> | Plantaginaceae | E |
| <i>Candollea</i> | Stylidiaceae | E | Chloris | Poaceae | M |
| <i>Canthium</i> | Rubiaceae | E | <i>Chlorophytum</i> | Hemerocallidaceae | M |
| Capsella | Brassicaceae | E | <i>Chondrilla</i> | Asteraceae | E |
| Cardamine | Brassicaceae | E | Chordifex | Restionaceae | M |
| <i>Cardaria</i> | Brassicaceae | E | Chorizandra | Cyperaceae | M |
| Carduus | Asteraceae | E | Chrysanthemoides | Asteraceae | E |
| Carex | Cyperaceae | M | <i>Chrysanthemum</i> | Asteraceae | E |
| Carpha | Cyperaceae | M | Chrysocephalum | Asteraceae | E |

| | | | | | |
|----------------------|------------------|---|--------------------|-------------------|---|
| <i>Chrysocoma</i> | Asteraceae | E | Cucumis | Cucurbitaceae | E |
| Cicendia | Gentianaceae | E | <i>Culcita</i> | Culcitaceae | P |
| Cichorium | Asteraceae | E | Cullen | Fabaceae | E |
| Cirsium | Asteraceae | E | <i>Cuphonotus</i> | Brassicaceae | E |
| Cistus | Cistaceae | E | Cuscuta | Convolvulaceae | E |
| Cladium | Cyperaceae | M | <i>Cyanicula</i> | Orchidaceae | M |
| <i>Claytonia</i> | Portulacaceae | E | Cyathea | Cyatheaceae | P |
| Clematis | Ranunculaceae | E | Cyathodes | Ericaceae | E |
| <i>Cnicus</i> | Asteraceae | E | <i>Cyclosorus</i> | Thelypteridaceae | P |
| Coleonema | Rutaceae | E | Cyclosporum | Apiaceae | E |
| <i>Collomia</i> | Polemoniaceae | E | Cynogeton | Juncaginaceae | M |
| Colobanthus | Caryophyllaceae | E | Cymbalaria | Plantaginaceae | E |
| Comesperma | Polygalaceae | E | Cymbonotus | Asteraceae | E |
| Conium | Apiaceae | E | <i>Cymodocea</i> | Cymodoceaceae | M |
| Conospermum | Proteaceae | E | Cynara | Asteraceae | E |
| Convolvulus | Convolvulaceae | E | Cynodon | Poaceae | M |
| Conyza | Asteraceae | E | Cynoglossum | Boraginaceae | E |
| Cooperhooia | Goodeniaceae | E | Cynosurus | Poaceae | M |
| Coprosma | Rubiaceae | E | Cyperus | Cyperaceae | M |
| Cordyline | Laxmanniaceae | M | Cyphanthera | Solanaceae | E |
| Coronidium | Asteraceae | E | Cyrtostylis | Orchidaceae | M |
| <i>Coronilla</i> | Fabaceae | E | <i>Cystanthe</i> | Ericaceae | E |
| <i>Coronopus</i> | Brassicaceae | E | Cystopteris | Athyriaceae | P |
| Correa | Rutaceae | E | Cytisus | Fabaceae | E |
| Cortaderia | Poaceae | M | <i>Dacrydium</i> | Podocarpaceae | G |
| Corunastylis | Orchidaceae | M | Dactylis | Poaceae | M |
| Corybas | Orchidaceae | M | Damasonium | Alismataceae | M |
| <i>Corysanthes</i> | Orchidaceae | M | Dampiera | Goodeniaceae | E |
| <i>Cota</i> | Asteraceae | E | Danthonia | Poaceae | M |
| Cotoneaster | Rosaceae | E | Daphne | Thymelaeaceae | E |
| Cotula | Asteraceae | E | Datura | Solanaceae | E |
| Cotyledon | Crassulaceae | E | Daucus | Apiaceae | E |
| <i>Crangonorchis</i> | Orchidaceae | M | <i>Davallia</i> | Culcitaceae | P |
| <i>Crantzia</i> | Apiaceae | E | Daviesia | Fabaceae | E |
| Craspedia | Asteraceae | E | <i>Decaspora</i> | Ericaceae | E |
| Crassula | Crassulaceae | E | Delairea | Asteraceae | E |
| Crataegus | Rosaceae | E | <i>Dendrobium</i> | Orchidaceae | M |
| <i>Crepidomanes</i> | Hymenophyllaceae | P | <i>Derwentia</i> | Plantaginaceae | E |
| Crepis | Asteraceae | E | Deschampsia | Poaceae | M |
| <i>Critesion</i> | Poaceae | M | Desmodium | Fabaceae | E |
| Crococsmia | Iridaceae | M | <i>Devauxia</i> | Centrolepidaceae | M |
| Cryptandra | Rhamnaceae | E | Deyeuxia | Poaceae | M |
| <i>Cryptostemma</i> | Asteraceae | E | Dianella | Hemerocallidaceae | M |
| Cryptostylis | Orchidaceae | M | Dianthus | Caryophyllaceae | E |
| <i>Ctenopteris</i> | Gleicheniaceae | P | <i>Dichanthium</i> | Poaceae | M |

| | | | | | |
|--------------------|-----------------|----|----------------------|------------------|---|
| Dichelachne | Poaceae | M | Echinochloa | Poaceae | M |
| Dichondra | Convolvulaceae | E | Echinopogon | Poaceae | M |
| <i>Dichopogon</i> | Laxmanniaceae | M | Echium | Boraginaceae | E |
| Dichosciadium | Apiaceae | E | Egeria | Hydrocharitaceae | M |
| Dicksonia | Dicksoniaceae | P | Ehrharta | Poaceae | M |
| <i>Didiscus</i> | Araliaceae | E | Einadia | Amaranthaceae | E |
| <i>Didymotheca</i> | Gyrostemonaceae | E | Elaeocarpus | Elaeocarpaceae | E |
| Digitalis | Plantaginaceae | E | Elatine | Elatinaceae | E |
| Digitaria | Poaceae | M | Eleocharis | Cyperaceae | M |
| <i>Dillenia</i> | Dilleniaceae | E | Eleusine | Poaceae | M |
| Dillwynia | Fabaceae | E | Elodea | Hydrocharitaceae | M |
| Dimorphotheca | Asteraceae | E | <i>Elymus</i> | Poaceae | M |
| Diplarrena | Iridaceae | M | <i>Elynanthus</i> | Cyperaceae | M |
| Diplaspis | Apiaceae | E | Elytrigia | Poaceae | M |
| <i>Diplax</i> | Poaceae | M | <i>Embothrium</i> | Proteaceae | E |
| Diplazium | Athyriaceae | P | Emex | Polygonaceae | E |
| <i>Diplodium</i> | Orchidaceae | M | Empodisma | Restionaceae | M |
| Diplotaxis | Brassicaceae | E | Epacris | Ericaceae | E |
| Dipodium | Orchidaceae | M | Epilobium | Onagraceae | E |
| Dipogon | Fabaceae | E | <i>Epipactis</i> | Orchidaceae | M |
| Dipsacus | Dipsacaceae | E | Equisetum | Equisetaceae | P |
| Disa | Orchidaceae | M | Eragrostis | Poaceae | M |
| <i>Disarrenum</i> | Poaceae | M | <i>Erechtites</i> | Asteraceae | E |
| Discaria | Rhamnaceae | E | Erica | Ericaceae | E |
| Diselma | Cupressaceae | G | Erigeron | Asteraceae | E |
| <i>Disperis</i> | Orchidaceae | M | Eriochilus | Orchidaceae | M |
| Disphyma | Aizoaceae | E | <i>Eriostemon</i> | Rutaceae | E |
| Distichlis | Poaceae | M | Erodium | Geraniaceae | E |
| Dittrichia | Asteraceae | E | Erophila | Brassicaceae | E |
| Diuris | Orchidaceae | M | <i>Eruca</i> | Brassicaceae | E |
| Dockrillia | Orchidaceae | M | Eryngium | Apiaceae | E |
| Dodonaea | Sapindaceae | E | <i>Erythraea</i> | Gentianaceae | E |
| <i>Dolichos</i> | Fabaceae | E | Erythranthe | Phrymaceae | E |
| Donatia | Donatiaceae | E | <i>Erythranthera</i> | Poaceae | M |
| <i>Doodia</i> | Blechnaceae | P | Eschscholzia | Papaveraceae | E |
| Draba | Brassicaceae | E | Eucalyptus | Myrtaceae | E |
| Dracophyllum | Ericaceae | E | Euchiton | Asteraceae | E |
| <i>Drapetes</i> | Thymelaeaceae | E | Eucryphia | Cunoniaceae | E |
| <i>Drimys</i> | Winteraceae | MG | Euonymus | Celastraceae | E |
| Drosanthemum | Aizoaceae | E | <i>Eupatorium</i> | Asteraceae | E |
| Drosera | Droseraceae | E | Euphorbia | Euphorbiaceae | E |
| Drymophila | Luzuriagaceae | M | Euphrasia | Orobanchaceae | E |
| Dryopoa | Poaceae | M | <i>Eurybia</i> | Asteraceae | E |
| Dysphania | Amaranthaceae | E | Eurychorda | Restionaceae | M |
| Ecballium | Cucurbitaceae | E | Euryomyrtus | Myrtaceae | E |

| | | | | | |
|--------------------|------------------|---|--------------------|-------------------|----|
| Euryops | Asteraceae | E | Glaucium | Papaveraceae | E |
| Eutaxia | Fabaceae | E | Gleichenia | Gleicheniaceae | P |
| Ewartia | Asteraceae | E | Glossodia | Orchidaceae | M |
| <i>Exacum</i> | Gentianaceae | E | Glossostigma | Phrymaceae | E |
| <i>Exarrhena</i> | Boraginaceae | E | Glyceria | Poaceae | M |
| Exocarpos | Santalaceae | E | Glycine | Fabaceae | E |
| <i>Fabricia</i> | Myrtaceae | E | Gnaphalium | Asteraceae | E |
| <i>Fagus</i> | Nothofagaceae | E | Gomphocarpus | Asclepidaceae | E |
| Fallopia | Polygonaceae | E | Gompholobium | Fabaceae | E |
| <i>Felicia</i> | Asteraceae | E | Gonocarpus | Haloragaceae | E |
| Festuca | Poaceae | M | Goodenia | Goodeniaceae | E |
| Ficinia | Cyperaceae | M | Goodia | Fabaceae | E |
| <i>Filago</i> | Asteraceae | E | <i>Grammitis</i> | Gleicheniaceae | P |
| <i>Fitzroya</i> | Cupressaceae | G | Gratiola | Plantaginaceae | E |
| Foeniculum | Apiaceae | E | Grevillea | Proteaceae | E |
| Forstera | Stylidiaceae | E | Griselinia | Griselinaceae | E |
| Frankenia | Frankeniaceae | E | Gunnera | Gunneraceae | E |
| Fraxinus | Oleaceae | E | <i>Gunnia</i> | Orchidaceae | M |
| Freesia | Iridaceae | M | <i>Gymnogramma</i> | Hemionitidiaceae | P |
| <i>Frenela</i> | Cupressaceae | G | Gymnoschoenus | Cyperaceae | M |
| Fuchsia | Onagraceae | E | Gynatrix | Malvaceae | E |
| Fumaria | Fumariaceae | E | Gypsophila | Caryophyllaceae | E |
| Furcraea | Asparagaceae | M | Gyrostemon | Gyrostemonaceae | E |
| <i>Fuscospora</i> | Nothofagaceae | E | Hackelia | Boraginaceae | E |
| Gahnia | Cyperaceae | M | Haemodorum | Haemodoraceae | M |
| Gaimardia | Centrolepidaceae | M | Hainardia | Poaceae | M |
| Galenia | Aizoaceae | E | Hakea | Proteaceae | E |
| Galinsoga | Asteraceae | E | Halophila | Hydrocharitaceae | M |
| Galium | Rubiaceae | E | Haloragis | Haloragaceae | E |
| Gamochaeta | Asteraceae | E | Hardenbergia | Fabaceae | E |
| Gastridium | Poaceae | M | <i>Hebe</i> | Plantaginaceae | E |
| Gastrodia | Orchidaceae | M | Hedera | Araliaceae | E |
| Gaudinia | Poaceae | M | Hedycarya | Monimiaceae | MG |
| Gaultheria | Ericaceae | E | <i>Hedybnois</i> | Asteraceae | E |
| Gazania | Asteraceae | E | <i>Hedysarum</i> | Fabaceae | E |
| Genista | Fabaceae | E | Helichrysum | Asteraceae | E |
| <i>Genoplesium</i> | Orchidaceae | M | <i>Helipterum</i> | Asteraceae | E |
| <i>Genosiris</i> | Iridaceae | M | <i>Helminthia</i> | Asteraceae | E |
| <i>Gentiana</i> | Gentianaceae | E | Helminthotheca | Asteraceae | E |
| Gentianella | Gentianaceae | E | <i>Helxine</i> | Urticaceae | E |
| Geococcus | Brassicaceae | E | Hemarthria | Poaceae | M |
| Geranium | Geraniaceae | E | Hemichroa | Amaranthaceae | E |
| Geum | Rosaceae | E | Herpolirion | Hemerocallidaceae | M |
| <i>Githago</i> | Caryophyllaceae | E | Hesperantha | Iridaceae | M |
| Gladiolus | Iridaceae | M | Hesperocyparis | Cupressaceae | G |

| | | | | | |
|---------------------|------------------|---|---------------------|------------------|---|
| Heterozostera | Zosteraceae | M | Ixia | Iridaceae | M |
| <i>Hewardia</i> | Iridaceae | M | <i>Ixiolaena</i> | Asteraceae | E |
| Hibbertia | Dilleniaceae | E | <i>Ixodia</i> | Asteraceae | E |
| <i>Hibiscus</i> | Malvaceae | E | <i>Jonesiopsis</i> | Orchidaceae | M |
| <i>Hieracium</i> | Asteraceae | E | Juncus | Juncaceae | M |
| Hierochloe | Poaceae | M | Kelleria | Thymelaeaceae | E |
| Hirschfeldia | Brassicaceae | E | Kennedia | Fabaceae | E |
| Histiopteris | Dennstaedtiaceae | P | Kickxia | Plantaginaceae | E |
| Holcus | Poaceae | M | Kniphofia | Asphodelaceae | M |
| <i>Homeria</i> | Iridaceae | M | Koeleria | Poaceae | M |
| Hookerochloa | Poaceae | M | <i>Kohlrauschia</i> | Caryophyllaceae | E |
| Hordeum | Poaceae | M | Kunzea | Myrtaceae | E |
| Hornungia | Brassicaceae | E | Laburnum | Fabaceae | E |
| Hovea | Fabaceae | E | Lachnagrostis | Poaceae | M |
| Humulus | Cannabaceae | E | Lactuca | Asteraceae | E |
| Huperzia | Lycopodiaceae | P | <i>Lagarosiphon</i> | Hydrocharitaceae | M |
| <i>Hutchinsia</i> | Brassicaceae | E | Lagarostrobos | Podocarpaceae | G |
| Hyalosperma | Asteraceae | E | Lagenophora | Asteraceae | E |
| <i>Hydatella</i> | Hydatellaceae | B | Lagurus | Poaceae | M |
| Hydrocotyle | Araliaceae | E | Lamium | Lamiaceae | E |
| Hydrorchis | Orchidaceae | M | <i>Lampocarya</i> | Cyperaceae | M |
| <i>Hylogyne</i> | Proteaceae | E | Lampranthus | Aizoaceae | E |
| <i>Hymenanthera</i> | Violaceae | E | Landoltia | Araceae | M |
| <i>Hymenochilus</i> | Orchidaceae | M | Lapsana | Asteraceae | E |
| <i>Hymenolobus</i> | Brassicaceae | E | Lasiopetalum | Malvaceae | E |
| Hymenophyllum | Hymenophyllaceae | P | Lasiospermum | Asteraceae | E |
| <i>Hyoscyamus</i> | Solanaceae | E | Lastreopsis | Dryopteridaceae | P |
| Hypericum | Hypericaceae | E | Lathyrus | Fabaceae | E |
| Hypochoeris | Asteraceae | E | Lavandula | Lamiaceae | E |
| Hypolaena | Restionaceae | M | <i>Lavatera</i> | Malvaceae | E |
| Hypolepis | Dennstaedtiaceae | P | Lawrencia | Malvaceae | E |
| Hypoxis | Hypoxidaceae | M | Laxmannia | Laxmanniaceae | M |
| Ilex | Aquifoliaceae | E | Leiocarpa | Asteraceae | E |
| Imperata | Poaceae | M | Leionema | Rutaceae | E |
| Indigofera | Fabaceae | E | Lemna | Araceae | M |
| <i>Inula</i> | Asteraceae | E | Leonotis | Lamiaceae | E |
| Ipomoea | Convolvulaceae | E | Leontodon | Asteraceae | E |
| Iris | Iridaceae | M | Lepidium | Brassicaceae | E |
| Isachne | Poaceae | M | Lepidosperma | Cyperaceae | M |
| Isoetes | Isoetaceae | P | Lepilaena | Potamogetonaceae | M |
| Isoetopsis | Asteraceae | E | Leptecophylla | Ericaceae | E |
| Isolepis | Cyperaceae | M | Leptinella | Asteraceae | E |
| Isophysis | Iridaceae | M | Leptocarpus | Restionaceae | M |
| Isopogon | Proteaceae | E | Leptoceras | Orchidaceae | M |
| Isotoma | Campanulaceae | E | <i>Leptocymus</i> | Fabaceae | E |

| | | | | | |
|---------------------|-----------------|---|---------------------|------------------|---|
| Leptomeria | Santalaceae | E | Lycopodium | Lycopodiaceae | P |
| Leptorhynchos | Asteraceae | E | <i>Lycopsis</i> | Boraginaceae | E |
| Leptospermum | Myrtaceae | E | Lycopus | Lamiaceae | E |
| <i>Lepturus</i> | Poaceae | M | <i>Lyonsia</i> | Apocynaceae | E |
| Lepyrodia | Restionaceae | M | Lyperanthus | Orchidaceae | M |
| Leucanthemum | Asteraceae | E | Lysimachia | Myrsinaceae | E |
| Leucochrysum | Asteraceae | E | Lythrum | Lythraceae | E |
| Leucojum | Amaryllidaceae | M | <i>Malaxis</i> | Orchidaceae | M |
| Leucophyta | Asteraceae | E | Malus | Rosaceae | E |
| Leucopogon | Ericaceae | E | Malva | Malvaceae | E |
| Levenhookia | Stylidiaceae | E | <i>Marianthus</i> | Pittosporaceae | E |
| Leycesteria | Caprifoliaceae | E | Marrubium | Lamiaceae | E |
| Leymus | Poaceae | M | Marsilea | Marsileaceae | P |
| Libertia | Iridaceae | M | Matricaria | Asteraceae | E |
| Ligustrum | Oleaceae | E | Matthiola | Brassicaceae | E |
| Lilaeopsis | Apiaceae | E | Maytenus | Celastraceae | E |
| <i>Limnanthemum</i> | Menyanthaceae | E | Mazus | Mazaceae | E |
| Limonium | Plumbaginaceae | E | <i>Mecodium</i> | Hymenophyllaceae | P |
| Limosella | Plantaginaceae | E | Medicago | Fabaceae | E |
| Linaria | Plantaginaceae | E | <i>Meionectes</i> | Haloragaceae | E |
| Lindsaea | Lindsaeaceae | P | Melaleuca | Myrtaceae | E |
| <i>Linguella</i> | Orchidaceae | M | <i>Melandrium</i> | Caryophyllaceae | E |
| Linum | Linaceae | E | Melicytus | Violaceae | E |
| Liparophyllum | Menyanthaceae | E | Melilotus | Fabaceae | E |
| Lissanthe | Ericaceae | E | Melissa | Lamiaceae | E |
| Lithospermum | Boraginaceae | E | Mentha | Lamiaceae | E |
| <i>Litobrochia</i> | Pteridaceae | P | Mesembryanthemum | Aizoaceae | E |
| Lobelia | Campanulaceae | E | <i>Mesomelaena</i> | Cyperaceae | M |
| Lobularia | Brassicaceae | E | Micranthemum | Picrodendraceae | E |
| Logfia | Asteraceae | E | Microcachrys | Podocarpaceae | G |
| Lolium | Poaceae | M | Microlaena | Poaceae | M |
| Lomandra | Asparagaceae | M | Microseris | Asteraceae | E |
| <i>Lomaria</i> | Blechnaceae | P | Microsorium | Polypodiaceae | P |
| Lomatia | Proteaceae | E | <i>Microstrobos</i> | Podocarpaceae | G |
| Lonicera | Caprifoliaceae | E | Microtidium | Orchidaceae | M |
| <i>Lophochloa</i> | Poaceae | M | Microtis | Orchidaceae | M |
| <i>Lophopyrum</i> | Poaceae | M | Milligania | Asteliaceae | M |
| <i>Lophozonia</i> | Nothofagaceae | E | Millotia | Asteraceae | E |
| Lotus | Fabaceae | E | <i>Mimosa</i> | Fabaceae | E |
| Lunaria | Brassicaceae | E | <i>Mimulus</i> | Phrymaceae | E |
| Lupinus | Fabaceae | E | Minuartia | Caryophyllaceae | E |
| Luzula | Juncaceae | M | Mirabilis | Nyctaginaceae | E |
| <i>Lychnis</i> | Caryophyllaceae | E | Mirbelia | Fabaceae | E |
| Lycium | Solanaceae | E | Misopates | Plantaginaceae | E |
| Lycopodiella | Lycopodiaceae | P | Mitrasacme | Loganiaceae | E |

| | | | | | |
|--------------------------|------------------|---|-----------------------|-----------------|---|
| <i>Mniarum</i> | Caryophyllaceae | E | Ononis | Fabaceae | E |
| Modiola | Malvaceae | E | Onopordum | Asteraceae | E |
| Moenchia | Caryophyllaceae | E | Opercularia | Rubiaceae | E |
| Molineriella | Poaceae | M | Ophioglossum | Ophioglossaceae | P |
| <i>Monerma</i> | Poaceae | M | Oreobolus | Cyperaceae | M |
| Monotoca | Ericaceae | E | Oreomyrrhis | Apiaceae | E |
| Montia | Portulacaceae | E | <i>Oreoporanthera</i> | Phyllanthaceae | E |
| Montitega | Ericaceae | E | Orites | Proteaceae | E |
| Moraea | Iridaceae | M | Ornduffia | Menyanthaceae | E |
| Muehlenbeckia | Polygonaceae | E | Ornithogalum | Hyacinthaceae | M |
| Muscari | Asparagaceae | M | Ornithopus | Fabaceae | E |
| Myoporum | Scrophulariaceae | E | Orobanche | Orobanchaceae | E |
| Myosotis | Boraginaceae | E | Orthoceras | Orchidaceae | M |
| Myosurus | Ranunculaceae | E | <i>Oryzopsis</i> | Poaceae | M |
| <i>Myriogyne</i> | Asteraceae | E | Oschatzia | Apiaceae | E |
| Myriophyllum | Haloragaceae | E | <i>Osteospermum</i> | Asteraceae | E |
| <i>Myrsiphyllum</i> | Asparagaceae | M | Ourisia | Plantaginaceae | E |
| Nablonium | Asteraceae | E | Oxalis | Oxalidaceae | E |
| <i>Nanozostera</i> | Zosteraceae | M | Oxylobium | Fabaceae | E |
| Narcissus | Amaryllidaceae | M | Ozothamnus | Asteraceae | E |
| Nardus | Poaceae | M | Pachycladon | Brassicaceae | E |
| Nassella | Poaceae | M | <i>Panax</i> | Araliaceae | E |
| Nasturtium | Brassicaceae | E | Pandorea | Bignoniaceae | E |
| Navarretia | Polemoniaceae | E | Panicum | Poaceae | M |
| <i>Nemacianthus</i> | Orchidaceae | M | Papaver | Papaveraceae | E |
| <i>Nematoceras</i> | Orchidaceae | M | Pappochroma | Asteraceae | E |
| Nematolepis | Rutaceae | E | <i>Paquerina</i> | Asteraceae | E |
| <i>Neopaxia</i> | Portulacaceae | E | Paracaleana | Orchidaceae | M |
| Nepeta | Lamiaceae | E | <i>Parahebe</i> | Plantaginaceae | E |
| <i>Nephrodium</i> | Dryopteridaceae | P | Parapholis | Poaceae | M |
| Nertera | Rubiaceae | E | Paraserianthes | Fabaceae | E |
| Nicotiana | Solanaceae | E | Parentucellia | Orobanchaceae | E |
| Nonea | Boraginaceae | E | Parietaria | Urticaceae | E |
| Notelaea | Oleaceae | E | Paronychia | Caryophyllaceae | E |
| Nothofagus | Nothofagaceae | E | Parsonsia | Apocynaceae | E |
| <i>Nothopanax</i> | Araliaceae | E | <i>Paspalidium</i> | Poaceae | M |
| Nothoscordum | Alliaceae | M | Paspalum | Poaceae | M |
| <i>Notodanthonia</i> | Poaceae | M | Passiflora | Passifloraceae | E |
| <i>Nymphoides</i> | Menyanthaceae | E | Pastinaca | Apiaceae | E |
| Odixia | Asteraceae | E | Patersonia | Iridaceae | M |
| Oenothera | Onagraceae | E | Pelargonium | Geraniaceae | E |
| Olearia | Asteraceae | E | Pellaea | Adiantaceae | P |
| <i>Oligochaetochilus</i> | Orchidaceae | M | <i>Pennisetum</i> | Poaceae | M |
| <i>Onobrychis</i> | Fabaceae | E | Pentachondra | Ericaceae | E |
| <i>Onoclea</i> | Blechnaceae | P | Pentaglottis | Boraginaceae | E |

| | | | | | |
|----------------------|-------------------|---|-------------------------|------------------|---|
| Pentapogon | Poaceae | M | Pneumatopteris | Thelypteridaceae | P |
| <i>Pernettya</i> | Ericaceae | E | Poa | Poaceae | M |
| Persicaria | Polygonaceae | E | Podalyria | Fabaceae | E |
| Persoonia | Proteaceae | E | Podocarpus | Podocarpaceae | G |
| <i>Petalochilus</i> | Orchidaceae | M | Podolepis | Asteraceae | E |
| Petasites | Asteraceae | E | <i>Podosperma</i> | Asteraceae | E |
| Petrorhagia | Caryophyllaceae | E | Podotheca | Asteraceae | E |
| Petroselinum | Apiaceae | E | Polycarpon | Caryophyllaceae | E |
| Phacelia | Boraginaceae | E | Polygala | Polygalaceae | E |
| Phalaris | Poaceae | M | Polygonum | Polygonaceae | E |
| Phebalium | Rutaceae | E | Polyphlebium | Hymenophyllaceae | P |
| Pheladenia | Orchidaceae | M | <i>Polypodium</i> | Polypodiaceae | P |
| Pherosphaera | Podocarpaceae | G | <i>Polypodium</i> | Grammitidaceae | P |
| Philotheca | Rutaceae | E | Polypogon | Poaceae | M |
| Phlegmariurus | Lycopodiaceae | P | <i>Polypompholyx</i> | Lentibulariaceae | E |
| Phleum | Poaceae | M | Polyscias | Araliaceae | E |
| Phormium | Hemerocallidaceae | M | Polystichum | Dryopteridaceae | P |
| Phragmites | Poaceae | M | Pomaderris | Rhamnaceae | E |
| Phyllachne | Stylidiaceae | E | Populus | Salicaceae | E |
| Phyllangium | Loganiaceae | E | Poranthera | Phyllanthaceae | E |
| Phyllanthus | Phyllanthaceae | E | Portulaca | Portulacaceae | E |
| Phyllocladus | Podocarpaceae | G | Posidonia | Posidoniaceae | M |
| Phylloglossum | Lycopodiaceae | P | Potamogeton | Potamogetonaceae | M |
| Phyllota | Fabaceae | E | Potentilla | Rosaceae | E |
| <i>Phymatodes</i> | Polypodiaceae | P | <i>Poterium</i> | Rosaceae | E |
| <i>Phymatosorus</i> | Polypodiaceae | P | Prasophyllum | Orchidaceae | M |
| Physalis | Solanaceae | E | <i>Pratia</i> | Campanulaceae | E |
| Phytolacca | Phytolaccaceae | E | Prionotes | Ericaceae | E |
| Picris | Asteraceae | E | Prostanthera | Lamiaceae | E |
| <i>Pilitis</i> | Ericaceae | E | Prunella | Lamiaceae | E |
| Pilosella | Asteraceae | E | Prunus | Rosaceae | E |
| Pilularia | Marsileaceae | P | <i>Psamma</i> | Poaceae | M |
| Pimelea | Thymelaeaceae | E | Pseudanthus | Picrodendraceae | E |
| Pinus | Pinaceae | G | <i>Pseudofumaria</i> | Fumariaceae | E |
| Piptatherum | Poaceae | M | <i>Pseudognaphalium</i> | Asteraceae | E |
| Pittosporum | Pittosporaceae | E | Pseudopanax | Araliaceae | E |
| <i>Plagianthus</i> | Malvaceae | E | <i>Psilotum</i> | Psilotaceae | P |
| Planocarpa | Ericaceae | E | Psoralea | Fabaceae | E |
| Plantago | Plantaginaceae | E | Psychrophila | Ranunculaceae | E |
| Platylobium | Fabaceae | E | Pteridium | Dennstaedtiaceae | P |
| <i>Platyloma</i> | Adiantaceae | P | Pteris | Pteridaceae | P |
| <i>Pleurandra</i> | Dilleniaceae | E | Pterostylis | Orchidaceae | M |
| Pleurophyllum | Asteraceae | E | Pterygopappus | Asteraceae | E |
| Pleurosorus | Aspleniaceae | P | Ptilotus | Amaranthaceae | E |
| <i>Plumatichilos</i> | Orchidaceae | M | Puccinellia | Poaceae | M |

| | | | | | |
|--------------------|-----------------|---|----------------------|------------------|---|
| Pultenaea | Fabaceae | E | <i>Sarothamnus</i> | Fabaceae | E |
| <i>Pygmea</i> | Plantaginaceae | E | Saxipoa | Poaceae | M |
| Pyrorchis | Orchidaceae | M | Scabiosa | Dipsacaceae | E |
| <i>Racosperma</i> | Fabaceae | E | Scaevola | Goodeniaceae | E |
| Ranunculus | Ranunculaceae | E | Scandix | Apiaceae | E |
| <i>Raoulia</i> | Asteraceae | E | <i>Schedonorus</i> | Poaceae | M |
| Raphanus | Brassicaceae | E | Schenkia | Gentianaceae | E |
| Rapistrum | Brassicaceae | E | Schizacme | Loganiaceae | E |
| Reseda | Resedaceae | E | Schizaea | Schizaeaceae | P |
| <i>Restio</i> | Restionaceae | M | <i>Schizostylis</i> | Iridaceae | M |
| <i>Reynoutria</i> | Polygonaceae | E | <i>Schoenodum</i> | Restionaceae | M |
| <i>Rhagadiolus</i> | Asteraceae | E | <i>Schoenoides</i> | Cyperaceae | M |
| Rhagodia | Amaranthaceae | E | Schoenoplectus | Cyperaceae | M |
| Rhamnus | Rhamnaceae | E | Schoenus | Cyperaceae | M |
| Rhodanthe | Asteraceae | E | Scilla | Hyacinthaceae | M |
| Rhododendron | Ericaceae | E | <i>Scirpus</i> | Cyperaceae | M |
| Rhytidosporum | Pittosporaceae | E | Scleranthus | Caryophyllaceae | E |
| Ribes | Grossulariaceae | E | Sclerochloa | Poaceae | M |
| <i>Richea</i> | Asteraceae | E | <i>Scleroleima</i> | Asteraceae | E |
| Richea | Ericaceae | E | <i>Sclerostegia</i> | Amaranthaceae | E |
| Ricinocarpos | Euphorbiaceae | E | Scorzonera | Asteraceae | E |
| <i>Roepera</i> | Zygophyllaceae | E | Scrophularia | Scrophulariaceae | E |
| Romulea | Iridaceae | M | Scutellaria | Lamiaceae | E |
| Rorippa | Brassicaceae | E | Sebaea | Gentianaceae | E |
| Rosa | Rosaceae | E | Secale | Poaceae | M |
| Rostraria | Poaceae | M | Securigera | Fabaceae | E |
| Rubus | Rosaceae | E | Sedum | Crassulaceae | E |
| Rumex | Polygonaceae | E | Selaginella | Selaginellaceae | P |
| Rumohra | Dryopteridaceae | P | Selliera | Goodeniaceae | E |
| Ruppia | Ruppiaceae | M | <i>Senebiera</i> | Brassicaceae | E |
| <i>Rutidosis</i> | Asteraceae | E | Senecio | Asteraceae | E |
| Rytidosperma | Poaceae | M | Setaria | Poaceae | M |
| Sagina | Caryophyllaceae | E | <i>Sheffieldia</i> | Theophrastaceae | E |
| Sagittaria | Alismataceae | M | Sherardia | Rubiaceae | E |
| <i>Salicornia</i> | Amaranthaceae | E | Sicyos | Cucurbitaceae | E |
| Salix | Salicaceae | E | <i>Sieglingia</i> | Poaceae | M |
| Salpichroa | Solanaceae | E | Sigesbeckia | Asteraceae | E |
| Salsola | Amaranthaceae | E | Silene | Caryophyllaceae | E |
| Salvia | Lamiaceae | E | Siloxerus | Asteraceae | E |
| Sambucus | Adoxaceae | E | Silybum | Asteraceae | E |
| Samolus | Theophrastaceae | E | <i>Simpliglottis</i> | Orchidaceae | M |
| Sanguisorba | Rosaceae | E | Sinapis | Brassicaceae | E |
| Saponaria | Caryophyllaceae | E | Sisymbrium | Brassicaceae | E |
| Sarcochilus | Orchidaceae | M | Sisyrinchium | Iridaceae | M |
| Sarcocornia | Amaranthaceae | E | <i>Skirrhophorus</i> | Asteraceae | E |

| | | | | | |
|-----------------------|-------------------|---|--------------------|-------------------|----|
| Solanum | Solanaceae | E | Symphotrichum | Asteraceae | E |
| Soleirolia | Urticaceae | E | Symphytum | Boraginaceae | E |
| Solenogyne | Asteraceae | E | Syzygium | Myrtaceae | E |
| Soliva | Asteraceae | E | Tanacetum | Asteraceae | E |
| <i>Sollya</i> | Pittosporaceae | E | Taraxacum | Asteraceae | E |
| Sonchus | Asteraceae | E | Tasmania | Winteraceae | MG |
| Sorbus | Rosaceae | E | <i>Taurantha</i> | Orchidaceae | M |
| Sorghum | Poaceae | M | <i>Taxanthemea</i> | Plumbaginaceae | E |
| Sowerbaea | Laxmanniaceae | M | Tecticornia | Amaranthaceae | E |
| Sparaxis | Iridaceae | M | Teesdalia | Brassicaceae | E |
| Spartina | Poaceae | M | Telopea | Proteaceae | E |
| Spartium | Fabaceae | E | Tetracarpaea | Tetracarpaeaceae | E |
| <i>Speculantha</i> | Orchidaceae | M | Tetragonia | Aizoaceae | E |
| Spergula | Caryophyllaceae | E | Tetralia | Cyperaceae | M |
| Spergularia | Caryophyllaceae | E | Tetrarrhena | Poaceae | M |
| <i>Sphaerocionium</i> | Hymenophyllaceae | P | Tetratheca | Elaeocarpaceae | E |
| Sphaerolobium | Fabaceae | E | Teucrium | Lamiaceae | E |
| <i>Spiculaea</i> | Orchidaceae | M | Thelionema | Hemerocallidaceae | M |
| Spinifex | Poaceae | M | Thelymitra | Orchidaceae | M |
| Spiranthes | Orchidaceae | M | Themeda | Poaceae | M |
| <i>Spirodela</i> | Araceae | M | Thesium | Santalaceae | E |
| Sporadanthus | Restionaceae | M | Thinopyrum | Poaceae | M |
| Sporobolus | Poaceae | M | Thismia | Burmanniaceae | M |
| Sprengelia | Ericaceae | E | Thlaspi | Brassicaceae | E |
| Spyridium | Rhamnaceae | E | <i>Thonandia</i> | Poaceae | M |
| Stachys | Lamiaceae | E | Threlkeldia | Amaranthaceae | E |
| Stackhousia | Celastraceae | E | Thryptomene | Myrtaceae | E |
| <i>Statice</i> | Plumbaginaceae | E | Thynninorchis | Orchidaceae | M |
| <i>Stegania</i> | Blechnaceae | P | Thyridia | Phrymaceae | E |
| <i>Stegostyla</i> | Orchidaceae | M | Thysanotus | Laxmanniaceae | M |
| Stellaria | Caryophyllaceae | E | <i>Tillaea</i> | Crassulaceae | E |
| Stenanthemum | Rhamnaceae | E | Tmesipteris | Psilotaceae | P |
| <i>Stenanthera</i> | Ericaceae | E | Todea | Osmundaceae | P |
| Stenopetalum | Brassicaceae | E | Tolpis | Asteraceae | E |
| Stenotaphrum | Poaceae | M | Torilis | Apiaceae | E |
| Sticherus | Gleicheniaceae | P | Townsonia | Orchidaceae | M |
| <i>Stipa</i> | Poaceae | M | Trachymene | Araliaceae | E |
| Stonesiella | Fabaceae | E | Tradescantia | Commelinaceae | M |
| Stuckenia | Potamogetonaceae | M | Tragopogon | Asteraceae | E |
| Stylidium | Stylidiaceae | E | <i>Trichinium</i> | Amaranthaceae | E |
| <i>Stypandra</i> | Hemerocallidaceae | M | <i>Trichomanes</i> | Hymenophyllaceae | P |
| Styphelia | Ericaceae | E | <i>Trichonema</i> | Iridaceae | M |
| Suaeda | Amaranthaceae | E | Tricoryne | Hemerocallidaceae | M |
| <i>Sullivania</i> | Orchidaceae | M | Tricostularia | Cyperaceae | M |
| Swainsona | Fabaceae | E | Trifolium | Fabaceae | E |

| | | | | | |
|-------------------|------------------|---|---------------------|------------------|---|
| Triglochin | Juncaginaceae | M | Viola | Violaceae | E |
| <i>Trigonella</i> | Fabaceae | E | Vittadinia | Asteraceae | E |
| <i>Triodia</i> | Poaceae | M | Vulpia | Poaceae | M |
| Tripleurospermum | Asteraceae | E | Wahlenbergia | Campanulaceae | E |
| Triptilodiscus | Asteraceae | E | Watsonia | Iridaceae | M |
| Trisetum | Poaceae | M | Westringia | Lamiaceae | E |
| Trithuria | Hydatellaceae | B | Wilsonia | Convolvulaceae | E |
| Triticum | Poaceae | M | Winifredia | Restionaceae | M |
| Tritonia | Iridaceae | M | Wolffia | Araceae | M |
| Trochocarpa | Ericaceae | E | Wurmbea | Colchicaceae | M |
| Tropaeolum | Tropaeolaceae | E | Xanthium | Asteraceae | E |
| Typha | Typhaceae | M | Xanthorrhoea | Xanthorrhoeaceae | M |
| Ulex | Fabaceae | E | Xanthosia | Apiaceae | E |
| Ulmus | Ulmaceae | E | Xerochrysum | Asteraceae | E |
| Uncinia | Cyperaceae | M | <i>Xerotes</i> | Asparagaceae | M |
| <i>Uniola</i> | Poaceae | M | Xyris | Xyridaceae | M |
| <i>Urochilus</i> | Orchidaceae | M | <i>Zaleya</i> | Aizoaceae | E |
| Urospermum | Asteraceae | E | <i>Zannichellia</i> | Potamogetonaceae | M |
| Urtica | Urticaceae | E | Zantedeschia | Araceae | M |
| Utricularia | Lentibulariaceae | E | Zieria | Rutaceae | E |
| <i>Vaccaria</i> | Caryophyllaceae | E | Zostera | Zosteraceae | M |
| Valerianella | Valerianaceae | E | Zoysia | Poaceae | M |
| Vallisneria | Hydrocharitaceae | M | Zygophyllum | Zygophyllaceae | E |
| <i>Valvaria</i> | Ranunculaceae | E | <i>×Agropogon</i> | Poaceae | M |
| Velleia | Goodeniaceae | E | <i>×Calassodia</i> | Orchidaceae | M |
| Vellereophyton | Asteraceae | E | <i>×Festulolium</i> | Poaceae | M |
| Verbascum | Scrophulariaceae | E | <i>×Glossadenia</i> | Orchidaceae | M |
| Verbena | Verbenaceae | E | <i>×Taurodium</i> | Orchidaceae | M |
| Veronica | Plantaginaceae | E | | | |
| Viburnum | Adoxaceae | E | | | |
| Vicia | Fabaceae | E | | | |
| <i>Villarsia</i> | Menyanthaceae | E | | | |
| Viminaria | Fabaceae | E | | | |
| Vinca | Apocynaceae | E | | | |

REFERENCES

- ABRS (Australian Biological Resources Study) (Australian Government: Department of the Environment, Water, Heritage and the Arts). <http://www.environment.gov.au/biodiversity/abrs/index.html>
- APG II (2003) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Botanical Journal of the Linnean Society* 141: 399–436.
- Baker ML, de Salas MF (2012) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania & Flora of Tasmania Online* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- Baker ML, de Salas MF (2013) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania & Flora of Tasmania Online* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- Baker ML, Duretto MF (2011) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania & Flora of Tasmania Online* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- Bentham G (1863) *Flora Australiensis*. Vol. I. (Lovell Reeve and Co.: London)
- Bentham G (1864) *Flora Australiensis*. Vol. II (Lovell Reeve and Co.: London)
- Bentham G (1867) *Flora Australiensis*. Vol. III (Lovell Reeve and Co.: London)
- Bentham G (1868) *Flora Australiensis*. Vol. IV (Lovell Reeve and Co.: London)
- Bentham G (1870) *Flora Australiensis*. Vol. V (Lovell Reeve and Co.: London)
- Bentham G (1873) *Flora Australiensis*. Vol. VI (Lovell Reeve and Co.: London)
- Bentham G (1878) *Flora Australiensis*. Vol. VII (Lovell Reeve and Co.: London)
- Brown R (1810) *Prodromus Florae Novae Hollandiae et Insulae van-Diemen* (Richard Taylor & Son: London)
- Buchanan AM (1995) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania*. Tasmanian Herbarium Occasional Publication No. 5 (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart)
- Buchanan AM (1999) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania*. 3rd edn. Tasmanian Herbarium Occasional Publication No. 6 (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart)
- Buchanan AM (2004) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- Buchanan AM (2005) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania*. 4th edn. Tasmanian Herbarium Occasional Publication No. 7 (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart)
- Buchanan AM (2007) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- Buchanan AM (2009) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- Buchanan AM, McGeary-Brown A, Orchard AE (1989) *A Census of the Vascular Plants of Tasmania*. Tasmanian Herbarium Occasional Publication No. 2 (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart)

- Burbidge, NT (1958) A monographic study of *Helichrysum* subgenus *Ozothamnus* (Compositae) and of two related genera formerly included therein. *Australian Journal of Botany* 6: 229–284
- Cheeseman TF (1919) The vascular flora of Macquarie Island. *Australasian Antarctic Expedition 1911–1914. Scientific Reports. Series C. Zoology and Botany. Volume VII, Part 3.* 63 pp. (Government Printer: Sydney)
- Cronquist AJ (1981) *An Integrated System of Classification of Flowering Plants.* (Columbia University Press: New York)
- Curtis (1956) *The Student's Flora of Tasmania Part I – Gymnospermae; Angiospermae: Ranunculaceae to Myrtaceae.* 1st edn. (Government Printer: Hobart)
- Curtis WM (1963) *The Student's Flora of Tasmania Part 2 – Angiospermae: Lythraceae to Epacridaceae.* (Government Printer: Hobart)
- Curtis WM (1967) *The Student's Flora of Tasmania Part 3 – Angiospermae: Plumbaginaceae to Salicaceae.* (Government Printer: Hobart)
- Curtis WM (1979) *The Student's Flora of Tasmania Part 4A – Angiospermae: Orchidaceae.* (Government Printer: Hobart)
- Curtis WM, Morris DI (1975) *The Student's Flora of Tasmania Part I – Gymnospermae; Angiospermae: Ranunculaceae to Myrtaceae.* 2nd edn. (Government Printer: Hobart)
- Curtis WM, Morris DI (1994) *The Student's Flora of Tasmania Part 4B – Angiospermae: Alismataceae to Burmanniaceae.* (St. Davids Park Publishing: Hobart)
- de Salas MF (2009+) *Flora of Tasmania Online* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart). <https://flora.tmag.tas.gov.au>
- de Salas MF, Baker ML (2014) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania & Flora of Tasmania Online* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- de Salas MF, Baker ML (2015) *A Census of the Vascular Plants of Tasmania including Macquarie Island* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- de Salas MF, Baker ML (2016) *A Census of the Vascular Plants of Tasmania including Macquarie Island* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- de Salas MF, Baker ML (2017) *A Census of the Vascular Plants of Tasmania including Macquarie Island* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- de Salas MF, Baker ML (2018) *A Census of the Vascular Plants of Tasmania including Macquarie Island* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) www.tmag.tas.gov.au
- Garrett M (1996) *The Ferns of Tasmania; Their Ecology and Distribution* (Tasmanian Forest Research Council: Hobart)
- Harris S, Buchanan A, Connolly A (2001) *One Hundred Islands: The Flora of the Outer Fureaux.* (Tasmanian DPIWE: Hobart). p. 197
- Henderson RJF in George AS (Ed) (1987) *Flora of Australia* 45: 209–214, 477–479 (ABRS: Canberra)
- Hewson HJ in George AS (Ed) (1982) *Flora of Australia* 8: 313, map 366 (ABRS: Canberra)
- Hooker JD (1844) *The Botany of the Antarctic Voyage of H. M. Discovery Ships Erebus and Terror in the Years 1839 – 1843 Under the Command of Captain Sir James Clark Ross. Part I: Flora Antarctica* (Reeve Brothers: London)

- Hooker JD (1847) Contributions towards a Flora of Van Diemen's Land. *London Journal of Botany* 6: 461 (bis)–479 (bis)
- Hooker JD (1860) *The Botany of the Antarctic Voyage of H. M. Discovery Ships Erebus and Terror in the Years 1839 – 1843 Under the Command of Captain Sir James Clark Ross. Part III: Flora Tasmaniae* (Reeve Brothers: London)
- Jarman SJ and Kantvilas GK (2017) *Leptecophylla* in Tasmania: a reassessment of four species. *Swainsona* 31: 1–16
- Labillardiere JJH de (1804–1807) *Novae Hollandiae Plantarum Specimen* (Typographia Dominae Huzard: Paris)
- Maiden JH, Betche E (1916) *A Census of New South Wales Plants* (W. A. Gullick, Government Printer: Sydney)
- McCarthy PM (1998) Ferns, Gymnosperms and Allied Groups. *Flora of Australia* 48. (ABRS: Canberra)
- Morris DI (1977) Miscellaneous notes on endemic Tasmanian plants in the genera *Olearia*, *Ixodia*, *Danthonia*, *Tetrarrhena*. *Records of the Queen Victoria Museum* 55: 1–5
- Raphael TD (1955) Tasmanian garden escapes. *Papers and Proceedings of the Royal Society of Tasmania* 89: 147–150
- Rodway L (1903) *The Tasmanian Flora*. (Government Printer: Hobart)
- Rye BL (1990) Thymelaeaceae. *Flora of Australia* 18: 181, Map 247 (ABRS: Canberra)
- Smith JJ (1912) Epacridaceae. *Nova Guinea* 8: 797
- Townrow JES (1969) A species list of and keys to the grasses of Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* 103: 69–96
- Townrow J (1978) *Papers and Proceedings of the Royal Society of Tasmania* 112
- Willis JH (1970) *A Handbook to Plants in Victoria*. (Melbourne Uni. Press: Melbourne)
- Willis JH (1973) *A Handbook to Plants of Victoria. Vol. 1 – Ferns, Conifers and Monocotyledons*. Ed. 2 (Melbourne Uni. Press: Melbourne)